

DESTABILISING COMPACT WARPED PRODUCT EINSTEIN MANIFOLDS

WAFAA BATAT, STUART HALL, AND THOMAS MURPHY

ABSTRACT. The linear stability of warped product Einstein metrics as fixed points of the Ricci flow is investigated. We generalise the results of Gibbons, Hartnoll and Pope and show that in sufficiently low dimensions, all warped product Einstein metrics are unstable. By exploiting the relationship between warped product Einstein metrics, quasi-Einstein metrics and Ricci solitons, we introduce a new destabilising perturbation (the Ricci variation) and show that certain infinite families of warped product Einstein metrics will be unstable in high dimensions.

1. INTRODUCTION

1.1. Main results. In 2003, Perelman made spectacular use of Hamilton's Ricci flow to prove Thurston's geometrization conjecture [30], [31] and [32]. Put simply, geometrization says that three-dimensional manifolds decompose into pieces that can each be endowed with a canonical geometry. In order to extend geometrization to higher dimensions, a crucial step is finding the right set of canonical geometries in each dimension. One candidate for these geometries are metrics that are the *stable* fixed points (up to diffeomorphism and scaling) of the Ricci flow. Fixed points of the Ricci flow are known as *Ricci solitons* and clearly include Einstein metrics. Roughly speaking, stability can be taken to mean that the Ricci flow starting at small perturbation of a Ricci soliton will return to the soliton.

The study of the linear stability of Ricci solitons was initiated by Cao, Hamilton, and Ilmanen [6] who considered the second variation of Perelman's ν entropy, a monotonic quantity for the flow. In the subsequent years, stability questions for many important classes of metric have been investigated such as: Einstein metrics admitting parallel spinors [12], compact symmetric spaces [7], and Kähler metrics [18].

In this article we take up the study of stability for a class of compact Einstein metrics known as warped products. These are generalisations of ordinary Riemannian products where the underlying manifold M decomposes as $M = B \times F$ for a base B and a fibre F but the metric on the fibre is 'warped' by a factor coming from the base. We refer the reader to Section 2 for the precise definition of these metrics. There are many important examples of such Einstein metrics in low dimensions including the inhomogenous families on $\mathbb{S}^3 \times \mathbb{S}^2$, $\mathbb{S}^3 \times \mathbb{S}^3$, and $\mathbb{S}^4 \times \mathbb{S}^2$ due to Böhm [3], and the warped product Einstein metric on $\mathbb{C}\mathbb{P}^2 \sharp \overline{\mathbb{C}\mathbb{P}^2} \times \mathbb{S}^2$ due to Lü, Page and Pope [27]. We completely settle the question of stability in low dimensions:

Theorem A. *Let (M^n, g) be a warped product Einstein manifold where $n \leq 6$. Then (M, g) is unstable as a fixed point of the Ricci flow.*

We remark that the proof of Theorem A also shows that warped products with three-dimensional base and a four-dimensional fibre are unstable too. However, this does not

account for all possible seven-dimensional products.

Many examples of Einstein warped products are constructed from families of Riemannian manifolds (B, \bar{g}_i, f_i, m_i) where the metrics \bar{g}_i (known as *quasi-Einstein metrics*) converge in the C^∞ topology as $i \rightarrow \infty$ to a Ricci soliton $(B, \bar{g}_\infty, \phi)$. For example, Case has demonstrated that the metrics of Lü–Page–Pope have this property [10]. For warped products coming from such a family, we prove an asymptotic instability result that shows that the Einstein metrics are also unstable if the dimension is large enough.

Theorem B. *Let (B, \bar{g}_i, f_i, m_i) be a sequence of quasi-Einstein metrics that converge in the C^∞ topology to a non-trivial Ricci soliton $(B, \bar{g}_\infty, \phi)$.*

Then there exists a $K \in \mathbb{N}$ such that the associated warped product Einstein metrics (M^k, g) are unstable for $k \geq K$.

As discussed in Section 2, the construction of a warped product Einstein metric requires an Einstein metric \tilde{g} on the fibre F . The following theorem makes precise the interaction between the stability of the fibre Einstein metric and that of the warped product.

Theorem C. *Let (M, g) be a warped product Einstein manifold with fibre (F, \tilde{g}) satisfying $\text{Ric}(\tilde{g}) = \mu\tilde{g}$. Let σ be a divergence-free, trace-free eigentensor of the Lichnerowicz Laplacian of (F, \tilde{g}) satisfying $\tilde{\Delta}_L \sigma = -\kappa\sigma$. If $\kappa < \mu$ then the warped product (M, g) is unstable.*

This theorem allows us to find a large class of Ricci flow unstable warped products. We will refer to warped products that are unstable in the manner of Theorem C as fibre-unstable.

Corollary D. *The following fibres yield fibre-unstable warped products:*

- when (F, \tilde{g}) is a Riemannian product $(F_1 \times F_2, g_1 \oplus g_2)$,
- when (F, \tilde{g}) is a Kähler-Einstein metric with $h^{(1,1)} > 1$,
- when (F, \tilde{g}) is a fibre-unstable warped product Einstein metric.

1.2. The stability of generalised black holes. A second motivation for studying stability, which historically precedes the Ricci flow, comes from the role compact Einstein metrics play in the theory of generalised black holes. This was the context of the pioneering study of the stability of Böhm’s Einstein metrics on low-dimensional products of spheres conducted by Gibbons, Hartnoll, and Pope [15]. In their study they exploited the fact that the Böhm metrics are invariant under a cohomogeneity one action by a compact Lie group. We generalise their results in low dimensions to arbitrary warped product Einstein metrics without any symmetry assumptions.

Theorem E. *Let M be a warped product Einstein metric with three-dimensional base and two or three-dimensional fibre. Then the associated Schwarzschild–Tangherlini black hole is unstable.*

1.3. Relation of Theorem E to other works on stability. The construction of Schwarzschild–Tangherlini black holes from Einstein metrics with positive Einstein constant can itself be viewed as a non-compact warped product. There are many other methods that build geometrically interesting manifolds from positive Einstein metrics. For example, in [16] a stability inequality for Ricci-flat cones over Einstein bases was investigated. In this setting, as in Theorem E, the stability of the cone is related to the spectral properties of the Lichnerowicz Laplacian of the base. The authors proved that the cones over product manifolds in dimension less than 10 are unstable (which

is somewhat similar to the situation in Theorem E in that the construction over low dimensional ‘products’ is unstable). In [16] authors also discussed the conjectural link, due to Ilmanen, between unstable cones and the non-uniqueness of Ricci flows emerging from them. Kröncke has made a more extensive study of such non-compact warped product constructions in [24] and [25].

The relation between the linear stability discussed in this article and the dynamic stability of the Ricci flow has been made precise by Sesum in [33]. We refer the reader to Section 2.2 for more details on this relationship.

1.4. Conventions. As the proofs of the theorems rest on certain quantities having a particular sign, it is important to state exactly the conventions used in the paper where there is room for ambiguity. On the product $B \times F$, uppercase letters denote general coordinates, lowercase Roman letters denote the coordinates on B and lowercase Greek letters denote the coordinates on the fibre F . We will denote the associated coordinate vector fields by ∂_A , ∂_a and ∂_α . In order to try and keep coordinate calculations uncluttered, we will often abuse notation by writing $\nabla_A T$ for $\nabla_{\partial_A} T$ and $\nabla_A B$ for $\nabla_{\partial_A} B$.

The divergence of a tensor T is given by

$$\operatorname{div}(T)(\cdot) = g^{AB}(\nabla_A T)(\partial_B, \cdot)$$

and the (connection) Laplacian ΔT is given by

$$\Delta T = -\nabla^* \nabla T = g^{AB}(\nabla_{A,B}^2 T) = \operatorname{div}(\nabla T).$$

With this convention, the spectrum of the Laplacian is non-positive. The convention we use for curvature is

$$R(X, Y, Z, W) = g(R(X, Y)Z, W) = g(\nabla_{Y,X}^2 Z - \nabla_{X,Y}^2 Z, W),$$

and the curvature operator $\operatorname{Rm} : s^2(TM) \rightarrow s^2(TM)$ is given by

$$\operatorname{Rm}(h, \cdot)_{AB} = R_{ACBD} h^{CD},$$

for $h \in s^2(TM)$.

Geometric objects on the base manifold B will usually be denoted using a bar, for example \bar{g} for the metric. Geometric objects on the fibre F are likewise denoted using a tilde, so we have \tilde{g} for the fibre metric. We will set $n = \dim(B)$ and $m = \dim(F)$.

Acknowledgements: The initial work on this project was conducted whilst WB and TM were visiting SH in Dec 2015 and Jan 2016. The visit of WB was funded by an LMS grant under scheme 4. TM was funded by a Cal. State Fullerton Startup Grant. We would like to thank the referees for their careful reading and very useful comments

2. BACKGROUND

2.1. Warped Product Einstein metrics. Let $M = B \times F$ be a product manifold. Equip M with the metric

$$g = \pi_B^* \bar{g} \oplus (f \circ \pi_B)^2 \pi_F^* \tilde{g},$$

where \bar{g} and \tilde{g} are Riemannian metrics on B and F respectively, $\pi_B : M \rightarrow B$ and $\pi_F : M \rightarrow F$ denote the natural projections, and $f \in C^\infty(B)$. The Riemannian manifold (M, g) is referred to as a *warped product*. We shall henceforth adopt the standard

abuse of notation and drop the references to the projections π_B and π_F . The manifold B is referred to as the base and F is referred to as the fibre. By taking the function f to be constant, one recovers a Riemannian product.

We will be concerned with the case where the warped product is an Einstein metric with positive scalar curvature. By Myers's theorem this immediately implies the manifolds B and F are compact. If a warped product (M, g) is an Einstein manifold with $\text{Ric}(g) = \lambda g$ for $\lambda > 0$, then the following is well known (e.g. Corollary 9.107 in [2]) :

(F, \tilde{g}) is an Einstein manifold with $\text{Ric}(\tilde{g}) = \mu \tilde{g}$ for some $\mu > 0$,

$$f\bar{\Delta}f + (m-1)|\bar{\nabla}f|^2 + \lambda f^2 = \mu, \quad (2.1)$$

$$\text{Ric}(\bar{g}) - mf^{-1}\bar{\nabla}^2 f = \lambda \bar{g}. \quad (2.2)$$

Riemannian manifolds (B, \bar{g}) that solve Equation (2.2) for some $f \in C^\infty(B)$ and $m > 0$ are known as *quasi-Einstein* manifolds and are studied in their own right as interesting generalisations of Einstein metrics. A foundational result of Kim and Kim [22] says that if (B, \bar{g}, f, m) solve (2.2), then there exists a $\mu > 0$ such that f solves Equation (2.1). Hence for integral values of $m \geq 2$, one can construct warped product Einstein manifolds from a quasi-Einstein metric on the base B .

We also consider Riemannian manifolds (M, g) where the metric g solves

$$\text{Ric}(g) + \nabla^2 \phi = \lambda g, \quad (2.3)$$

where $\phi \in C^\infty(M)$ and $\lambda \in \mathbb{R}$. Metrics solving (2.3) are called *gradient Ricci solitons*. One can view Equation (2.3) as the formal limit as $m \rightarrow \infty$ of (2.2) by setting $\phi_m = -m \log f$. More detailed results on the sense in which gradient Ricci solitons are the limits of quasi-Einstein metrics can be found in the work of Case [11].

At the time of writing, there are very few general methods for constructing of compact, warped product Einstein metrics. Much more is known about constructions of non-compact warped product Einstein metrics, see for example Chapter 9 of [2] where there are examples of quasi-Einstein metrics on non-compact surfaces with the parameter $m \in (1, \infty)$. As mentioned in Section 1, the first compact examples that were found are due to Böhm [3] and occur on the product $\mathbb{S}^n \times F^m$ with $2 \leq m \leq 6$ and $3 \leq n \leq 9 - m$. The second family of examples come from a construction due to Lü, Page and Pope [27] of quasi-Einstein metrics on $\mathbb{C}\mathbb{P}^1$ -bundles over a Fano Kähler-Einstein base. This construction (and its generalisation due to the second author [17]) produce quasi-Einstein metrics for all $m > 1$ and hence infinitely many examples of warped product Einstein manifolds. The lowest dimensional examples of the Lü-Page-Pope construction occur when the base is the non-trivial $\mathbb{C}\mathbb{P}^1$ -bundle over $\mathbb{C}\mathbb{P}^1$. In this case one can view the base as $B = \mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2}$. A very explicit construction of the Lü-Page-Pope metrics on this manifold was given in [1]. As mentioned already, Case [10] showed that the Lü-Page-Pope quasi-Einstein metrics converge as $m \rightarrow \infty$ to the Koiso-Cao Kähler-Ricci soliton constructed independently in [4] and [23]. The metrics constructed in [17] should converge to generalisations of the Koiso-Cao soliton due to Dancer and Wang [13] which are known as Dancer-Wang Kähler-Ricci solitons.

2.2. Linear stability for Ricci flow. Ricci solitons first arose as the fixed points up to gauge of the Ricci flow

$$\frac{\partial g}{\partial t} = -2\text{Ric}(g).$$

In particular, Einstein metrics evolve via homothetic rescaling. Perelman [30] introduced a quantity ν which is monotonically increasing along a Ricci flow and stationary only at shrinking Ricci solitons and, in particular, at Einstein metrics with positive Einstein constant λ . If the second variation of ν at an Einstein metric is positive, then a small perturbation of the metric will increase ν and the Ricci flow cannot flow back to the Einstein metric. Hence the Einstein metric will be unstable. The linear stability of Ricci solitons was considered by Cao, Hamilton and Ilmanen [6], Cao and Zhu [8], and the second and third authors [18]. In order to state their theorem we need to introduce the operators

$$\operatorname{div}_\phi(\cdot) := e^{-\phi} \operatorname{div}(e^\phi \cdot) = \operatorname{div}(\cdot) - \iota_{\nabla\phi}(\cdot),$$

and

$$\Delta_\phi(\cdot) := \Delta(\cdot) - \nabla_{\nabla\phi}(\cdot).$$

Theorem 2.1 (Cao–Hamilton–Ilmanen [6], Cao–Zhu [8]). *Let (M, g, ϕ) be a gradient Ricci soliton with potential function ϕ and constant λ . Let $h \in s^2(TM)$. Then*

$$\frac{d^2}{ds^2} \nu(g + sh)|_{s=0} = \frac{(2\lambda)^{-1}}{(8\pi\lambda)^{n/2}} \int_M \langle N(h), h \rangle e^{-\phi} dV_g,$$

where

$$N(h) = \frac{1}{2} \Delta_\phi(h) + \operatorname{Rm}(h, \cdot) + \operatorname{div}^* \operatorname{div}_\phi h + \frac{1}{2} \nabla^2 v_h - \left(\frac{\int_M \langle \operatorname{Ric}(g), h \rangle e^{-\phi} dV_g}{\int_M \operatorname{scal}(g) e^{-\phi} dV_g} \right) \operatorname{Ric}(g), \quad (2.4)$$

$\operatorname{scal}(g)$ is the scalar curvature of g , and v_h is the unique solution to

$$\Delta_\phi v_h + \lambda v_h = \operatorname{div}_\phi \operatorname{div}_\phi h.$$

As Perelman's ν -entropy is invariant under homothetic rescaling and diffeomorphisms of the metric, we restrict to perturbations given by tensors h satisfying the gauge-fixing conditions

$$\operatorname{div}_\phi(h) = 0 \text{ and } \int_M \langle \operatorname{Ric}(g), h \rangle e^{-\phi} dV_g = 0.$$

In the Einstein case (where the soliton potential ϕ is constant), the stability operator (2.4) restricted to these tensors is given by

$$N(h) = \frac{1}{2} (\Delta_L h + 2\lambda h),$$

where Δ_L is the Lichnerowicz Laplacian

$$\Delta_L h = \Delta h + 2\operatorname{Rm}(h, \cdot) - \operatorname{Ric} \cdot h - h \cdot \operatorname{Ric}. \quad (2.5)$$

Hence we can state a stability criterion for Einstein metrics in terms of the spectrum of the Lichnerowicz Laplacian.

Definition 2.2 (Linear stability of Einstein metrics [6], [7]). *Let (M, g) be a compact Einstein manifold satisfying $\operatorname{Ric}(g) = \lambda g$ and let $-\kappa$ be the largest eigenvalue of the Lichnerowicz Laplacian restricted to the space of divergence-free, g -orthogonal tensors.*

- (1) If $\kappa > 2\lambda$, g is called *linearly stable*.
- (2) If $\kappa = 2\lambda$, g is called *neutrally linearly stable*.
- (3) If $\kappa < 2\lambda$, g is called *linearly unstable*.

Sesum has related the notions of linear stability and the dynamical stability of the Ricci flow [33]. In particular, an Einstein metric g_0 is dynamically stable if there exists a C^k -neighbourhood ($k \geq 3$) U of g_0 such that the λ -rescaled Ricci flows

$$\frac{\partial g}{\partial t} = -2\text{Ric}(g) + 2\lambda g$$

converge to g_0 for all initial metrics $g \in U$. If no such neighbourhood exists then we say g_0 is unstable. This article is concerned with unstable metrics and we note the following result of Sesum.

Proposition 2.3 ([33]). *Let (M, g) be a compact Einstein manifold satisfying $\text{Ric}(g) = \lambda g$. If the Lichnerowicz Laplacian has a divergence-free, g -orthogonal eigentensor with eigenvalue $-\kappa$ satisfying*

$$\kappa < 2\lambda,$$

then g is unstable as a fixed point of the Ricci flow.

It is expected that Einstein metrics which are stable under the Ricci flow are quite special. In dimension four Richard Hamilton has conjectured that the only linearly stable examples of positively curved Einstein metrics (or Ricci solitons) are \mathbb{S}^4 and $\mathbb{C}\mathbb{P}^2$ with their standard metrics. The Fubini–Study on $\mathbb{C}\mathbb{P}^n$ metric is neutrally linearly stable as the eigentensors of the Lichnerowicz Laplacian achieve the bound -2λ . Recent work by Kröncke [26] showed the surprising result that the Fubini–Study metric on $\mathbb{C}\mathbb{P}^n$ is not dynamically stable. Various works [12], [16], [18], [19], [20] have provided case-by-case evidence for Hamilton’s conjecture but as yet, very little general theory exists. Recent work by Pali has addressed this in the case of Kähler–Ricci solitons and Kähler–Ricci flow [28], [29]. By looking at compact symmetric spaces, Cao and He showed that there do exist a wider variety of stable Einstein metrics in higher dimensions [7].

In the case that function f in Equations (2.1) and (2.2) is constant, we recover the usual notion of a product Einstein metric. It is well-known that ordinary products can be destabilised by ‘inflating’ one of the factors. The destabilising perturbations we use to prove the main theorems follow a similar idea but of course the presence of the non-constant warping factor f complicates this procedure.

We end this section with a lemma that will prove useful in subsequent calculations.

Lemma 2.4. *Let (M^n, g) be an Einstein manifold with Einstein constant $\lambda > 0$ and let h be a divergence-free tensor. Then:*

(1) *The tensor $h + cg$ satisfies*

$$\int_M \langle h + cg, g \rangle dV_g = 0,$$

where

$$c = \frac{-\int_M \text{tr}(h)dV_g}{n\text{Vol}(M)}.$$

(2) *The stability integral for $h + cg$ is given by*

$$\langle N(h + cg), h + cg \rangle_{L^2(g)} = \langle \frac{1}{2}\Delta h + \text{Rm}(h, \cdot), h \rangle_{L^2(g)} - \lambda \frac{(\int_M \text{tr}(h)dV_g)^2}{n\text{Vol}(M)}.$$

Proof. (1) is a trivial calculation. To see (2) note that by Equation 2.4, the stability operator N is given by

$$N(h + cg) = \frac{1}{2}\Delta(h + cg) + \text{Rm}(h + cg, \cdot).$$

Hence we see

$$\langle N(h + cg), h + cg \rangle_{L^2(g)} = \langle \frac{1}{2} \Delta(h + cg) + \text{Rm}(h + cg, \cdot), h + cg \rangle_{L^2(g)}.$$

Using the fact that $\Delta(cg) = 0$ and, as g is an Einstein metric,

$$\langle \text{Rm}(h + cg, \cdot), cg \rangle_{L^2(g)} = c\lambda \int_M \text{tr}(h + cg) dV_g = 0,$$

we obtain

$$\langle N(h + cg), h + cg \rangle_{L^2(g)} = \langle \frac{1}{2} \Delta h + \text{Rm}(h, \cdot), h \rangle_{L^2(g)} + \langle \text{Rm}(cg, \cdot), h \rangle_{L^2(g)}.$$

The claim follows by noting $\text{Rm}(cg, \cdot) = c\text{Ric}(g) = c\lambda g$ and the value of c from part (1). \square

2.3. Black hole stability. In [14] the authors developed the stability theory of generalised Schwarzschild–Tangherlini black holes. These are metrics of the form

$$d\hat{s}^2 = - \left[1 - \left(\frac{l}{r} \right)^{d-1} \right] dt^2 + \frac{dr^2}{\left[1 - \left(\frac{l}{r} \right)^{d-1} \right]} + r^2 ds_d^2,$$

where l is a constant and ds_d^2 is the metric on a d dimensional compact Einstein manifold normalised so that its Einstein constant is $d-1$. They found a stability criterion involving the spectrum of the Lichnerowicz Laplacian restricted to the divergence-free (called transverse in the physics literature), trace-free tensors on the Einstein manifold. We state their stability criterion with respect to our convention that the ordinary Laplacian has a non-positive spectrum (this is the opposite convention to that taken in [14]).

Proposition 2.5 (Black hole linear stability [14]). *Let (M^n, g) be a compact Einstein manifold satisfying $\text{Ric}(g) = \lambda g$. Then the associated Schwarzschild–Tangherlini black hole is linearly unstable if the Lichnerowicz Laplacian has a divergence-free, trace-free eigentensor with eigenvalue $-\kappa$ satisfying*

$$\kappa < \frac{\lambda}{n-1} \left(4 - \frac{(5-n)^2}{4} \right) = \frac{(9-n)\lambda}{4}.$$

Note that if an Einstein metric is unstable in the black hole sense, then it is unstable as a fixed point of the Ricci flow. We also note that we require genuinely trace-free perturbations in this definition rather than perturbations which are L^2 -orthogonal to the metric (i.e. the integral of the trace of the perturbation is zero).

In [15], Gibbons, Hartnoll and Pope investigated the linear stability of the Böhm warped product metrics. They proved that the Böhm metrics (or rather the associated Schwarzschild–Tangherlini black holes) on $\mathbb{S}^3 \times \mathbb{S}^m$ for $m = 2, 3$ are unstable. Their proof used the cohomogeneity one symmetry that the Böhm metrics exhibit. Our Theorem E is a generalisation of this result to an arbitrary warped product metric on these spaces.

2.4. A heuristic for destabilising one parameter families of warped products. The methods for proving the Theorems A and B are inspired by considering how to destabilise a product gradient Ricci soliton. Let (B, \bar{g}, ϕ) be a gradient Ricci soliton satisfying

$$\text{Ric}(\bar{g}) + \bar{\nabla}^2 \phi = \lambda \bar{g},$$

and let (F, \tilde{g}) be an Einstein manifold satisfying

$$\text{Ric}(\tilde{g}) = \lambda \tilde{g},$$

where $\lambda > 0$. Then the metric $g = \bar{g} \oplus \tilde{g}$ is a gradient Ricci soliton on $M = B \times F$ with potential function $\phi \circ \pi_B$ (as usual, we will drop the reference to the projection from now on and also denote this function ϕ).

As mentioned previously, it is natural to use gauge-fixed tensors to destabilise. For a Ricci soliton, this means choosing tensors which are div_ϕ -free and L^2 -orthogonal, with respect to the weighted volume form $e^{-\phi} dV_{\bar{g}}$, to the Ricci tensor. On a product soliton there are two natural tensors satisfying the gauge-fixing conditions. The first is the tensor

$$h_1 = e^\phi \left(\frac{\bar{g}}{n} \oplus -\frac{1}{m} \tilde{g} \right).$$

The second is the tensor

$$h_2 = \text{Ric}(\bar{g}) \oplus c \tilde{g},$$

with the constant c chosen so that

$$\int_M \langle \text{Ric}(g), h_2 \rangle e^{-\phi} dV_{\bar{g}} = 0.$$

One can compute the stability integral in Theorem 2.1 for each of the perturbations.

Proposition 2.6. *Let $(B \times F, g = \bar{g} \oplus \tilde{g}, \phi)$ be a gradient product Ricci soliton and let h_1 and h_2 be as above. Then*

$$\langle N(h_1), h_1 \rangle_{L^2(e^{-\phi} dV_g)} = \text{Vol}(F) \int_B \left(-\left(\frac{1}{2n} + \frac{1}{2m} - \frac{1}{n^2} \right) |\bar{\nabla} e^\phi|^2 + \left(\frac{1}{n} + \frac{1}{m} \right) \lambda e^{2\phi} \right) e^{-\phi} dV_{\bar{g}},$$

and

$$\langle N(h_2), h_2 \rangle_{L^2(e^{-\phi} dV_g)} = \lambda \|h_2\|_{L^2(e^{-\phi} dV_g)}^2.$$

Proof. As h_1 satisfies the gauge-fixing conditions $\text{div}_\phi(h_1) = 0$ and $\langle \text{Ric}(g), h_1 \rangle_{L^2(e^{-\phi} dV_g)} = 0$, equation (2.4) reduces to

$$N(h_1) = \frac{1}{2} \Delta_\phi h_1 + \text{Rm}(h_1, \cdot).$$

As we are working with the rescaled volume form $e^{-\phi} dV_g$ we can integrate the first term by parts and get

$$\frac{1}{2} \langle \Delta_\phi h_1, h_1 \rangle_{L^2(e^{-\phi} dV_g)} = \int_M -\frac{1}{2} |\nabla h_1|^2 e^{-\phi} dV_g.$$

We then compute

$$|\nabla h_1|^2 = |\bar{\nabla} e^\phi|^2 \left| \frac{\bar{g}}{n} \oplus \frac{-\tilde{g}}{m} \right|^2 = |\bar{\nabla} e^\phi|^2 \left(\frac{1}{n} + \frac{1}{m} \right).$$

Hence

$$\frac{1}{2} \langle \Delta_\phi h_1, h_1 \rangle_{L^2(e^{-\phi} dV_g)} = \text{Vol}(F) \int_B -|\bar{\nabla} e^\phi|^2 \left(\frac{1}{2n} + \frac{1}{2m} \right) e^{-\phi} dV_{\bar{g}}.$$

The curvature operator term is given by

$$\text{Rm}(h_1, \cdot) = e^\phi \left(\frac{\text{Ric}(\bar{g})}{n} \oplus -\lambda \frac{\tilde{g}}{m} \right),$$

thus, using the fact \bar{g} is a gradient Ricci soliton,

$$\langle \text{Rm}(h_1, \cdot), h_1 \rangle = \left(-\frac{\bar{\Delta}\phi}{n^2} + \lambda \left(\frac{1}{n} + \frac{1}{m} \right) \right) e^{2\phi}.$$

We note the identity $|\bar{\nabla}\phi|^2 e^\phi = |\bar{\nabla}e^\phi|^2 e^{-\phi}$ and so integrating by parts yields

$$\langle \text{Rm}(h_1, \cdot), h_1 \rangle_{L^2(e^{-\phi}dV_g)} = \text{Vol}(F) \int_B \left(\frac{|\bar{\nabla}e^\phi|^2}{n^2} + \lambda e^{2\phi} \left(\frac{1}{n} + \frac{1}{m} \right) \right) e^{-\phi} dV_{\bar{g}}.$$

Combining the Laplacian and curvature operator terms yields the identity for the h_1 tensor.

For the variation h_2 we note that the results of Cao and Zhu [8] show that h_2 is div_ϕ -free and of course, from the choice of c , is $L^2(e^{-\phi}dV_g)$ -orthogonal to $\text{Ric}(g)$. Hence we compute as with the tensor h_1 ,

$$\frac{1}{2}\Delta_\phi h_2 + \text{Rm}(h_2, \cdot) = \left(\frac{1}{2}\bar{\Delta}_\phi(\text{Ric}(\bar{g})) + \overline{\text{Rm}}(\text{Ric}(\bar{g}), \cdot) \right) \oplus c \left(\frac{1}{2}\tilde{\Delta}(\tilde{g}) + \widetilde{\text{Rm}}(\tilde{g}, \cdot) \right).$$

We again use the results of Cao and Zhu [8] who show that (as \bar{g} is a gradient Ricci soliton)

$$\frac{1}{2}\bar{\Delta}_\phi(\text{Ric}(\bar{g})) + \overline{\text{Rm}}(\text{Ric}(\bar{g}), \cdot) = \lambda \text{Ric}(\bar{g}).$$

Finally, as \tilde{g} is Einstein,

$$\frac{1}{2}\tilde{\Delta}(\tilde{g}) + \widetilde{\text{Rm}}(\tilde{g}, \cdot) = \lambda c \tilde{g},$$

and we see $N(h_2) = \lambda h_2$. The result follows. \square

While it is clear from Proposition 2.6 that the perturbation h_2 always destabilises, it is not clear that this is true for h_1 . Note that

$$\int_B |\bar{\nabla}e^\phi|^2 e^{-\phi} dV_{\bar{g}} = \frac{1}{2} \int_B \langle \bar{\nabla}\phi, \bar{\nabla}e^{2\phi} \rangle e^{-\phi} dV_{\bar{g}}.$$

It is well-known [5] that it is also possible to normalise ϕ so that

$$\bar{\Delta}_\phi \phi = -2\lambda\phi.$$

In this case

$$\int_B |\bar{\nabla}e^\phi|^2 e^{-\phi} dV_{\bar{g}} = \lambda \int_B \phi e^\phi dV_{\bar{g}},$$

and the perturbation h_1 is destabilising if

$$\int_B \phi e^\phi dV_{\bar{g}} < \frac{2n(m+n)}{n^2 + mn - 2m} \int_B e^\phi dV_{\bar{g}}.$$

Such an inequality does not in general hold for functions ψ satisfying

$$\int_B \psi e^{-\psi} dV_{\bar{g}} = 0,$$

and so it is not possible to conclude that h_1 is a destabilising perturbation of a product Ricci soliton.

Suppose that there is a sequence of quasi-Einstein metrics (M, \bar{g}_i, f_i, m_i) solving Equations (2.2) and (2.1) for some fixed λ and μ_i (we are always free to fix λ by rescaling the g_i). Setting $\phi_i = -m_i \log f_i$, Equation (2.2) becomes

$$\text{Ric}(\bar{g}_i) + \bar{\nabla}^2 \phi_i - \frac{d\phi_i \otimes d\phi_i}{m_i} = \lambda \bar{g}_i.$$

If as $i \rightarrow \infty$, $m_i \rightarrow \infty$, the metrics \bar{g}_i and the functions ϕ_i converge (as we are only outlining some heuristic reasoning we do not make this notion of convergence precise), then the limiting metric \bar{g}_∞ and function φ solve the gradient Ricci soliton equation

$$\text{Ric}(\bar{g}_\infty) + \bar{\nabla}^2 \varphi = \lambda \bar{g}_\infty.$$

For large values of m_i , we can make the following approximations

$$\bar{g}_i \approx \bar{g}_\infty, \quad f_i \approx 1, \quad \text{and} \quad f_i^{m_i} \approx e^{-\varphi}.$$

Hence the warped product Einstein metric $\bar{g}_i \oplus f_i^2 \tilde{g}$ can be approximated by the metric $\bar{g}_\infty \oplus \tilde{g}$. We shall see in Section 5 that $\mu_i \approx \lambda$, and so, for large values of m_i , the metric $\bar{g}_i \oplus \tilde{g}$ is almost a product Ricci soliton.

In Sections 4 and 5 we define two different tensors. The first are the GHP variations (Definition 4.1) which are the analogues of the tensor h_1 . The fact that h_1 is not obviously universally destabilising goes some way to explain why using GHP variations fails to destabilise warped products in all but the lowest dimensions. The Ricci variation (Definition 5.1) is the analogue of h_2 and Theorem B could be paraphrased as saying that, providing the metrics \bar{g}_i and \bar{g} are close to each other, the fact that h_2 is universally destabilising means the Ricci variation also destabilises the warped product for large values of i .

3. GEOMETRIC OPERATORS FOR WARPED PRODUCTS

In this section we collect some useful identities that are used in the proof of the main theorems. All of the theorems involve choosing a potentially destabilising tensor $h \in s^2(TM)$ and then computing the Rayleigh quotient

$$\frac{\int_M \langle \Delta_L h, h \rangle dV_g}{\int_M |h|^2 dV_g},$$

which provides a lower bound for the least negative eigentensor of Δ_L . The class of destabilising tensors that we consider can be written in the form

$$h = \bar{h} \oplus \psi \tilde{h}, \tag{3.1}$$

where $\bar{h} \in s^2(TB)$, $\tilde{h} \in s^2(TF)$ and $\psi \in C^\infty(B)$.

One fundamental calculation is of the Christoffel symbols for the Levi-Civita connection of a warped product metric.

Lemma 3.1 (Christoffel symbols of g). *Let $M = B \times F$ be a product manifold and let $g = \bar{g} \oplus f^2 \tilde{g}$ be a warped product metric on M . Then the Christoffel symbols for the*

Levi-Civita connection of g are given by:

$$\begin{aligned}\Gamma_{ab}^c &= \bar{\Gamma}_{ab}^c, \\ \Gamma_{\alpha\beta}^c &= -\tilde{g}_{\alpha\beta} f \bar{g}^{cd} (\bar{\nabla}_d f), \\ \Gamma_{a\beta}^\gamma &= (\bar{\nabla}_a \log f) \delta_{\beta}^\gamma, \\ \Gamma_{\alpha\beta}^\gamma &= \tilde{\Gamma}_{\alpha\beta}^\gamma.\end{aligned}$$

All other symbols are zero.

It will also be useful to know an explicit form of the curvature tensor.

Lemma 3.2 (Curvature tensor of g). *Let $M = B \times F$ be a product manifold and let $g = \bar{g} \oplus f^2 \tilde{g}$ be a warped product metric on M . Then the curvature tensor for g can be described by:*

$$\begin{aligned}R_{abcd} &= \bar{R}_{abcd}, \\ R_{a\beta\gamma d} &= f \tilde{g}_{\beta\gamma} (\bar{\nabla}^2 f)_{ad}, \\ R_{\alpha\beta\gamma\delta} &= f^2 \widetilde{\text{Rm}}_{\alpha\beta\gamma\delta} - f^2 |\bar{\nabla} f|^2 (\tilde{g}_{\alpha\gamma} \tilde{g}_{\beta\delta} - \tilde{g}_{\alpha\delta} \tilde{g}_{\beta\gamma}).\end{aligned}$$

All other components are zero.

As mentioned in Section 2, one need only check stability on divergence-free tensors. The next lemma computes the divergence of tensors of the form (3.1).

Lemma 3.3 (Divergence of h). *Let $(B \times F^m, \bar{g} \oplus f^2 \tilde{g})$ be a warped product manifold and let h be of the form (3.1). Then*

$$\begin{aligned}\text{div}(h)(\cdot) &= \bar{\text{div}}(\bar{h})(\cdot) + m \bar{h}(\bar{\nabla} \log f, \cdot) \\ &\quad - f^{-2} \psi (\text{tr}(\tilde{h})) d \log f(\cdot) + f^{-2} \psi \widetilde{\text{div}}(\tilde{h})(\cdot).\end{aligned}$$

Proof. In coordinates, as $g^{a\beta} = 0$, we have

$$\text{div}(h)_C = g^{AB} (\nabla_A h)(\partial_B, \partial_C) = \bar{g}^{ab} (\nabla_a h)(\partial_b, \partial_C) + f^{-2} \tilde{g}^{\alpha\beta} (\nabla_\alpha h)(\partial_\beta, \partial_C).$$

Using Lemma 3.1 and the fact $h_{a\beta} = 0$ we see

$$(\nabla_a h)(\partial_b, \partial_\gamma) = 0.$$

Hence $\bar{g}^{ab} (\nabla_a h)(\partial_b, \partial_C) = \bar{\text{div}}(\bar{h})(\partial_C)$. For the next term we again use the Christoffel symbols of Lemma 3.1

$$f^{-2} \tilde{g}^{\alpha\beta} (\nabla_\alpha h)(\partial_\beta, \partial_c) = f^{-2} \tilde{g}^{\alpha\beta} (\nabla_\alpha h_{\beta c} - \Gamma_{\alpha\beta}^d h_{cd} - \Gamma_{\alpha c}^\delta h_{\beta\delta}),$$

which as, $h_{\beta c} = 0$, $\Gamma_{\alpha\beta}^d = -\tilde{g}_{\alpha\beta} f \bar{g}^{de} (\bar{\nabla}_e f)$, and $\Gamma_{\alpha c}^\delta = (\bar{\nabla}_c \log f) \delta_{\alpha}^\delta$, yields

$$f^{-2} \tilde{g}^{\alpha\beta} (\nabla_\alpha h)(\partial_\beta, \partial_c) = m \bar{h}(\bar{\nabla} \log f, \partial_c) - f^{-2} \psi \text{tr}(\tilde{h})(\bar{\nabla}_c \log f).$$

Finally, we consider

$$f^{-2} \tilde{g}^{\alpha\beta} (\nabla_\alpha h)(\partial_\beta, \partial_\gamma) = f^{-2} \tilde{g}^{\alpha\beta} (\nabla_\alpha h_{\beta\gamma} - \Gamma_{\alpha\beta}^\delta h_{\gamma\delta} - \Gamma_{\alpha\gamma}^\delta h_{\beta\delta}).$$

Hence as $h_{\alpha\beta} = \psi \tilde{h}_{\alpha\beta}$,

$$f^{-2} \tilde{g}^{\alpha\beta} (\nabla_\alpha h)(\partial_\beta, \partial_\gamma) = f^{-2} \psi \widetilde{\text{div}}(\tilde{h})(\partial_\gamma).$$

Combining each piece finishes the proof. \square

To break down the calculation of the Lichnerowicz Laplacian, we first compute the connection Laplacian of the tensors h .

Lemma 3.4 (Connection Laplacian). *Let $(B \times F^m, \bar{g} \oplus f^2 \tilde{g})$ be a warped product manifold and let h be of the form (3.1). Then*

$$\begin{aligned} \Delta h &= \bar{\Delta} \bar{h} + 2f^{-2} \psi \tilde{\text{tr}}(\tilde{h})(d \log f \otimes d \log f) \\ &\quad - m(d \log f \otimes \iota_{\bar{\nabla} \log f} \bar{h} + \iota_{\bar{\nabla} \log f} \bar{h} \otimes d \log f - (\bar{\nabla}_{\bar{\nabla} \log f} \bar{h})) \\ &\quad + (\bar{\Delta} \psi - 2\psi \bar{\Delta} \log f + (m-4)\bar{g}(\bar{\nabla} \psi, \bar{\nabla} \log f) + 2(1-m)\psi |\bar{\nabla} \log f|^2) \tilde{h} \\ &\quad + f^{-2} \psi \tilde{\Delta} \tilde{h} + 2\bar{h}(\bar{\nabla} f, \bar{\nabla} f) \tilde{g} \\ &\quad - 2\psi f^{-2} (d \log f \otimes \tilde{\text{div}}(\tilde{h}) + \tilde{\text{div}}(\tilde{h}) \otimes d \log f). \end{aligned}$$

Proof. We begin by noting that, for any tensor T , $\nabla_{A,B}^2 T = \nabla_A \nabla_B T - \nabla_{\nabla_A B} T$. We need to compute

$$\Delta h = g^{AB} \nabla_{A,B}^2 h = \bar{g}^{ab} \nabla_{a,b}^2 h + f^{-2} \tilde{g}^{\alpha\beta} \nabla_{\alpha,\beta}^2 h.$$

The proof proceeds by using the coordinate description of the connection given in Lemma 3.1. We will compute each separate part of Δh .

We begin by computing $(\Delta h)_{cd}$. We can easily verify that

$$(\nabla_a \nabla_b h)_{cd} = (\bar{\nabla}_a \bar{\nabla}_b \bar{h})_{cd} \quad (3.2)$$

and

$$(\nabla_{\nabla_a b} h)_{cd} = (\bar{\nabla}_{\bar{\nabla}_a b} \bar{h})_{cd}. \quad (3.3)$$

To compute the $(\nabla_{\alpha,\beta}^2 h)_{cd}$ terms, we expand

$$(\nabla_\alpha \nabla_\beta h)_{cd} = \nabla_\alpha (\nabla_\beta h)_{cd} - (\nabla_\beta h)(\nabla_\alpha c, \partial_d) - (\nabla_\beta h)(\partial_c, \nabla_\alpha d),$$

which yields

$$(\nabla_\alpha \nabla_\beta h)_{cd} = \nabla_\alpha (\nabla_\beta h)_{cd} - (\bar{\nabla}_c \log f)(\nabla_\beta h)_{\alpha d} - (\bar{\nabla}_d \log f)(\nabla_\beta h)_{c\alpha}.$$

A straightforward check shows

$$(\nabla_\beta h)_{cd} = 0,$$

and, for example,

$$(\nabla_\beta h)_{\alpha d} = \tilde{g}_{\alpha\beta} f \bar{h}(\bar{\nabla} f, \partial_d) - \psi(\bar{\nabla}_d \log f) \tilde{h}_{\alpha\beta}.$$

Putting all this together we obtain

$$(\nabla_\alpha \nabla_\beta h)_{cd} = 2\psi \tilde{h}_{\alpha\beta} (\bar{\nabla}_c \log f \bar{\nabla}_d \log f) - \tilde{g}_{\alpha\beta} f ((\bar{\nabla}_c \log f) \bar{h}(\bar{\nabla} f, \partial_d) + (\bar{\nabla}_d \log f) \bar{h}(\bar{\nabla} f, \partial_c)). \quad (3.4)$$

In a similar manner we see that

$$(\nabla_{\nabla_\alpha \beta} h)_{cd} = -\tilde{g}_{\alpha\beta} f (\bar{\nabla}_{\bar{\nabla}_f} \bar{h})_{cd}. \quad (3.5)$$

Combining Equations (3.2) and (3.3), and (3.4) and (3.5), then taking traces yields

$$\begin{aligned} (\Delta h)_{cd} &= (\bar{\Delta} \bar{h})_{cd} + 2f^{-2} \psi \tilde{\text{tr}}(\tilde{h})(d \log f \otimes d \log f)_{cd} \\ &\quad - m(d \log f \otimes \iota_{\bar{\nabla} \log f} \bar{h} + \iota_{\bar{\nabla} \log f} \bar{h} \otimes d \log f - (\bar{\nabla}_{\bar{\nabla} \log f} \bar{h}))_{cd}. \end{aligned}$$

We now compute $(\Delta h)_{\gamma\delta}$. The term

$$(\nabla_a \nabla_b h)_{\gamma\delta} = \nabla_a (\nabla_b h)_{\gamma\delta} - (\nabla_b h)(\nabla_a \gamma, \partial_\delta) - (\nabla_b h)(\partial_\gamma, \nabla_a \delta),$$

is given by

$$(\nabla_a \nabla_b h)_{\gamma\delta} = \nabla_a (\nabla_b h)_{\gamma\delta} - 2(\bar{\nabla}_a \log f)(\nabla_b h)_{\gamma\delta}.$$

We expand

$$(\nabla_b h)_{\gamma\delta} = (\bar{\nabla}_b \psi - 2\psi \bar{\nabla}_b \log f) \tilde{h}_{\gamma\delta},$$

which yields

$$\begin{aligned} & (\nabla_a \nabla_b h)_{\gamma\delta} = \\ & (\bar{\nabla}_a \bar{\nabla}_b \psi - 2(\bar{\nabla}_a \psi \bar{\nabla}_b \log f + \bar{\nabla}_b \psi \bar{\nabla}_a \log f) - 2\psi \bar{\nabla}_a \bar{\nabla}_b \log f + 4\psi \bar{\nabla}_a \log f \bar{\nabla}_b \log f) \tilde{h}_{\gamma\delta}. \end{aligned} \quad (3.6)$$

Similarly, we obtain

$$(\nabla_{\nabla_a b} h)_{\gamma\delta} = (\bar{\nabla}_{\bar{\nabla}_a b} \psi - 2\psi \bar{\nabla}_{\bar{\nabla}_a b} \log f) \tilde{h}_{\gamma\delta}. \quad (3.7)$$

To compute $(\nabla_\alpha \nabla_\beta h)_{\delta\gamma}$ we expand,

$$\begin{aligned} & (\nabla_\alpha \nabla_\beta h)_{\delta\gamma} = \nabla_\alpha (\nabla_\beta h)_{\gamma\delta} - (\nabla_\beta h) (\nabla_\alpha \gamma, \partial_\delta) - (\nabla_\beta h) (\partial_\gamma, \nabla_\alpha \delta), \\ & = \nabla_\alpha (\nabla_\beta h)_{\gamma\delta} - (\nabla_\beta h) (\tilde{\nabla}_\alpha \gamma, \partial_\delta) - (\nabla_\beta h) (\partial_\gamma, \tilde{\nabla}_\alpha \delta) + f \tilde{g}_{\alpha\gamma} (\nabla_\beta h) (\bar{\nabla} f, \partial_\delta) + f \tilde{g}_{\alpha\delta} (\nabla_\beta h) (\partial_\gamma, \bar{\nabla} f). \end{aligned}$$

Noting that $(\nabla_\beta h)_{\delta\gamma} = \psi (\tilde{\nabla}_\beta \tilde{h})_{\delta\gamma}$ and, for example,

$$(\nabla_\beta h) (\bar{\nabla} f, \partial_\delta) = -f |\bar{\nabla} \log f|^2 \psi \tilde{h}_{\beta\delta} + f \tilde{g}_{\beta\delta} \bar{h} (\bar{\nabla} f, \bar{\nabla} f),$$

we obtain

$$\begin{aligned} & (\nabla_\alpha \nabla_\beta h)_{\gamma\delta} = \\ & \psi \tilde{\nabla}_\alpha (\tilde{\nabla}_\beta \tilde{h})_{\gamma\delta} + f^2 (\tilde{g}_{\alpha\gamma} \tilde{g}_{\beta\delta} + \tilde{g}_{\alpha\delta} \tilde{g}_{\beta\gamma}) \bar{h} (\bar{\nabla} f, \bar{\nabla} f) - f^2 |\bar{\nabla} \log f|^2 \psi ((\tilde{g}_{\alpha\gamma} \tilde{h}_{\beta\delta} + \tilde{g}_{\alpha\delta} \tilde{h}_{\beta\gamma})). \end{aligned} \quad (3.8)$$

In a similar manner we see that

$$(\nabla_{\nabla_\alpha \beta} h)_{\gamma\delta} = \psi (\tilde{\nabla}_{\tilde{\nabla}_\alpha \beta} \tilde{h})_{\gamma\delta} + \tilde{g}_{\alpha\beta} f^2 (2\psi |\bar{\nabla} \log f|^2 - \bar{g} (\bar{\nabla} \log f, \bar{\nabla} \psi)) \tilde{h}_{\gamma\delta}. \quad (3.9)$$

Combining Equations (3.6) and (3.7), and (3.8) and (3.9), then taking traces yields

$$\begin{aligned} (\Delta h)_{\gamma\delta} & = \\ & (\bar{\Delta} \psi - 2\psi \bar{\Delta} \log f + (m-4)\bar{g} (\bar{\nabla} \psi, \bar{\nabla} \log f) + 2(1-m)\psi |\bar{\nabla} \log f|^2) \tilde{h}_{\gamma\delta} \\ & + f^{-2} \psi (\tilde{\Delta} \tilde{h})_{\gamma\delta} + 2\bar{h} (\bar{\nabla} f, \bar{\nabla} f) \tilde{g}_{\gamma\delta}. \end{aligned}$$

Finally, we calculate $(\Delta h)_{c\delta}$. As $(\nabla_b h)_{c\delta} = 0$, we find

$$(\nabla_a \nabla_b h)_{c\delta} = 0 \quad \text{and} \quad (\nabla_{\nabla_a b} h)_{c\delta} = 0.$$

Expanding $(\nabla_\alpha \nabla_\beta h)_{c\delta}$ yields

$$(\nabla_\alpha \nabla_\beta h)_{c\delta} = \nabla_\alpha (\nabla_\beta h)_{c\delta} - (\bar{\nabla}_c \log f) (\nabla_\beta h)_{\alpha\delta} - (\nabla_\beta h) (\partial_c, \tilde{\nabla}_\alpha \delta) + f \tilde{g}_{\alpha\delta} (\nabla_\beta h) (\partial_c, \bar{\nabla} f).$$

Computing each term yields

$$\begin{aligned} (\nabla_\beta h)_{c\delta} & = -\bar{\nabla}_c \log f \psi \tilde{h}_{\beta\delta} + f \tilde{g}_{\beta\delta} \bar{h} (\partial_c, \bar{\nabla} f), \\ (\nabla_\beta h)_{\alpha\delta} & = \psi (\tilde{\nabla}_\beta \tilde{h})_{\alpha\delta}, \\ (\nabla_\beta h) (\partial_c, \tilde{\nabla}_\alpha \delta) & = -(\bar{\nabla}_c \log f) \psi \tilde{h} (\partial_\beta, \tilde{\nabla}_\alpha \delta) + f \tilde{g} (\partial_\beta, \tilde{\nabla}_\alpha \delta) \bar{h} (\partial_c, \bar{\nabla} f), \end{aligned}$$

and

$$(\nabla_\beta h) (\partial_c, \bar{\nabla} f) = 0.$$

We also have

$$(\nabla_{\nabla_\alpha \beta} h)_{c\delta} = -(\bar{\nabla}_c \log f) \psi \tilde{h} (\tilde{\nabla}_\alpha \beta, \partial_\delta) + f \tilde{g} (\tilde{\nabla}_\alpha \beta, \partial_\delta) \bar{h} (\partial_c, \bar{\nabla} f).$$

Putting this all together, we have

$$(\nabla_{\alpha,\beta}^2 h)_{c\delta} = -(\bar{\nabla}_c \log f) \psi \left((\tilde{\nabla}_\alpha \tilde{h})_{\beta\delta} + (\tilde{\nabla}_\beta \tilde{h})_{\alpha\delta} \right) + f \bar{h} (\bar{\nabla} f, \partial_c) (\tilde{\nabla}_\alpha \tilde{g})_{\beta\delta}.$$

Taking traces and noting that $\widetilde{\nabla}\tilde{g} = 0$ gives

$$(\Delta h)_{c\delta} = -2\psi f^{-2}(\widetilde{\nabla}_c \log f)\widetilde{\text{div}}(\tilde{h})_\delta.$$

□

The other component of the Lichnerowicz Laplacian is the curvature operator.

Lemma 3.5 (Curvature operator). *Let $(B \times F^m, \bar{g} \oplus f^2\tilde{g})$ be a warped product manifold. and let h be of the form (3.1). Then*

$$\begin{aligned} \text{Rm}(h, \cdot) &= \overline{\text{Rm}}(\bar{h}, \cdot) - f^{-3}\psi\tilde{\text{tr}}(\tilde{h})(\widetilde{\nabla}^2 f) + f^{-2}\psi\widetilde{\text{Rm}}(\tilde{h}, \cdot) \\ &\quad - \psi|\widetilde{\nabla} \log f|^2(\tilde{\text{tr}}(\tilde{h})\tilde{g} - \tilde{h}) - f\langle \widetilde{\nabla}^2 f, \bar{h} \rangle_{\bar{g}}\tilde{g}. \end{aligned}$$

Proof. The curvature operator is given by

$$\text{Rm}(h, \cdot)_{AB} = R_{ACBD}h^{CD},$$

which, given the form of h , can be written

$$\text{Rm}(h, \cdot)_{AB} = R_{AcBd}\bar{h}^{cd} + \psi f^{-4}R_{A\gamma B\delta}\tilde{h}^{\gamma\delta}.$$

(The factor of f^{-4} appears as we are raising two indices of \tilde{h}). We now use the explicit components of the curvature tensor given in Lemma 3.2.

As $R_{\alpha\beta\gamma d} = R_{abc\delta} = 0$ we see that $\{A, B\} = \{a, b\}$ or $\{A, B\} = \{\alpha, \beta\}$. For the former set we obtain

$$\begin{aligned} \text{Rm}(h, \cdot)_{ab} &= R_{acbd}\bar{h}^{cd} + \psi f^{-4}R_{a\gamma b\delta}\tilde{h}^{\gamma\delta} = \bar{R}_{acbd}\bar{h}^{cd} - \psi f^{-4}fg_{\gamma\delta}(\widetilde{\nabla}^2 f)_{ab}\tilde{h}^{\gamma\delta}, \\ &= \overline{\text{Rm}}(\bar{h}, \cdot)_{ab} - f^{-3}\psi\tilde{\text{tr}}(\tilde{h})(\widetilde{\nabla}^2 f)_{ab}. \end{aligned}$$

For the latter set we obtain

$$\begin{aligned} \text{Rm}(h, \cdot)_{\alpha\beta} &= R_{\alpha c\beta d}\bar{h}^{cd} + \psi f^{-4}R_{\alpha\gamma\beta\delta}\tilde{h}^{\gamma\delta}, \\ &= -f\tilde{g}_{\alpha\beta}(\widetilde{\nabla}^2 f)_{cd}\bar{h}^{cd} + \psi f^{-2}\tilde{R}_{\alpha\gamma\beta\delta}\tilde{h}^{\gamma\delta} - \psi f^{-2}|\widetilde{\nabla} f|^2(\tilde{g}_{\alpha\beta}\tilde{g}_{\gamma\delta} - \tilde{g}_{\alpha\gamma}\tilde{g}_{\beta\delta})\tilde{h}^{\gamma\delta}. \end{aligned}$$

This gives

$$\text{Rm}(h, \cdot)_{\alpha\beta} = -f\langle \widetilde{\nabla}^2 f, \bar{h} \rangle_{\bar{g}}\tilde{g}_{\alpha\beta} + \psi f^{-2}\widetilde{\text{Rm}}(\tilde{h}, \cdot)_{\alpha\beta} - \psi|\widetilde{\nabla} \log f|^2(\tilde{\text{tr}}(\tilde{h})\tilde{g}_{\alpha\beta} - \tilde{h}_{\alpha\beta}),$$

and the result now follows. □

4. THE PROOFS OF THEOREMS A, C AND E

In order to prove Theorems A and E, we consider a generalisation of the variation considered by Gibbons, Hartnoll, and Pope in [15]. Using the terminology of [15], the perturbation is a ‘balloning mode’ which generalises the manner one destabilises an ordinary Riemannian product by changing the relative volumes of the base and fibre.

Definition 4.1 (GHP variations). Let $(B \times F^m, \bar{g} \oplus f^2\tilde{g})$ be a warped product manifold. The GHP variations are the tensors

$$h = \left(\frac{f^k}{n} \bar{g} \right) \oplus \left(\frac{(m+k)f^{k+2}}{mn} \tilde{g} \right), \quad (4.1)$$

where $k \in \mathbb{R} \setminus \{0\}$.

Proposition 4.2. *Let $(B^n \times F^m, \bar{g} \oplus f^2 \tilde{g})$ be a warped product Einstein manifold. Then any GHP variation h is divergence-free. Furthermore if c is the constant defined in Lemma 2.4, then*

$$\langle N(h + cg), h + cg \rangle_{L^2(g)} = C_{n,m,k}^1 \int_B f^{2k+m-2} |\bar{\nabla} f|^2 dV_{\bar{g}} + \lambda \left(\|h\|^2 - \frac{(\int_M \text{tr}(h) dV_g)^2}{\|g\|^2} \right),$$

where

$$C_{n,m,k}^1 = -\frac{\text{Vol}(F)}{2} \left(\frac{k^2(4k + 2m + mn + (m+k)^2)}{n^2 m} \right).$$

Proof. Consider a variation

$$h = Af^k \bar{g} \oplus Bf^{k+2} \tilde{g}.$$

A routine calculation using Lemma 3.3 yields

$$\text{div}(h) = ((m+k)A - mB) f^{k-1} df.$$

Hence if the constants A and B are chosen so that

$$A = \frac{m}{m+k} B,$$

then h is divergence-free. The GHP variation (4.1) has $A = 1/n$ and $B = (m+k)/mn$ and so it is divergence-free. Using Lemma 3.4 with

$$\bar{h} = \frac{f^k}{n} \bar{g}, \quad \tilde{h} = \tilde{g} \quad \text{and} \quad \psi = \frac{(m+k)f^{k+2}}{mn},$$

we obtain

$$\langle \Delta h, h \rangle = \frac{f^k \bar{\Delta} f^k}{n} + \frac{(m+k)^2}{mn^2} f^{k-2} \bar{\Delta} f^{k+2} - 2 \frac{(m+k)^2}{mn^2} f^{2k} \bar{\Delta} \log f + C_{n,m,k}^2 f^{2k-2} |\bar{\nabla} f|^2,$$

where

$$C_{n,m,k}^2 = \frac{km^3 - 4km^2 - 8k^2m - 8km + k^3m - 6k^2 - 4k^3 - 4m^2 + 2k^2m^2 + km^2n}{mn^2}.$$

Integrating by parts yields

$$\int_M \langle \Delta h, h \rangle dV_g = -\text{Vol}(F) \left(\frac{k^2(k^2 + 2km + m^2 + nm + 2)}{mn^2} \right) \int_B f^{2k+m-2} |\bar{\nabla} f|^2 dV_{\bar{g}}.$$

Using Lemma 3.5 (and simplifying using Equations (2.1) and (2.2)) we obtain

$$\langle \text{Rm}(h, \cdot), h \rangle = \left(\frac{(m+k)^2}{mn^2} - \frac{m+2k}{n^2} \right) f^{2k-1} \bar{\Delta} f + \lambda f^{2k} \left(\frac{1}{n} + \frac{(m+k)^2}{mn^2} \right).$$

Integrating by parts yields

$$\int_M \langle \text{Rm}(h, \cdot), h \rangle dV_g = \text{Vol}(F) \left(\frac{-k^2(2k+m-1)}{mn^2} \right) \int_B f^{2k+m-2} |\bar{\nabla} f|^2 dV_{\bar{g}} + \lambda \|h\|_{L^2(g)}^2.$$

The result follows immediately using the simplification given in Lemma 2.4. \square

The proofs of Theorems A and E now follow by taking special values of the parameter k .

Proof of Theorem A. As $k \neq 0$, no GHP variation is a multiple of the metric g and the Cauchy-Schwarz inequality applied to $\langle h, g \rangle_{L^2(g)}$ implies that

$$\|h\|^2 - \frac{(\int_M \text{tr}(h) dV_g)^2}{\|g\|^2} > 0.$$

Hence we wish to select k such that the term $C_{n,m,k}^1$ in Proposition 4.2 is non-negative. Clearly this can only occur if

$$(4k + 2m + mn + (m + k)^2) \leq 0.$$

Viewed as a quadratic in k , $(4k + 2m + mn + (m + k)^2)$ is minimised when $k = -(2 + m)$ giving a value of $((n - 2)m - 4)$. Hence the coefficient $C_{n,m,k}^1$ is only non-negative when $(n - 2)m \leq 4$. The result follows noting that either $n = 3$ and $m \in \{2, 3, 4\}$ or $n = 4$ and $m = 2$. This covers all possible six-dimensional products as if the base has dimension 2 then, by the rigidity theorem of Case–Shu–Wei [9], the function f is constant and the product is trivially unstable. The fibre of a warped product Einstein manifold with positive Einstein constant must be at least two-dimensional, as, by Myers’s theorem the product cannot have infinite fundamental group. \square

Proof of Theorem E. We take $k = -(m + n)$ in Proposition 4.2. In this case the GHP variation h is divergence-free and trace-free. This means that the constant $c = 0$ and so

$$\langle N(h), h \rangle_{L^2(g)} = \frac{1}{2} \langle \Delta_L h + 2\lambda h, h \rangle_{L^2(g)}.$$

Computing using Proposition 4.2 and rearranging yields

$$\langle \Delta_L h, h \rangle_{L^2(g)} = \text{Vol}(F) \left(\frac{(m + n)^2(4n + 2m - n^2 - mn)}{n^2 m} \right) \int_B f^{-2n-m-2} |\bar{\nabla} f|^2 dV_{\bar{g}}.$$

This is manifestly non-negative when $n = 3$ and $m = 2$ or $m = 3$. Hence the Lichnerowicz Laplacian, when restricted to divergence-free, trace-free tensors, has a non-negative eigenvalue $-\kappa$ for some $\kappa \leq 0$. The instability follows from Proposition 2.5 as

$$\kappa \leq 0 < \frac{(9 - (3 + m))\lambda}{4}$$

when $m = 2$ or $m = 3$. \square

We use a similar method to prove Theorem C.

Proof of Theorem C. We consider the perturbation

$$h = 0 \oplus \sigma.$$

Decomposing the variation h in the manner of (3.1) we see that

$$\bar{h} = 0, \quad \tilde{h} = \sigma \text{ and } \psi = 1.$$

This immediately yields $\text{div}(h) = 0$ by Lemma 3.3 and it is clear h is trace-free as σ is trace-free. Using Lemma 3.4 and Lemma 3.5 we obtain

$$\Delta h = f^{-2} \tilde{\Delta} \sigma + (2(1 - m) |\bar{\nabla} \log f|^2 - 2\bar{\Delta} \log f) \sigma,$$

and

$$\text{Rm}(h, \cdot) = f^{-2} \widetilde{\text{Rm}}(\sigma, \cdot) + |\bar{\nabla} \log f|^2 \sigma.$$

Hence

$$\langle \Delta_L h, h \rangle_g = f^{-6} \langle \tilde{\Delta}_L \sigma, \sigma \rangle_{\bar{g}} + 2(\mu f^{-6} - \lambda f^{-4} + (2 - m) f^{-6} |\bar{\nabla} f|^2 - f^{-4} \bar{\Delta} \log f) |\sigma|_{\bar{g}}^2.$$

Integrating by parts and using the fact $\tilde{\Delta}_L \sigma = -\kappa \sigma$ we obtain

$$\int_M \langle \Delta_L h, h \rangle_g dV_g = 2 \|\sigma\|_{\bar{g}}^2 \int_B \left(\mu - \frac{\kappa}{2} \right) f^{m-6} - \lambda f^{m-4} - 2f^{m-6} |\bar{\nabla} f|^2 dV_{\bar{g}}. \quad (4.2)$$

Multiplying Equation (2.1) by f^{m-6} and integrating we see that

$$\int_B \frac{\mu}{2} f^{m-6} - 2f^{m-6} |\bar{\nabla} f|^2 dV_{\bar{g}} = \frac{\lambda}{2} \int_B f^{m-4} dV_{\bar{g}}.$$

Hence if $\kappa < \mu$, we can substitute into Equation (4.2) and get the inequality

$$\int_M \langle \Delta_L h, h \rangle dV_g > -\lambda \|\sigma\|_g^2 \int_B f^{m-4} dV_{\bar{g}} = -\lambda \|h\|_g^2. \quad (4.3)$$

Hence the result follows. \square

We can now prove Corollary D.

Proof of Corollary D. Einstein products and Kähler-Einstein metrics with $h^{1,1} > 1$ (all with Einstein constant $\mu > 0$) admit divergence-free, trace-free eigentensors of the Lichnerowicz Laplacian with eigenvalue 0 (see [6], [8], and [18]). Equation (4.3) shows that a fibre-unstable warped product must also have a divergence-free, trace-free eigentensor satisfying the destabilising conditions and so must be unstable as a fibre. \square

5. THE PROOF OF THEOREM B

In [21] He, Petersen, and Wylie introduced the following function ρ and symmetric $(0, 2)$ -tensor P associated to a quasi-Einstein metric (B^n, \bar{g}, f, m) solving Equation (2.2):

$$\rho = \frac{1}{m-1} ((n-1)\lambda - \text{scal}(\bar{g})),$$

$$P = \text{Ric}(\bar{g}) - \rho \bar{g},$$

where $\text{scal}(\bar{g})$ is the scalar curvature of \bar{g} . Using the fact that

$$\text{Ric}(\bar{g}) = m f^{-1} \bar{\nabla}^2 f + \lambda \bar{g},$$

and

$$\text{scal}(\bar{g}) = m f^{-1} \bar{\Delta} f + n\lambda,$$

we can write the tensor fP as

$$fP = m \bar{\nabla}^2 f + \frac{m}{m-1} (\bar{\Delta} f + \lambda f) \bar{g}.$$

If the quasi-Einstein metrics appear as a family $(B^n, \bar{g}_i, f_i, m_i)$ with $\bar{g}_i \rightarrow \bar{g}_\infty$ and $m_i \rightarrow \infty$ as $i \rightarrow \infty$, then the tensors $P_i \rightarrow \text{Ric}(\bar{g}_\infty)$. As we saw in Proposition 2.6, the Ricci tensor of a gradient soliton plays an important role in the destabilising of product solitons. Hence we expect the tensor P to play a similar role for warped product Einstein metrics. As we shall see, it is the tensor fP that is divergence-free and so we have the following definition.

Definition 5.1 (Ricci variation). Let $(B^n \times F^m, \bar{g} \oplus f^2 \tilde{g})$ be a warped product Einstein manifold with Einstein constant λ . Then the Ricci variation is the tensor

$$h = \left(m \bar{\nabla}^2 f + \frac{m}{m-1} (\bar{\Delta} f + \lambda f) \bar{g} \right) \oplus 0. \quad (5.1)$$

Lemma 5.2. *The Ricci variation (5.1) is divergence-free.*

Proof. We begin by noting if $\bar{\eta} \in s^2(TB)$ satisfies $\bar{\text{div}}(\bar{\eta}) = 0$, then using Lemma 3.3,

$$\eta = f^{-m} \bar{\eta} \oplus 0,$$

satisfies $\text{div}(\eta) = 0$. In Proposition 5.4 of [21], the authors showed that $\bar{\text{div}}(f^{m+1}P) = 0$ and so $h = fP \oplus 0$ satisfies $\text{div}(h) = 0$. \square

The hypothesis of Theorem B will be used to guarantee that the functions f_i have certain limiting behaviours. We collect what we will need in a lemma. Note that (B, \bar{g}_i, f_i, m_i) converging the C^∞ topology to a non-trivial Ricci soliton $(B, \bar{g}_\infty, \varphi)$ means $g_i \rightarrow g_\infty$, $f_i^{m_i} \rightarrow e^{-\varphi}$, and $m_i \rightarrow \infty$ as $i \rightarrow \infty$ where the convergence is uniform with respect to any C^k -norm.

Lemma 5.3. *Let (B, \bar{g}_i, f_i, m_i) be a one parameter family of quasi-Einstein metrics that converge in the C^∞ topology to a non-trivial Ricci soliton $(B, \bar{g}_\infty, \varphi)$ as in Theorem B. Then*

- (1) *The functions $f_i \rightarrow 1$ as $i \rightarrow \infty$.*
- (2) *The one-forms $m_i^{1/2} d \log f_i \rightarrow 0$ as $i \rightarrow \infty$.*
- (3) *The associated constants μ_i from Equation (2.1) satisfy $\mu_i \rightarrow \lambda$ as $i \rightarrow \infty$.*

Proof. The first two items follow trivially from the requirement that $f_i^{m_i} \rightarrow e^{-\varphi}$. Item (3) is proved by Case in [10] Proposition 4.11. \square

We remark that the normalisation of the limiting soliton potential φ is actually fixed by $\lim_{i \rightarrow \infty} m_i(\lambda - \mu_i)$. This is also proved by Case in Proposition 4.11 of [10]. Next, we compute the stability operator N applied to the Ricci variation.

Lemma 5.4. *Let $(B^n \times F^m, \bar{g} \oplus f^2 \tilde{g})$ be a warped product Einstein manifold with λ and μ as in equation (2.1) and let h be the Ricci variation (5.1). Then*

$$\left\langle \frac{1}{2} \Delta h + \text{Rm}(h, \cdot), h \right\rangle_{L^2(g)} = \text{Vol}(F) \left(\lambda \|h\|_{L^2(f^m dV_{\bar{g}})}^2 + \frac{m}{m-1} \left\langle \frac{1}{2} \Delta(\bar{\Delta} f + \lambda f) \bar{g} + m \mu (f^{-3} df \otimes df), h \right\rangle_{L^2(f^m dV_{\bar{g}})} \right).$$

Proof. For an arbitrary function Φ we recall that in coordinates

$$\nabla^2 \Phi_{AB} = \frac{\partial^2 \Phi}{\partial x_A \partial x_B} - \Gamma_{AB}^C \frac{\partial \Phi}{\partial x_C}.$$

Hence we can use the formula

$$\Gamma_{\alpha\beta}^c = -\tilde{g}_{\alpha\beta} f \tilde{g}^{cd} (\bar{\nabla}_d f),$$

from Lemma 3.1 to obtain

$$\nabla^2 \Phi = \bar{\nabla}^2 \Phi \oplus f \langle \bar{\nabla} \Phi, \bar{\nabla} f \rangle \tilde{g}.$$

Using this, and the fact that $g = \bar{g} \oplus f^2 \tilde{g}$, we can write the tensor h in the following manner,

$$h = m \nabla^2 f + \frac{m}{m-1} (\bar{\Delta} f + \lambda f) g - \frac{m}{m-1} (f^2 \bar{\Delta} f + \lambda f^3 + (m-1) f |\bar{\nabla} f|^2) \tilde{g}.$$

Thus equation (2.1) yields $f^2 \bar{\Delta} f + \lambda f^3 + (m-1) f |\bar{\nabla} f|^2 = f \mu$ and so

$$h = m \nabla^2 f + \frac{m}{m-1} (\bar{\Delta} f + \lambda f) g - \frac{\mu m f}{m-1} \tilde{g}.$$

We will now compute the integral $\langle \frac{1}{2} \Delta h + \text{Rm}(h, \cdot), h \rangle_{L^2(g)}$ term-by-term. Let

$$\mathcal{T}_1 = m \nabla^2 f, \quad \mathcal{T}_2 = \frac{m}{m-1} (\bar{\Delta} f + \lambda f) g \quad \text{and} \quad \mathcal{T}_3 = \frac{\mu m f}{m-1} \tilde{g}.$$

We begin by computing $\langle N(\mathcal{T}_1), h \rangle_{L^2(g)}$. For an arbitrary function Φ , we note the identity

$$\langle \nabla^2 \Phi, H \rangle_{L^2(g)} = \langle \operatorname{div}^*(\nabla \Phi), H \rangle_{L^2(g)} = \langle \nabla \Phi, \operatorname{div}(H) \rangle_{L^2(g)},$$

for any $H \in TM^* \otimes TM^*$. Hence, as it is divergence-free by Lemma 5.2, h is L^2 -orthogonal to the Hessian of any function. In the proof of their Lemma 3.5 in [7], Cao–He show that for an Einstein metric

$$\frac{1}{2} \Delta \nabla^2 \Phi + \operatorname{Rm}(\nabla^2 \Phi, \cdot) = \nabla^2 \Delta \Phi + \lambda \nabla^2 \Phi,$$

for any function Φ . Applying these two identities to \mathcal{T}_1 we obtain

$$\left\langle \frac{1}{2} \Delta(m \nabla^2 f) + \operatorname{Rm}(m \nabla^2 f, \cdot), h \right\rangle_{L^2(g)} = \langle m \nabla^2(\Delta f + \lambda f), h \rangle_{L^2(g)} = 0 = \lambda \langle m \nabla^2 f, h \rangle_{L^2(g)}.$$

Thus

$$\left\langle \frac{1}{2} \Delta \mathcal{T}_1 + \operatorname{Rm}(\mathcal{T}_1, \cdot), h \right\rangle_{L^2(g)} = \lambda \langle \mathcal{T}_1, h \rangle_{L^2(g)}.$$

To compute the integral for the second term \mathcal{T}_2 we note that

$$\Delta(\Phi g) = (\Delta \Phi)g,$$

for any function Φ and any Riemannian metric g . We also have

$$\operatorname{Rm}(\Phi g, \cdot) = \lambda \Phi g,$$

for any function Φ any Einstein metric g with Einstein constant λ . Hence we conclude

$$\left\langle \frac{1}{2} \Delta \mathcal{T}_2 + \operatorname{Rm}(\mathcal{T}_2, \cdot), h \right\rangle_{L^2(g)} = \lambda \langle \mathcal{T}_2, h \rangle_{L^2(g)} + \frac{m}{m-1} \left\langle \frac{1}{2} \Delta(\bar{\Delta} f + \lambda f) \right\rangle g.$$

To deal with the final term \mathcal{T}_3 we note that pointwise $\langle \tilde{g}, h \rangle = 0 = \langle \mathcal{T}_3, h \rangle$. This is clear using the original definition of h in Equation (5.1). Using Lemma 3.4 and Lemma 3.5 we obtain

$$\left\langle \frac{1}{2} \Delta(f \tilde{g}) + \operatorname{Rm}(f \tilde{g}, \cdot), h \right\rangle_{L^2(g)} = \langle m(f^{-3} df \otimes df - f^{-2} \bar{\nabla}^2 f), h \rangle_{L^2(g)}.$$

Hence

$$\left\langle \frac{1}{2} \Delta \mathcal{T}_3 + \operatorname{Rm}(\mathcal{T}_3, \cdot), h \right\rangle_{L^2(g)} = \lambda \langle \mathcal{T}_3, h \rangle_{L^2(g)} + \frac{m\mu}{m-1} \left\langle (m(f^{-3} df \otimes df - f^{-2} \bar{\nabla}^2 f), h) \right\rangle_{L^2(g)}.$$

Putting the three calculations for the \mathcal{T}_i together we see that h is almost a λ -eigentensor for the stability operator N .

$$\begin{aligned} \left\langle \frac{1}{2} \Delta h + \operatorname{Rm}(h, \cdot), h \right\rangle_{L^2(g)} &= \lambda \|h\|_{L^2(g)}^2 + \frac{m}{m-1} \left\langle \frac{1}{2} \Delta(\bar{\Delta} f + \lambda f) \bar{g} \right. \\ &\quad \left. - m\mu(f^{-3} df \otimes df - f^{-2} \bar{\nabla}^2 f), h \right\rangle_{L^2(g)}. \end{aligned} \quad (5.2)$$

We now observe two further identities for an arbitrary function $\Phi \in C^\infty(B)$, firstly the general identity

$$\nabla^2 \Phi^{-1} = -\frac{1}{\Phi^2} \nabla^2 \Phi + \frac{2}{\Phi^3} d\Phi \otimes d\Phi.$$

The second is that, as h is both divergence-free and pointwise orthogonal to tensors of the form $0 \oplus \tilde{h}$, we have

$$0 = \langle \nabla^2 \Phi, h \rangle_{L^2(g)} = \langle \bar{\nabla}^2 \Phi \oplus f \langle \bar{\nabla} \Phi, \bar{\nabla} f \rangle \tilde{g}, h \rangle_{L^2(g)} = \langle \bar{\nabla}^2 \Phi, h \rangle_{L^2(g)}.$$

Hence equation (5.2) becomes

$$\left\langle \frac{1}{2} \Delta h + \text{Rm}(h, \cdot), h \right\rangle_{L^2(g)} = \lambda \|h\|_{L^2(g)}^2 + \frac{m}{m-1} \left\langle \left(\frac{1}{2} \Delta (\bar{\Delta} f + \lambda f) \bar{g} + m \mu (f^{-3} df \otimes df) \right), h \right\rangle_{L^2(g)}.$$

The result follows by computing the integrals over the base B and fibre F . \square

We can now prove Theorem B.

Proof of Theorem B. Using Lemma (2.4), we see that the warped product $(B^n \times F^{m_i}, \bar{g}_i \oplus f_i^2 \tilde{g})$ is unstable if

$$\left\langle \frac{1}{2} \Delta h_i + \text{Rm}(h_i, \cdot), h_i \right\rangle_{L^2(g_i)} - \lambda \frac{\left(\int_M \text{tr}(h_i) dV_{g_i} \right)^2}{(n + m_i) \text{Vol}(M)} > 0,$$

where the h_i are the Ricci variations defined by the metrics \bar{g}_i , the functions f_i and the constants m_i . Thus we consider the behaviour, as $i \rightarrow \infty$ of the sequence

$$a_i = \text{Vol}(F_i)^{-1} \left(\left\langle \frac{1}{2} \Delta h_i + \text{Rm}(h_i, \cdot), h_i \right\rangle_{L^2(g_i)} - \lambda \frac{\left(\int_M \text{tr}(h_i) dV_{g_i} \right)^2}{(n + m_i) \text{Vol}(M)} \right).$$

The sequence a_i does not in fact depend at all on the fibres F_i .

Using Lemma 5.4 we need to consider the limit as $i \rightarrow \infty$ of

$$\lambda \|h_i\|_{L^2(\bar{g}_i)}^2 + \frac{m_i}{m_i - 1} \left\langle \left(\frac{1}{2} \Delta (\bar{\Delta} f_i + \lambda f_i) \bar{g}_i + m_i \mu (f_i^{-3} df_i \otimes df_i) \right), h_i \right\rangle_{L^2(\bar{g}_i)}.$$

As Lemma 5.3 gives $f_i \rightarrow 1$, $d \log f_i^{m_i^{1/2}} \rightarrow 0$, and $\mu_i \rightarrow \lambda$ we see

$$\left\langle \left(\frac{1}{2} \Delta (\bar{\Delta} f_i + \lambda f_i) \bar{g}_i + m_i \mu_i (f_i^{-3} df_i \otimes df_i) \right), h_i \right\rangle_{L^2(f_i^{m_i} dV_{\bar{g}_i})} \rightarrow 0.$$

We also have

$$\text{Vol}(F_i)^{-1} \lambda \frac{\left(\int_M \text{tr}(h_i) dV_{g_i} \right)^2}{(m_i + n) \text{Vol}(M)} = \lambda \frac{\left(\int_B \text{tr}(h_i) f_i^{m_i} dV_{\bar{g}_i} \right)^2}{(m_i + n) \text{Vol}(B, f^{m_i} dV_{\bar{g}_i})} \rightarrow 0.$$

The limit of the a_i is thus

$$\lambda \|h_\infty\|_{L^2(e^{-\varphi} dV_{\bar{g}_\infty})}^2,$$

where $h_\infty = \text{Ric}(\bar{g}_\infty)$ is the limit of the h_i as $i \rightarrow \infty$ (the limit exists as g_i converges in the C^∞ topology). Hence we see for large enough i ,

$$\int_M \langle N(h_i + cg_i), h_i + cg_i \rangle dV_g > 0,$$

and the result follows. \square

REFERENCES

- [1] BATAT, W., HALL, S. J., JIZANY, A., AND MURPHY, T. Conformally Kähler geometry and quasi-Einstein metrics. *Münster Journal of Mathematics* 8 (2015), 211–228.
- [2] BESSE, A. L. *Einstein manifolds*. Classics in Mathematics. Springer-Verlag, Berlin, 2008. Reprint of the 1987 edition.
- [3] BÖHM, C. Inhomogeneous Einstein metrics on low-dimensional spheres and other low-dimensional spaces. *Invent. Math.* 134, 1 (1998), 145–176.
- [4] CAO, H.-D. Existence of gradient Kähler-Ricci solitons. In *Elliptic and parabolic methods in geometry (Minneapolis, MN, 1994)*. A K Peters, Wellesley, MA, 1996, pp. 1–16.
- [5] CAO, H.-D. Recent progress on Ricci solitons. In *Recent advances in geometric analysis*, vol. 11 of *Adv. Lect. Math. (ALM)*. Int. Press, Somerville, MA, 2010, pp. 1–38.

- [6] CAO, H.-D., HAMILTON, R., AND ILMANEN, T. Gaussian densities and stability for some Ricci solitons. - (2004). preprint, arXiv:math/0404165 [math.DG].
- [7] CAO, H.-D., AND HE, C. Linear stability of Perelman's ν -entropy on symmetric spaces of compact type. *J. Reine Angew. Math.* 709 (2015), 229–246.
- [8] CAO, H.-D., AND ZHU, M. On second variation of Perelman's Ricci shrinker entropy. *Math. Ann.* 353, 3 (2012), 747–763.
- [9] CASE, J., SHU, Y.-J., AND WEI, G. Rigidity of quasi-Einstein metrics. *Differential Geom. Appl.* 29, 1 (2011), 93–100.
- [10] CASE, J. S. Smooth metric measure spaces and quasi-Einstein metrics. *Internat. J. Math.* 23, 10 (2012), 1250110, 36.
- [11] CASE, J. S. The energy of a smooth metric measure space and applications. *J. Geom. Anal.* 25, 1 (2015), 616–667.
- [12] DAI, X., WANG, X., AND WEI, G. On the stability of Riemannian manifold with parallel spinors. *Invent. Math.* 161, 1 (2005), 151–176.
- [13] DANCER, A., AND WANG, M. On Ricci solitons of cohomogeneity one. *Ann. Global Anal. Geom.* 39 (2011), 259–292.
- [14] GIBBONS, G. W., AND HARTNOLL, S. A. Gravitational instability in higher dimensions. *Phys. Rev. D (3)* 66, 6 (2002), 064024, 17.
- [15] GIBBONS, G. W., HARTNOLL, S. A., AND POPE, C. N. Bohm and Einstein-Sasaki metrics, black holes, and cosmological event horizons. *Phys. Rev. D (3)* 67, 8 (2003), 084024, 24.
- [16] HALL, S., HASLHOFER, R., AND SIEPMANN, M. The stability inequality for Ricci-flat cones. *J. Geom. Anal.* 24, 1 (2014), 472–494.
- [17] HALL, S. J. Quasi-Einstein metrics on hypersurface families. *J. Geom. Phys.* 64 (2013), 83–90.
- [18] HALL, S. J., AND MURPHY, T. On the linear stability of Kähler-Ricci solitons. *Proc. Amer. Math. Soc.* 139, 9 (2011), 3327–3337.
- [19] HALL, S. J., AND MURPHY, T. On the spectrum of the Page and the Chen-LeBrun-Weber metrics. *Ann. Global Anal. Geom.* 46, 1 (2014), 87–101.
- [20] HALL, S. J., AND MURPHY, T. Numerical approximations to extremal toric Kähler metrics with arbitrary Kähler class. *Proc. Edinb. Math. Soc. (2)* 60, 4 (2017), 893–910.
- [21] HE, C., PETERSEN, P., AND WYLIE, W. On the classification of warped product Einstein metrics. *Comm. Anal. Geom.* 20, 2 (2012), 271–311.
- [22] KIM, D.-S., AND KIM, Y. H. Compact Einstein warped product spaces with nonpositive scalar curvature. *Proc. Amer. Math. Soc.* 131, 8 (2003), 2573–2576.
- [23] KOISO, N. On rotationally symmetric Hamilton's equation for Kähler-Einstein metrics. In *Recent topics in differential and analytic geometry*, vol. 18 of *Adv. Stud. Pure Math.* Academic Press, Boston, MA, 1990, pp. 327–337.
- [24] KRÖNCKE, K. Stable and unstable Einstein warped products. *Trans. Amer. Math. Soc.* 369, 9 (2017), 6537–6563.
- [25] KRÖNCKE, K. Stability of sin-cones and cosh-cylinders. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 18, 3 (2018), 1155–1187.
- [26] KRÖNCKE, K. Stability of Einstein metrics under Ricci flow. *Comm. Anal. Geom.* (to appear). arXiv:1312.2224 [math.DG].
- [27] LÜ, H., PAGE, D. N., AND POPE, C. N. New inhomogeneous Einstein metrics on sphere bundles over Einstein-Kähler manifolds. *Phys. Lett. B* 593, 1-4 (2004), 218–226.
- [28] PALI, N. The soliton-Ricci flow with variable volume forms. *Complex Manifolds* 3 (2016), Art. 3.
- [29] PALI, N. Variational stability of Kähler-Ricci solitons. *Adv. Math.* 290 (2016), 15–35.
- [30] PERELMAN, G. The entropy formula for the Ricci flow and its geometric applications. - (2002). preprint, arXiv:math/0211159 [math.DG].
- [31] PERELMAN, G. Finite extinction time for the solutions to the Ricci flow on certain three-manifolds. - (2003). preprint, arXiv:math/0307245 [math.DG].
- [32] PERELMAN, G. Ricci flow with surgery on three-manifolds. - (2003). preprint, arXiv:math/0303109 [math.DG].
- [33] SESUM, N. Linear and dynamical stability of Ricci-flat metrics. *Duke Mathematical Journal* 133, 1 (2006), 1–26.

ECOLE NATIONALE POLYTECHNIQUE D'ORAN, B.P 1523 EL M'NAOUAR, 31000 ORAN, ALGERIA
E-mail address: `batatwafa@yahoo.fr`

SCHOOL OF MATHEMATICS AND STATISTICS, HERSCHEL BUILDING, NEWCASTLE UNIVERSITY, NEWCASTLE-
UPON-TYNE, NE1 7RU
E-mail address: `stuart.hall@ncl.ac.uk`

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY FULLERTON, 800 N. STATE COLLEGE
BLD., FULLERTON, CA 92831, USA.
E-mail address: `tmurphy@fullerton.edu`