

The definitive version of this article is published and available online as:

Devereux, C.L., Denny, M.J.H., Whittingham, M.J., Minimal effects of wind turbines on the distribution of wintering farmland birds, *Journal of Applied Ecology*, Volume 45, Issue 6, December 2008, Pages 1689-1694
<http://www3.interscience.wiley.com/journal/121427776/abstract>

Title: Minimal effects of wind turbines on the distribution of wintering farmland birds

Claire L. Devereux*, Matthew J.H. Dennyⁿ & Mark J. Whittingham*[&].

* *School of Biology, Ridley Building, Newcastle University, Newcastle-Upon-Tyne, NE1 7RU*

ⁿ *Baker Shepherd Gillespie, Worton Rectory Park, Oxford, OX29 4SX*

[‡] *Present address: Denny Ecology, Vine Cottage, Hailey, Witney, Oxon, OX29 9UB*

[&] *Corresponding author: e-mail: m.j.whittingham@ncl.ac.uk*

Short title: windfarms and farmland birds

No of words in abstract: 299

No of words in text: 5180

[&] *Correspondance author: Dr Mark Whittingham, School of Biology, Ridley Building, Newcastle University, Newcastle-Upon-Tyne, NE1 7RU. E-mail: m.j.whittingham@ncl.ac.uk, tel: 0191 2226599, fax 0191 2225229*

Summary

1. Energy production from wind power is increasing rapidly in Europe to help combat the threats from global warming. For example, the European Commission have set a target for 20% of EU energy to come from renewable sources by 2020. In recent decades biodiversity on European farmland has fallen dramatically due to agricultural intensification. Agri-environment schemes (AES) have been implemented across the EU, in part at least, to combat these declines. Significant numbers of turbines are, and will be, built on farmland. There is therefore a potential conflict between wind turbines and AESs on farmland.
2. Various mechanisms potentially cause wind turbines to alter bird distribution including noise and physical structure.
3. We show that turbine location (controlling for other effects such as boundary location and crop type) did not affect the distribution of four functional groups of wintering farmland birds (seed-eaters, corvids, gamebirds and Eurasian skylarks) at differing distances from wind turbines ranging from 0-150 m to 600-750 m. The only species for which distribution was related to the presence of wind turbines was the largest and least manoeuvrable (common pheasant *Phasianus colchicus* L.).
4. In a further analysis of data collected at 0-75 m and 75-150 m from turbines we found no evidence to suggest that farmland birds in our study avoided areas close to wind turbines.
5. *Synthesis and applications.* This is the first evidence suggesting that the present and future location of large numbers of wind turbines on European farmland is unlikely to have detrimental effects on farmland birds (at least for those species included in our study). This should be welcome news for nature conservationists, wind energy companies and policy-makers. However, our work is only a first step as there may be potential influences of wind turbines on bird distribution during the breeding season.

Keywords: climate change; disturbance; sustainable agriculture; farmland biodiversity; agri-environment schemes; noise pollution; wind energy

Introduction

Global climate change has resulted in worldwide calls for 'renewable' energy sources to reduce carbon emissions. Wind power provides the largest share in the renewable energy sector in Europe. The Global Wind Energy Council is forecasting that the global wind market will grow by over 155% to reach 240 Giga-Watts of total installed capacity by 2012 (<http://www.gwec.net/>). The European Commission have set a target for 20% of EU energy to come from renewable sources by 2020 (EC proposal 52008PC0019). The major land use within the European Union is agriculture (44% of land area was categorised as farmed in 1997 with woodland the next largest landholding area at 32% – EC website:

http://ec.europa.eu/agriculture/envir/report/en/terr_en/report.htm), and so it seems likely that many of these turbines will be located on farmland. For example, in the UK it is predicted that approximately 10% of UK electricity production by 2020 will come from onshore wind power sites (<http://www.bwea.com>); from interpolation using current technology parameters (c.2.5MW turbines), this equates to approximately 4400 new turbines. Within the EU a large amount of resources are given over to agri-environment schemes designed to enhance the environment on farmland (e.g. between 1993 and 2003 €24 Billion was spent by the EU on agri-environment schemes; Kleijn & Sutherland 2003). One of the major goals of these schemes is to enhance biodiversity on farmland (Whittingham 2007), which has been widely reported as declining (e.g. Krebs *et al.* 1999; Donald, Green & Heath 2001).

There are major concerns regarding the impacts of wind turbines on animals, especially birds (e.g. Percival 2005; Osborn *et al.* 2000), but other taxa such as bats have also been shown to be affected (e.g. Arnett *et al.* 2008). Much terrestrial research into the effects of wind turbines on birds has focussed on geese, waders and raptors: all groups that are likely to suffer relatively high potential direct mortality risk (Barrios & Rodriguez 2004; De Lucas *et al.* 2004; Percival 2005). In the UK, raptor, wader and wintering goose populations are highest in upland and coastal areas, and there is increasing conservation concern regarding the impact of windfarms on these species in these areas. Consequently applications to build new turbines are increasingly focussing on other sites, particularly lowland farmland landscapes in central and eastern England (see British Wind Energy Association website, <http://www.bwea.com/ukwed/>, which shows maps of existing and future windfarm sites).

It is not uncommon for farmland bird species to fly at altitudes above 20 m (the bottom reach of the turbines in our study) but there is virtually no evidence-based evaluation in the published literature of the potential displacement that could result from wind turbines. One study from the USA found that songbird species living in a grassland/farmland mixture occurred at up to four times higher densities further than 180 m from wind turbines during the breeding season (Leddy, Higgins & Naugle 1999); however this study did not appear to control for differences in habitats at differing distances from wind turbines. We could find no evidence from the European literature of the effects on songbirds during the winter. Farmland bird populations have declined severely across Europe, e.g. a 40% decline between 1980 and 2005 across 20 EU countries

(<http://www.ebcc.info/index.php?ID=299>). Consequently the population index of farmland bird species across Europe was recently adopted by the European Union as a structural and sustainable development indicator.

Should wind turbines affect the distribution of farmland birds then it would be logical to spatially separate agri-environmental measures aimed at birds and the location of wind turbines. There is thus a potential conflict between uptake of agri-environment schemes and the placement of wind turbines on lowland farmland in the UK. This is likely to apply to many other European countries in which farmland birds are also a major target of agri-environment schemes (Kleijn & Sutherland 2003).

We tested whether wind turbines affect the distribution of lowland farmland birds during the winter, a critical time in the annual cycle (Payne & Wilson 1999). To our knowledge our study is the first to address the issue of whether wind turbines located on lowland agricultural fields effect the distribution (and hence habitat quality) of European farmland birds. *A priori* the most likely mechanism of altering habitat quality is via disturbance. Birds might perceive wind turbines as disturbance for a number of reasons. Increased noise levels have been shown to affect bird distribution and behaviour in a variety of habitats (e.g. Reijnen *et al.* 1995; Reijnen, Foppen & Meeuwsen 1996; Habib, Bayne & Boutin 2007) and within cities this has been shown to be linked with decreased ability to communicate vocally (e.g. Katti & Warren 2004). Other work has shown that increased noise levels are associated with higher predation risk by birds (Quinn *et al.* 2006) which is likely to result in changing patch quality (e.g. Butler *et al.* 2005). Wind turbines create noise and the amount varies with wind speed or rpm (rotations per minute). The noise created by a 3MW turbine can be about 100 dB at 14 rpm and 108 dB at 20 rpm (Klug 2002), although more modern wind turbines emit less noise. Work on farmland has shown that bird distribution is affected at levels of noise higher than around 40-50dB (Reijnen *et al.* 1996). It is thus plausible that noise from wind turbines could result in lower habitat quality close to wind turbines (e.g. Habib *et al.* 2007). Alternatively: (i) birds might avoid flying near to turbines to avoid potential collision, resulting in an effective 'exclusion' zone around turbines; (ii) human presence around turbines could affect birds, although on lowland English farmland this seems unlikely given the amount of disturbance by farmers, walkers, etc.; (iii) birds that prefer 'open' landscapes might avoid tall structures irrespective of potential collision risk. In general, disturbance reduces habitat quality for birds and thus, all other things being equal, birds may preferentially use patches away from wind turbines as predicted by Optimal Foraging Theory (Sutherland 1996).

Methods

Bird survey data were collected from farmland immediately surrounding two operational wind-farms in East Anglia in January and February 2007. Each had eight turbines arranged in two parallel rows separated by 900 m (four and four at Site 1; three and five at Site 2) and spaced at 300 m intervals. The turbines were all Repower MM82 2.0 MW tri-blade type with a hub height of 60 m and total tip height of 100 m. Both sites have been operational since early 2006 and the wind turbines were all 'turning' throughout our surveys.

Many factors are known to affect the distribution of farmland birds, e.g. crop type, non-cropped habitats (such as grass margins, hedge dimensions, trees within hedges, woodland, etc.) and spraying of pesticides (Hinsley & Bellamy 2000; Whittingham & Evans 2004; Whittingham *et al.* in press). Our study sites were relatively homogenous in nature, comprising cereal fields (47%; mostly winter wheat), oil seed rape (23%; patchily distributed and similar visually to bare plough) and bare plough (30%), but with variation in the distribution of ditches and hedges.

The windfarms were located on fenland arable farmland managed by five landowners. Maps of each site were overlaid with 150 m grids after bird counts had been undertaken. Each 150 x 150 m grid block was scored for its distance to the nearest turbine as follows: <150 m, 150-300 m, 300-450 m, 450-600 m, 600-750 m. No blocks greater than 750 m from a turbine were surveyed because of access issues. The predominant crop type and presence of boundary features (hedge or vegetated ditch) was recorded for each block (Table 1). We took the proportion of winter wheat as an indicator of crop type (% cereals in Table 1). Both sites were flat (no slopes >2 degrees).

Eleven bird survey visits were made to the two sites (five at Site 1 and six at Site 2). Surveys were performed at least two days apart between 0900 and 1500 h, at least one hour after dawn and before dusk (to avoid times when birds may have either departed to roost or before they had arrived from roosts). Surveys were not conducted in reduced visibility conditions such as mist, fog and persistent rain or when turbines were not operating (e.g. at very low wind speeds). Before each survey, a route of line transects was planned, covering a distance of approximately 7.5 km with approximately a third of the blocks in each distance category (thus 82.5 km of transects were walked in total). Different combinations of blocks and different routes were selected at random for each survey. The routes were walked at a speed of 2.5–3 kmh⁻¹. All birds flushed by the observer and birds within c100m were identified by sight and sound, and individual positions marked onto a map. The large majority of birds were recorded by flushing and care was taken to avoid duplicate counts of these birds through observations of movements of previously flushed individuals.

We analysed data for those bird species that were recorded in sufficient abundance on their own, and for other species (where appropriate) we assigned them to functional groups based on ecological and taxonomic characteristics (Butler *et al.* 2005) (see Table S1, Supplementary Material, for species assignment to functional groups). Note that common species names throughout this paper follow Gill & Wright (2006) and associated updates (<http://www.worldbirdnames.org/index.html>). The effect of turbine proximity (distance) on the abundance and distribution of the four species/functional groups was tested using Generalized Linear Mixed Models (GLMMs) under a binomial distribution using a logit link function in Genstat v8 (VSN International). Distance was specified as a continuous variable, but the results did not change when the analysis was repeated with distance considered as categorical. Due to the logistics of each site it was not possible to include the same number of blocks in each 'distance from turbine' category in each visit (see Table S2, Supplementary Material). Therefore, we controlled for this variation in our analysis by including the number of blocks visited in each distance category during each survey as the binomial

denominator and the number of blocks containing each functional group as the response variable. For example, if granivores were present in 4 blocks out of 14 visited in the 150-300 m distance category on a survey then 4 was specified as the response variable and 14 as the denominator (thus 14 was equivalent to the number of 'trials' undertaken in a Binomial model). This method of abundance analysis represents a biologically realistic approach as birds in flocks are unlikely to select foraging habitats independently of conspecifics, but frequency of occurrence is often related to total number of individuals recorded, which is likely to indicate the relative value of a foraging site (see Perkins *et al.* 2000; Moorcroft *et al.* 2002).

A two-level factor 'site' was included as a random effect in all models to control for variation resulting from unmeasured site-specific parameters. To control for seasonal effects the term 'visit', which correlated with Julian date, was incorporated into the model and considered as a continuous variable. Crop type and presence of boundary features were standardized by the number of blocks visited in each distance category during each survey and were included as fixed effects (see Table 1).

The significance of distance, crop type and boundary were assessed using the change in deviance (ΔD), which is distributed asymptotically as χ^2 on removal of each term from a model including all of the predictors. Models were also re-run with visit excluded as it did not explain any of the variation in the distribution of most functional groups, and its exclusion gave very similar results (Table 2). Our approach was designed to test the effect of distance from turbines whilst checking that any result is not an artefact of some confounding effect caused by differences between crop type and boundary at varying distances from turbines. Thus we acknowledge there are more sensitive ways to test for the effects of boundaries and crop types on our data set but this was not the aim of our analysis. The fit of the model to the assumptions of a binomial distribution can be approximated by comparing the ratio of residual deviance / residual degrees of freedom (Crawley 1993). Ratios close to one indicate a reasonable fit to the data, whereas ratios greater than 2.5 indicate a poor, overdispersed fit (Crawley 1993). All probabilities quoted are two-tailed. Means and standard errors are presented in the form mean \pm 1 standard error.

Results

Thirty-three bird species were observed during surveys of the two windfarms totalling 2845 individuals (Table S2, Supplementary Material). Three additional species were seen flying over the site (western barn owl *Tyto alba* L., lesser black-backed gull *Larus fuscus* L. and black-headed gull *Larus ridibundus* L.). The proportions of each species at different distances from the wind turbines are given in Table S1, Supplementary Material.

Overall, for all functional groups there was no effect of turbine proximity on the probability of the functional group's occurrence in a block (Figure 1, Table 2). Granivores were more likely to occur in 150 x 150 m blocks that contained boundary features (e.g. hedges and vegetated ditches) but their occurrence was not related to turbine proximity ($\chi^2 = 0.26$, $P = 0.56$). Eurasian skylarks *Alauda arvensis* L. tended to avoid blocks containing boundary features ($\chi^2 = 3.82$, $P = 0.05$) but were not affected by turbine proximity ($\chi^2 =$

0.53, $P = 0.47$). Corvids were more likely to be found in blocks containing a higher proportion of cereal crops ($\chi^2 = 5.46$, $P = 0.02$) but were not affected by turbine proximity ($\chi^2 = 2.64$, $P = 0.10$). Gamebirds were unaffected by turbine proximity ($\chi^2 = 1.56$, $P = 0.21$) or any of the other predictors. However, when inspecting the raw data it was apparent that the two gamebird species responded differently. Turbine proximity had no effect on red-legged partridges *Alectoris rufa* L. ($\chi^2 = 0.24$, $P = 0.62$) but the probability of common pheasant *Phasianus colchicus* L. presence increased with increasing distance from turbines ($\chi^2 = 8.04$, $P = 0.005$) (see Figure 2).

One potential criticism of our study is that we did not look at distances less than 150 m from turbines. In order to address this issue we subdivided our data from survey maps into two further units. One category included only blocks within 75 m of a turbine and the other blocks between 75 and 150 m from a turbine. We then analysed the data in the same way as for the main data set with turbine distance included as a two level factor, site as a two-level factor and proportion of boundary and proportion of cereals both as variables (data are presented in Table S3, Supplementary material). There was no significant effect of turbine distance on granivores ($\chi^2 = 1.08$, d.f. = 1, $P = 0.35$) or gamebirds ($\chi^2 = 1.98$, d.f. = 1, $P = 0.19$). However, both corvid spp. ($\chi^2 = 4.39$, d.f. = 1, $P = 0.04$) and Eurasian skylarks ($\chi^2 = 8.41$, d.f. = 1, $P < 0.005$) were more likely to occur in blocks close to turbines than those further away.

Discussion

Overall our study suggests that wind turbines of the size we studied (2.0 MW, 60 m from ground level at hub height, 100 m at top of blade and 20 m at bottom of blade) had little impact on the distribution of farmland birds on our study sites during winter. To our knowledge this is the first examination of the effects of wind turbines on the distribution of lowland farmland birds.

There has been considerable interest in halting and reversing recent farmland bird population declines both in the media and by the UK and other EU governments (Fuller 2000; Vickery *et al.* 2004). Measures to effect this change both in the UK and more widely in the other EU countries (Sutherland & Kleijn 2003) have been implemented through government-funded agri-environment schemes (Whittingham 2007). Our work suggests that the benefits of agri-environment schemes for wintering farmland birds are unlikely to be compromised by the presence of wind turbines at least for those species in our study.

We recorded four species of granivorous farmland birds that are currently 'red' listed as of high conservation concern: yellowhammer *Emberiza citrinella* L., Eurasian tree sparrow *Passer montanus* L., common reed bunting *Emberiza schoeniclus* L. and corn bunting *Miliaria calandra* L. (Gregory *et al.* 2002). Although individual counts of some granivorous species were low, cumulatively we recorded 400 individual seed-eaters. Two pieces of evidence suggest that the distribution of granivorous birds is not affected by wind turbines. First, across the group as a whole (including seven species) we found no statistical relationship between the distance from turbines and occurrence. Secondly, none of the seven individual species (see Table S1, Supplementary Material) shows a pattern indicative of avoidance of areas close to turbines.

Visual inspection of individual species presence and abundance in each of the remaining three functional groups (see Tables S1, S2, Supplementary Material) also failed to reveal any patterns of avoidance of wind turbines. Inspection of the individual species counts not included within the four functional groups did reveal two species with low counts in the 0-150 m area. Interestingly both species (mallard *Anas platyrhynchos* L. and common wood pigeon *Columba palumbus* L.) were larger less manoeuvrable species; however, not too much should be inferred from their abundances (Table S2, Supplementary Material) given the low number of presences recorded (Table S1, Supplementary Material): in other words a few large flocks may have biased the results.

Our study did not assess the collision risk to farmland birds from wind turbines as has been reported for larger birds like raptors (Barrios & Rodriguez 2004). However, it seems likely that larger birds, which are generally less manoeuvrable in flight, are likely to be at higher risk of collision than small farmland birds. Two recent studies have both shown that ducks and geese make far less use of areas within several hundred metres of wind turbines at sea (Desholm & Kahlert 2005; Larsen & Guillemette 2007). The results of our study also show that the largest (and least manoeuvrable) species, common pheasant, which we were able to analyse, tended to use areas further from wind turbines less than would be expected by chance.

Our analysis suggests that turbines were not a major deterrent to birds using areas within 150 m and supports our main findings at a larger scale. The underlying reason for the increased use of areas close to turbines by corvid spp. and Eurasian skylarks is not clear although it may be related to food. Nevertheless it is clear that a range of farmland birds are not influenced by wind turbines even at close proximity.

A potential criticism of our study is whether the data collected has sufficient power to detect effects of turbines. This is difficult to refute comprehensively but several points suggest this is not the case. First, the data on the effect of turbine distance on common pheasants was significant but the sample size was relatively small compared to other species/ groups included in the analysis. Secondly, the results showing that both Eurasian skylark and corvids preferred areas close to turbines at small spatial scales were statistically significant: thus even within one distance category sufficient data existed to pick up a significant result. Thirdly, inspection of the raw data (see Supplementary Material) did not reveal any consistent patterns of turbine avoidance across species/species groups.

With large numbers of windfarms needing to be built on lowland areas, the cumulative impacts on farmland bird species has the potential to be a significant constraint to development. At the European level both the Environmental Impact Assessment (EIA) and Strategic Environmental Assessment (SEA) Directives (and the associated UK EIA regulation – Town and Country Planning (EIA) Regulations (1999)) specify the need to assess the cumulative environmental impacts of development projects. Our study has provided a first step in assessing the potential impact of wind turbines on farmland birds. Our study suggests, at least during the winter period, that there is likely to be little impact of wind turbines on the farmland bird species included in our study and this should be welcome news for nature conservationists, wind energy companies and policy-

makers. However, we recommend that further studies of the effects of wind turbines on farmland birds, particularly during the breeding season, are carried out.

Acknowledgements

Many thanks to Baker Shepherd Gillespie Ecological Consultants for providing the funding to carry out this study and to BBSRC for funding MJW through a David Phillips Fellowship. Thanks to the landowners for allowing access to the windfarm sites and to Chris Elphick, Davy McCracken and two anonymous referees for providing helpful suggestions to improve the MS.

References

Arnett, E.B., Kent Brown, W., Erikson, W.P., Fiedler, J.K., Hamilton, B.L., Henry, T.H., Jain, A., Johnson, G.D., Kerns, J., Koford, R.R., Nicholson, C.P., O'Connell, T.J., Piorkowski, M.D. & Tankersley, J.R. (2008) Patterns of bat fatalities at wind energy facilities in North America. *Journal of Wildlife Management*, **72**, 61-78.

Barrios, L. & Rodriguez, A. (2004) Behavioural and environmental correlates of soaring-bird mortality at on-shore wind turbines. *Journal of Applied Ecology*, **41**, 72-81.

Butler, S.J., Bradbury, R.B. & Whittingham, M.J. (2005) Stubble height manipulation causes differential spatial use of stubble fields by farmland birds. *Journal of Applied Ecology*, **42**, 469-476.

Crawley, M.J. (1993) *GLIM for Ecologists*. Blackwell Scientific Publications, Oxford, UK.

De Lucas, M., Janss, G.F.E. & Ferrer, M. (2004) The effects of a wind farm on birds in a migration point: the Strait of Gibraltar. *Biodiversity and Conservation*, **13**, 395-407.

Desholm, M. & Kahlert, J. (2005) Avian collision risk at an offshore windfarm. *Biology Letters*, **1**, 296-298.

Donald, P.F., Green, R.E. & Heath, M.F. (2001) Agricultural intensification and the collapse of Europe's

farmland bird populations. *Proceedings of the Royal Society, Series B.*, **268**, 25-29.

Fuller, R.J. (2000) Relationships between recent changes in lowland British agriculture and farmland bird populations: an overview. *Ecology and Conservation of Lowland Farmland Birds* (eds N.J. Aebischer, A.D. Evans, P.V. Grice, & J.A. Vickery,) pp. 5-16. British Ornithologists' Union, Tring.

Gill, F. & Wright, M. (2006) *Birds of the World: recommended English names*. Princeton University Press, Princeton, USA.

Gregory, R.D., Wilkinson, N.I., Noble, D.G., Robinson, J.A., Brown, A.F., Hughes, J., Procter, D.A., Gibbons, D.W. & Galbraith, C.A. (2002) The population status of birds in the United Kingdom, Channel Islands and Isle of Man: an analysis of conservation concern 2002-2007. *British Birds*, **95**, 410-450.

Habib, L., Bayne, E.M. & Boutin, S. (2007) Chronic industrial noise affects pairing success and age structure of ovenbirds *Seiurus aurocapilla*. *Journal of Applied Ecology*, **44**, 176-184.

Hinsley, S.A. & Bellamy, P.E. (2000) The influence of hedge structure, management and landscape context on the value of hedgerows to birds: A review. *Journal of Environmental Management*, **60**, 33-49.

Katti, M. & Warren, P.S. (2004) Tits, noise and bioacoustics. *Trends in Ecology and Evolution*, **19**, 109-110.

Kleijn, D. & Sutherland, W.J. (2003) How effective are European agri-environment schemes in conserving and promoting biodiversity? *Journal of Applied Ecology*, **40**, 947-969.

Klug, H. (2002) Noise from wind turbines: standards and noise reduction procedures. Paper from the Forum Acousticum 2002, Sevilla, Spain.

- Krebs, J.R., Wilson, J.D., Bradbury, R.B. & Siriwardena, G.M. (1999). The second silent spring? *Nature*, **401**, 611-612.
- Larsen, J.K. & Guillemette, M. (2007) Effects of wind turbines on flight behaviour of wintering common eiders: implications for habitat use and collision risk. *Journal of Applied Ecology*, **44**, 516-522.
- Leddy, K.L., Higgins, K.F. & Naugle, D.E. (1999) Effects of wind turbines on upland nesting birds in Conservation Research Programme grasslands. *Wilson Bulletin*, **111**, 100-104.
- Moorcroft, D., Whittingham, M.J., Bradbury, R.B. & Wilson, J.D. (2002) The selection of stubble fields by wintering granivorous birds reflects vegetation cover and food abundance. *Journal of Applied Ecology*, **39**, 535-547.
- Osborn, R.G., Higgins, K.F., Usgaard, R.E., Dieter, C.D. & Neiger, R.D. (2000) Bird mortality associated with wind turbines at the Buffalo Ridge Wind Resource Area, Minnesota. *The American Midland Naturalist*, **143**, 41-52.
- Payne, R.J.H. & Wilson, J.D. (1999) Resource limitation in seasonal environments. *Oikos*, **87**, 303-314.
- Percival, S.M. (2005) Birds and windfarms: what are the real issues? *British Birds*, **98**, 194-204.
- Perkins, A.J., Whittingham, M.J., Bradbury, R.B., Wilson, J.D., Morris, A.J. & Barnett, P.R. (2000) Habitat characteristics affecting use of lowland agricultural grasslands by birds in winter. *Biological Conservation*, **95**, 279-294.
- Quinn, J.L., Whittingham, M.J., Butler, S.J. & Cresswell, W. (2006) Noise, predation risk compensation and vigilance in the chaffinch *Fringilla coelebs*. *Journal of Avian Biology*, **37**, 601-608.

Reijnen, R., Foppen, R. & Terbraak, C. & Thissen, J. (1995) The effects of car traffic on breeding bird populations in woodland. 3. Reduction of density in relation to the proximity of main roads. *Journal of Applied Ecology*, **32**, 187-202.

Reijnen, R., Foppen, R. & Meeuwsen, H. (1996) The effects of traffic on the density of breeding birds in Dutch agricultural landscapes. *Biological Conservation*, **75**, 255-260.

Sutherland, W.J. (1996) *From Individual Behaviour to Population Ecology*. Oxford University Press, Oxford, UK.

Vickery, J.A., Evans, A.D., Grice, P., Aebischer, N.A. & Brand-Hardy, R. (2004) The Ecology and Conservation of Lowland Farmland Birds II: The Road to Recovery. *Ibis Supplement*.

Whittingham, M.J. (2007) Will agri-environment schemes deliver substantial biodiversity gain and if not why not? *Journal of Applied Ecology*, **44**, 1-5.

Whittingham, M.J. & Evans, K.L. (2004) A review of the effects of habitat structure on predation risk of birds in agricultural landscapes. In *Ecology and Conservation of Farmland Birds II: the road to recovery*. *Ibis*, 146, 211-222.

Whittingham, M.J., Krebs, J.R., Swetnam, R.D., Thewlis, R. Wilson, J.D. & Freckleton, R.P. (in press) Habitat-associations of British breeding farmland birds. *Bird Study*.

Table 1. Descriptions of predictors included in statistical models. * - Note that as visit was not significant and its inclusion or omission made little difference to the final models, Table 2 is presented without visit (see methods for further details).

Predictor name	Description
Distance from turbine	Variable from 1 to 5. 150 x 150 m blocks of land were surveyed at varying distances from wind turbines in the following categories: (1) <150 m; (2) 150-300 m; (3) 300-450 m; (4) 450-600 m; (5) 600-750 m.
% Boundary	Variable from 0 to 100%. Each block visited at each survey visit was scored as a '1' if it had a boundary feature present (e.g. hedge, ditch), and a '0' otherwise. Total blocks in each distance category were collated, e.g. visit 1 recorded boundary features in 9 out of 16 blocks thus % boundary = 0.56.
% cereals	Variable from 0 to 100%. The large majority of crops were either oil-seed rape or winter wheat and so the proportion of cereals was used as a predictor of differences due to crops. The proportion of cereals was recorded in the same way as % boundary.
Visit	Variable included as Julian date.*
Site	Random effect with two levels. 1 = Glassmoor; 2 = Deeping St.Nicholas

Table 2. Results of binomial error models for four functional groups of birds surveyed at two windfarm sites in East Anglia (each with 8 wind turbines). Results of the deletion of each predictor from a model containing all predictors (i.e. a full model) are presented. Wald statistics are presented (degrees of freedom was 1 in every case), along with significance values (P) (note that P -values are calculated using the χ^2 distribution) and the direction of the relationship is shown in brackets (+ or -) if the relationship was significant. * indicates P -values <0.05 , ** $P<0.01$.

Species or functional group	Distance from turbine	% Boundary	% cereals
Granivores	0.26, $p = 0.61$	4.19, $p = 0.04 (+)^*$	0.45, $p=0.50$
Eurasian skylark	0.53, $p = 0.47$	3.82, $p = 0.05 (-)^*$	0.14, $p = 0.71$
Corvids	2.64, $p = 0.10$	1.39, $p = 0.24$	5.46, $p = 0.02 (+)^*$
Gamebirds	1.56, $p = 0.21$	0.05, $p = 0.82$	0.19, $p = 0.66$
Common pheasant only	8.04, $p = 0.005 (+)^**$	0.23, $p = 0.63$	1.05, $p = 0.31$

Figure 1. Proportion of visits in which one or more of each functional group was recorded in different distance bands from wind turbines. For example, 21% of visits to patches within 150 m of wind turbines contained one or more Eurasian skylarks, but only 18% of visits to patches 600-750 m from wind turbines recorded one or more Eurasian skylarks. Raw data are presented here but the inclusion of crop type and boundaries did not alter the conclusions drawn (Table 2).




Figure 2. The probability of finding common pheasants (% common pheasant) in a block increased with increasing distance from wind turbines, but there was no difference for red-legged partridges (% partridge).

