Brandt K, Srednicka-Tober D, Baranski M, Sanderson R, Leifert C, Seal CJ.
Methods for Comparing Data across Differently Designed Agronomic Studies:
Examples of Different Meta-analysis Methods Used to Compare Relative
Composition of Plant Foods Grown Using Organic or Conventional Production
Methods and a Protocol for a Systematic Review.
Journal of Agricultural and Food Chemistry 2013, 61(30), 7173-7180

Copyright:

This document is the Accepted Manuscript version of a Published Work that appeared in final form in *Journal of Agricultural and Food Chemistry*, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <http://dx.doi.org/10.1021/jf4008967>

Date deposited:

03/09/2013

This work is licensed under a [Creative Commons Attribution-NonCommercial 3.0 Unported License](http://creativecommons.org/licenses/by-nc/3.0/)

Methods for Comparing Data Across Differently Designed Agronomic Studies: Examples of Different Meta-Analysis Methods Used to Compare Relative Composition of Plant Foods Grown Using Organic or Conventional Production Methods, and a Protocol for a Systematic Review.

Kirsten Brandt,^{#,*} Dominika Średnicka-Tober,[†] Marcin Barański,[†] Roy Sanderson,[‡] Carlo Leifert,[†] and Chris Seal[#]

[#]Human Nutrition Research Centre, School of Agriculture, Food and Rural Development, Newcastle University, Newcastle upon Tyne, NE1 7RU, United Kingdom

[†]Nafferton Ecological Farming Group, School of Agriculture, Food and Rural Development, Newcastle University, Stocksfield, Northumberland, NE43 7XD, United Kingdom

[‡]School of Biology, Newcastle University, Newcastle upon Tyne, NE1 7RU, United Kingdom.

*Phone: +44 0191 222 5852. Fax: +44 0191 222 6720. E-mail: kirsten.brandt@ncl.ac.uk.

ABSTRACT

Meta-analyses are methods to combine outcomes from different studies in order to investigate consistent effects of relatively small magnitude, which are difficult to distinguish from random variation within a single study. Several published meta-analyses addressed whether organic and conventional production methods affect the composition of plant foods differently. The meta-analyses were carried out using different options for the methodology, and resulted in different conclusions. The types of designs of field trials and farm comparisons widely used in horticultural and agronomic research differ substantially from the clinical trials and epidemiological studies that most meta-analysis methodologies were developed for. Due to this it is proposed to carry out a systematic review and meta-analysis aiming to develop a consolidated methodology. If successful, this methodology can then be used to determine effects of different production systems on plant food composition as well as other comparable factors with small but systematic effects across studies.

KEYWORDS

Vitamins, Minerals, Horticulture, Food Quality

17 INTRODUCTION

18 Background and Justification

19 Meta-analyses of data from multiple primary studies are used widely to improve the efficiency of the
20 scientific process. Most often these methods are used to assess general applicability of the findings of
21 randomized placebo controlled clinical trials in medicinal research, but they are also increasingly used
22 for other types of studies¹, which include investigations of climate change² and trends in ecological
23 research.³ Meta-analyses are particularly useful to study consistent effects of relatively small magnitude,
24 which are difficult to distinguish from random variation within a single study. One such topic in
25 agricultural chemistry is whether production systems influence food composition, and if so by how
26 much. Numerous single-factor studies show that the composition of plant foods can be affected if the
27 plants are exposed to substantial differences in growing conditions.⁴ It is much less certain, however,
28 whether organic and conventional production methods result in significant differences in composition of
29 the plant products produced in these two systems, in particular regarding compounds that are
30 synthesized by the plant or naturally accumulate within it (in contrast to e.g. pesticide residues). Until
31 now seven published studies⁵⁻¹¹ have included meta-analyses of the relative contents of all or some of
32 the following groups of compounds: vitamins, minerals, macronutrients and secondary metabolites in
33 plant foods grown using organic or conventional production methods. All of these studies compared data
34 from primary publications reporting composition of comparable products produced in the two systems,
35 and in all of these studies the aim was to determine whether the production methods result in significant
36 differences in composition of the plant foods, primarily relating to nutritional value. The seven reviews
37 came to very different conclusions (see Table 1), even in cases where comparable compounds were
38 included in the analyses. Whilst to some extent differences in conclusions may reflect differences in
39 interpretation of the outcomes ('glass half full' versus 'glass half empty'), there are also definitive
40 numeric differences between studies in the calculated significance levels and magnitudes of the effects

of production system, which can only have been caused by the use of different methods for the meta-analysis.

The 7 studies were all based on the principle of meta-analysis, where information from several independent datasets from primary studies are converted into a common metric, which allows the assessment of a treatment effect independently of the specific characteristics of each dataset. Then the metric is used to assess significance of the treatment effect across all datasets. So if for example a treatment is applied to plants from several different species, and most of them respond with a reduction in the content of the nutrient, then the metric will allow the analysis of this effect using the data from all the species, even if the range of nutrient content values within one species is different from that of another species. If relevant for the study designs, the analysis may include a weighting procedure to compensate for objective differences in data reliability, for example giving more weight to studies with several independent replications than to non-replicated ones. The primary advantage of the meta-analysis principle is that it greatly increases the statistical power, and therefore allows statistical evaluation of effects of much smaller magnitude than what can be definitively assessed within a single experimental study. The key steps of a meta-analysis are: 1. choosing the primary studies to include; 2. choosing the type of metric and calculating it; and 3. analyzing the metric and reporting the outcomes. However, a range of different options is available at each step of the procedure, and the optimal choice depends on the types of data analyzed (for example if they are from controlled experiments or from surveys) as well as the aims of the investigation (to estimate the magnitude of an effect or only to assess its significance). A key issue is that most of the development of best practice has focused on research on humans, either in medical or social sciences,^{12, 13, 1} making it relatively more difficult for researchers in the food and agriculture areas to select the most appropriate methodology for meta-analyses in their area. This is reflected in the observation that each of the 7 studies carried out their meta-analysis using different specific methods (see Table 2). The differences in methodology between the reviews are so substantial that this can fully explain why the outcomes were so different, despite similar aims and

substantial overlap in the primary studies included. In fact any pair of two studies differs on more than one major element of the methodology, so it is not possible to assign the differences in outcomes to any specific elements of the methodologies. The choice of method options may substantially affect the outcome, potentially introducing a bias to the conclusions. The robustness of conclusions can however be objectively assessed by sensitivity analyses, comparing the outcomes resulting from different key options in the methodology. Consensus statements about best practice for meta-analyses^{12, 1} strongly emphasize the importance of clearly defined and transparently described methodology, and consistent use of sensitivity analyses wherever the appropriate procedures are not self-evident. In particular this is important for meta-analyses of observational data,¹ which often comprise individual studies with different designs and/or where design and reporting has been done without consideration of the suitability of the data for subsequent meta-analyses. A particular element of best practice¹² is that the protocol for a planned meta-analysis should be published before the analysis is carried out, to provide a definitive documentation of the starting point for the work. This provides a means to monitor that all subsequent adaptations to the protocol are logged and explained. Publication of the protocol also serves to allow constructive criticism from the scientific community, such as suggestions for improvements to the planned procedures.

However, in each of the seven above-mentioned reviews⁵⁻¹¹ only one method was used to select the data to include in the meta-analysis, only one of the studies published a protocol in advance⁷ and only one or two alternative methods were used to analyze the data. In every study the criteria for selection of data to include in analyses were defined solely based on the authors' preferences (not sensitivity analyses), in most cases without any specific explanation for the choices of methods. Only two studies reported that sensitivity analyses were performed. In Brandt et al.,⁶ the outcomes were calculated with or without dry matter adjustment. In Smith-Spangler et al.,¹⁰ different estimated values for sample size were explored for studies where this value was not available, as well as sub-group analyses of differences between plant species or dry matter v. fresh weigh measurements for a few compounds. In all these cases it was

shown that the tested adjustments did not change the significance of the conclusions, but the precise effects on the outcomes were not reported. This clearly indicates how a consolidated methodology for meta-analyses of agronomic/horticultural data would be beneficial.

We propose to use the same approach as used in environmental research² when a similar situation occurred regarding the effect of elevated carbon dioxide on soil carbon (4 independent meta-analyses of the same type of data reported substantially different outcomes). By analyzing similarities and differences among the previous studies, a more robust and authoritative methodology was developed, which succeeded to resolve the question and also explain the reasons for the discrepancies.²

Objectives of the Review

- To systematically collect all primary data on paired comparisons of contents of vitamins, minerals, macronutrients and secondary metabolites in plant foods grown using organic or conventional production methods.
- To carry out relevant sensitivity analyses of the methods used for selection of studies of appropriate quality, to determine the advantages and disadvantages of each of the methods and assess the effect of the outcomes of using more or less optimal methods for this purpose.
- To carry out relevant sensitivity analyses of the methods used for selection and adjustment of data from those studies to include in meta-analyses, to determine the advantages and disadvantages of each of the methods and assess the effect of the outcomes of using more or less optimal methods for this purpose.
- To carry out relevant sensitivity analyses of the methods used for comparison of these data, to determine the advantages and disadvantages of each of the methods and assess the effect of the outcomes of using more or less optimal methods for this purpose.
- To determine the significances and magnitudes of differences found using selected combinations of methods including those considered optimal based on the sensitivity analysis.

116

117 **MATERIALS AND METHODS**118 **1. Inclusion Criteria**

119 The review will include all original studies that compare primary data on the content of vitamins,
120 minerals, macronutrients and secondary metabolites in plant foods grown using organic or conventional
121 production methods. Studies published on paper or electronically (including e.g. PhD thesis repositories
122 and web-only conference proceedings) after January 1958, in all languages, both peer-reviewed and non-
123 peer reviewed, will be included initially.

124 **2. Exclusion Criteria**

125 The review will not include publications containing data from only one production system. In case of
126 duplicates of primary data the least detailed version will be excluded. Studies will be excluded if the
127 essential information (see Table 3) is missing from the published description and cannot be obtained by
128 contact with the author. Specific languages may be excluded if it is not possible to locate sufficiently
129 qualified assistance within the project group and its affiliates to allow extraction of the essential
130 information.

131 **3. Literature Search**

132 Papers will be identified through an initial search of the literature with Web of Knowledge and Scopus,
133 followed by further hand search of reference lists of identified reviews and original publications.
134 Corresponding authors of the identified publications will also be contacted by email and asked to
135 provide further relevant publications.

136 **4. Search Terms for Electronic Databases**

137 Titles of publications will be identified through the search of the literature using the search term
138 structure: (organic* or ecologic* or biodynamic*) and (conventional* or integrated) and (names of 72
139 relevant crops and foods) and (names of 51 relevant compounds and constituents) for the period 1958 –
140 present.

5. Data Extraction

Information from the identified publications will be extracted to an MS Access database as detailed in Table 3. A set of paired values is defined as the values provided for the concentration of one 'compound' in the same product from the same variety (or cultivar if variety name is unavailable) and treatment. A 'compound' can be a chemically defined entity (e.g. iodine) or several chemically related compounds measured together (e.g. 'sugars' or 'hydroxycinnamic acids'). For initial data collection, each 'compound' will be recorded as defined by the author, although where more than one synonym is used for the same 'compound' in different articles, the most frequently used name will subsequently be applied for the data from all articles. Some of the subsequent analyses will involve the grouping of 'compounds' according to the different principles used in each of the published meta-analyses (see 'Handling of multiple forms of compounds' in Table 2). Data presented in graphical form on arithmetic scales will be recorded manually, while graphical data presented on logarithmic scales will be excluded.

6. Quality Control

For most publications the data will be extracted by one person. To reduce the risk of spelling errors (including errors in the publications), for data recorded as a limited number of repeated words or phrases (e.g. author names, research units, names of compounds and analysis methods etc.), each new occurrence of such a word or phrase will be added to a list, while repeated occurrences will be copied from the list. After completion of the data extraction, an initial test for outliers will be used to find and correct obvious errors. Those authors whose work has been included in the review may then be contacted and asked to check whether the extracted information is accurate and also invited to provide additional information if relevant. In particular this will be done in cases where information is missing or unclear in the publication, including data presented only as graphics or only as averages of factors such as years. For such publications, if the response is insufficient or not received, the data will be checked by repeated extraction by a different person.

165 Once the meta-analysis data have been published, the full extracted datasets will be made available
166 online, allowing any interested party to check and reproduce all the published calculations. The website
167 will include a facility for reporting of errors or additional data. If errors in data or calculations are
168 reported, which may affect the conclusions of the meta-analyses, the affected analyses will be repeated
169 and the corrected results presented on the website.

170 **7. Comparison of Analysis Methods**

171 The methods used by each of the previous authors (see Table 2) will be compared using iterative
172 sensitivity analysis, changing one element of a method at a time where feasible, in those cases where the
173 description is sufficiently detailed to allow it to be replicated. Methods from other studies on related
174 topics, such as the paper by Seufert et al.¹⁴ on comparison of crop yields in organic and conventional
175 agriculture, will also be considered and included where relevant. Specifically, in this paper the variable
176 used was the ln-transformed response ratio weighted by the inverse of the mixed-model variance,
177 excluding studies with only one data pair. In contrast, another recent review by Hoefkens et al.¹⁵ with a
178 meta-analysis of composition of organic and conventional vegetables, used a fundamentally different set
179 of methods, which we probably will not be able to compare with the other meta-analyses. This study was
180 to a large extent based on primary data from studies on composition of vegetables from only one
181 production system ('unpaired' data), and rather than assessing the physiological responses of the plants
182 to the production methods, the aim was described as 'to investigate the scientific validity of nutrition
183 claims such as "no vegetable/potato has higher amounts of nutrient X than organic vegetables/potatoes".
184 This study's methods are so different from the other reviews that it was not possible to include them in
185 Table 2.

186 Prior to this step, the authors of all the meta-analysis studies will be contacted to ask if they are
187 interested to provide additional information about the methodology they used, as well as if they have
188 additional suggestions for which methods we should include in the comparison. The methods used as the
189 reference in these comparisons will be the ones used in the study by Brandt et al.⁶ Comparisons will be

done in groups of elements, first comparing the methods for extraction and analysis of the data, then the methods for selection of publications to include, and finally combinations of relevant methods in each group.

8. Examples of Sensitivity Analyses:

- Quality criteria for the publications included: e.g. peer-reviewed papers only vs. both peer-reviewed and non-peer reviewed; large sample size versus low or unreported sample size; etc.
- Handling of data from studies with different designs: e.g. data for separate years included as separate data points vs. data for separate years averaged; with or without weighting based on standard variation; multiple cultivars treated as separate samples or as single samples; etc.

EXPECTED RESULTS AND DISCUSSION

The study will develop and implement concepts from other research areas^{2, 3, 16} to provide a comprehensive meta-analysis of the effects of production systems on the composition of plant foods regarding contents of vitamins, minerals, macronutrients and secondary metabolites, incorporating the new original data published since the most recent reviews covering each of these types of compounds, as well as the accumulated experience with different approaches for such meta-analyses. In addition, the multiple sensitivity analyses and subgroup comparisons will provide systematic information about the consequences of using different methods for selection of publications, extraction of information from them and analysis of the data obtained.

We expect that the primary reason why different methods have caused different outcomes is that the number of independent primary studies is relatively low, with a variety of different design principles and potentials for within-study correlations. Therefore over time, as more independent original studies are published, the differences between the overall outcomes caused by using the different methods will be reduced, as found in other comparable areas.³ If this is confirmed, it also means that the (combinations of) methods that are best able to estimate the overall outcome from relatively small data volumes (for

example using reduced timespans or limited selections of crop species) are objectively better than other methods, thus providing a concept for unbiased ranking of the quality of the methodologies.

We therefore aim to derive a set of objectively justified general guidelines for future meta-analyses, which will be used for future updates to include new original data in the specific area. They may also turn out to be useful to analyze other types of external effects (e.g. climate) on other qualities of cultivated plants (e.g. resistance to diseases), where experimental designs show the same types of variations as for studies comparing organic and conventional crops.

The outcome of the present study will also provide objectively justified guidelines for the design and reporting of future primary studies. It will quantify and illustrate the importance of multi-year recordings, appropriately replicated experiments, meticulous experimental protocols and reporting of relevant statistical outcomes, providing evidence that can be used by educators and journal editors to encourage best practice whenever possible. However it will also demonstrate appropriate exploitation of the data resource provided by multiple independent non-replicated experiments, each of which are often insufficient to justify publication on its own, and may thus encourage initiatives to make more good quality unpublished datasets available for secondary research.

REFERENCES

- (1) Stroup, D. F.; Berlin, J. A.; Morton, S. C.; Olkin, I.; Williamson, G. D.; Rennie, D.; Moher, D.; Becker, B. J.; Sipe, T. A.; Thacker, S. B., Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *Jama* **2000**, 283, 2008-12.
- (2) Hungate, B. A.; van Groenigen, K. J.; Six, J.; Jastrow, J. D.; Luo, Y. Q.; de Graaff, M. A.; van Kessel, C.; Osenberg, C. W., Assessing the effect of elevated carbon dioxide on soil carbon: a comparison of four meta-analyses. *Global Change Biol* **2009**, 15, 2020-2034.

- (3) Leimu, R.; Koricheva, J., Cumulative meta-analysis: a new tool for detection of temporal trends and publication bias in ecology. *P Roy Soc B-Biol Sci* **2004**, *271*, 1961-1966.
- (4) Poiroux-Gonord, F.; Bidel, L. P.; Fanciullino, A. L.; Gautier, H.; Lauri-Lopez, F.; Urban, L., Health Benefits of Vitamins and Secondary Metabolites of Fruits and Vegetables and Prospects To Increase Their Concentrations by Agronomic Approaches. *J Agric Food Chem* **2010**, *58*, 12065–12082.
- (5) Benbrook, C.; Zhao, X.; Davies, N.; Andrews, P. *New evidence confirms the nutritional superiority of plant-based organic foods*; The Organic Center: 2008. pp 49.
- (6) Brandt, K.; Leifert, C.; Sanderson, R.; Seal, C. J., Agroecosystem management and nutritional quality of plant foods: the case of organic fruits and vegetables. *Critical Reviews in Plant Sciences* **2011**, *30*, 177-197.
- (7) Dangour, A. D.; Dodhia, S. K.; Hayter, A.; Allen, E.; Lock, K.; Uauy, R., Nutritional quality of organic foods: A systematic review. *Am J Clin Nutr* **2009**, *90*, 680-685.
- (8) Heaton, S. *Organic farming, food quality and human health. A review of the evidence.*; Soil Association: 2001, pp 88.
- (9) Hunter, D.; Foster, M.; McArthur, J. O.; Ojha, R.; Petocz, P.; Samman, S., Evaluation of the micronutrient composition of plant foods produced by organic and conventional agricultural methods. *Critical Reviews in Food Science and Nutrition* **2011**, *51*, 571-582.
- (10) Smith-Spangler, C.; Brandeau, L. M.; Hunter, G. E.; Bavinger, J. C.; Pearson, M., Are Organic Foods Safer or Healthier Than Conventional Alternatives? A Systematic Review. *Annals of Internal Medicine* **2012**, *157*, 348-366.
- (11) Worthington, V., Nutritional quality of organic versus conventional fruits, vegetables, and grains. *Journal of Alternative and Complementary Medicine* **2001**, *7*, 161-173.
- (12) Moher, D.; Liberati, A.; Tetzlaff, J.; Altman, D. G., Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* **2009**, *6*, e1000097.

- 264 (13) Petticrew, M.; Roberts, H., *Systematic reviews in the social sciences: a practical guide*. Blackwell
265 Pub.: Malden, MA; Oxford, 2006; p xv, 336 p.
- 266 (14) Seufert, V.; Ramankutty, N.; Foley, J. A., Comparing the yields of organic and conventional
267 agriculture. *Nature* **2012**, *485*, 229-232.
- 268 (15) Hoefkens, C.; Vandekinderen, I.; De Meulenaer, B.; Devlieghere, F.; Baert, K.; Sioen, I.; De
269 Henauw, S.; Verbeke, W.; Van Camp, J., A literature-based comparison of nutrient and contaminant
270 contents between organic and conventional vegetables and potatoes. *British Food Journal* **2009**, *111*,
271 1078-1097.
- 272 (16) Quemada, M.; Baranski, M.; Nobel-de Lange, M. N. J.; Vallejo, A.; Cooper, J. M., Meta-analysis of
273 strategies to control nitrate leaching in irrigated agricultural systems and their effects on crop yield.
274 *Agriculture, Ecosystems and Environment* **2013**, *In press*, accepted 15 April 2013.
- 275 (17) Hedges, L. V.; Gurevitch, J.; Curtis, P. S., The meta-analysis of response ratios in experimental
276 ecology. *Ecology and Society* **1999**, *80*, 1150-1156.
- 277 (18) Brickell, C.; Alexander, C.; David, J. C.; Hetterscheid, W. L. A.; Leslie, A. C.; Malecot, V.; Jin, X.;
278 Cubey, J. J., *International code of nomenclature for cultivated plants : (ICNCP or cultivated plant code)*
279 *incorporating the rules and recommendations for naming plants in cultivation : adopted by the*
280 *International Union of Biological Sciences, International Commission for the Nomenclature of*
281 *Cultivated plants*. 8th ed. / prepared and edited by C.D. Brickell ... [et al.]. ed.; ISHS: Leuven, 2009.

282

283

TABLES

Table 1. Summary of outcomes from previous studies on relative nutrient content in organic and conventional plant foods, resulting from analysis of literature data from multiple comparative studies.							
Lead author	Benbrook	Brandt	Dangour	Heaton	Hunter	Smith-Spangler	Worthington
Year published	2008	2011	2009	2001	2011	2012	2001
Types of compounds included	Vitamins, minerals and secondary metabolites	Secondary metabolites and vitamin C	Vitamins, minerals and secondary metabolites	Vitamins, minerals and secondary metabolites	Vitamins and minerals	Vitamins, minerals and secondary metabolites	Vitamins and minerals
Number of compound groups compared	11	7 groups + 3 combined groups	11	5	24 groups + 4 combined groups	13	5, and 32 without test
Number of compound groups statistically significantly different	No test	5 groups + 3 combined groups	3	No test	1 group + 2 combined groups	2	5
Overall conclusion	'organic plant-based foods are ... more nutritious'	'higher than in corresponding conventional samples'	'no evidence of a difference in nutrient quality'	'organically grown crops are significantly different'	'levels of micronutrients were higher in organic foods'	'lacks strong evidence that organic foods are significantly more nutritious'	'genuine differences in the nutrient content'
Full dataset published	No	Yes	Yes	No	No	No	No

Table 2. Summary of methods used in previous studies							
Lead author	Benbrook	Brandt	Dangour	Heaton	Hunter	Smith-Spangler	Worthington
Year published	2008	2011	2009	2001	2011	2012	2001
Publication year for data	1980-2007	1992-2009	1993 (1958 in protocol) - 2007	1974-2000	1980-2007	1978-2012	1951-1997
Selection criteria for organic treatment	A combination of the following using points: Produced under qualified organic management. Stated appropriate conversion period; stated adherence to appropriate regulation or certification body; input list provided that conforms with what is allowed by appropriate regulation or certification body	At least one of the following: Stated adherence to appropriate regulation or certification body; input list provided that conforms with what is allowed by appropriate regulation or certification body; the word 'organic' used in a 'legally defined context': a time and place where it is illegal to use it for production methods that do not conform with what is allowed by appropriate regulation	Stated adherence to appropriate regulation or certification body. Non-standard organic certified systems such as 'biodynamic' were excluded in principle, but included if organic certification was stated	Produced under qualified organic management including explicitly stated appropriate conversion period	Described as 'organic' or 'biodynamic' or utilized 'organic manure' without the use of synthetic inputs. Plant protection methods not specifically mentioned, but presumably excluded synthetic pesticides	Included biodynamic and defined by investigators' stated adherence, must include 'farming practice' (probably plant protection)	Stated use of organic fertilizer or organic system. Plant protection methods not specifically mentioned
Selection criteria for conventional	As defined by author where more than one	As defined by author where more than one	Only explicitly 'conventional	Required to reflect typical	Described as 'conventional', 'integrated', 'low	Stated use of conventional fertilizer or	Stated use of conventional fertilizer or

treatment	non-organic system is included in a study. Commonly used certified non-organic systems ('Integrated', 'Good Agricultural Practice' etc.) considered conventional, unless explicitly contrasted with other conventional practice	non-organic system is included in a study. Commonly used certified non-organic systems ('Integrated', 'Good Agricultural Practice' etc.) considered conventional, unless explicitly contrasted with other conventional practice	' practice considered conventional. Non-organic certified systems such as 'Integrated' were excluded in principle, but included if certification was not mentioned	conventional practice, but also accepted inclusion of otherwise acceptable studies that only compared fertilization but not crop protection methods	input' or 'mineral fertilizers'. Plant protection methods not specifically mentioned	conventional system. Plant protection methods not specifically mentioned	conventional system. Plant protection methods not specifically mentioned
Selection criteria for general quality of publications	Included if method descriptions sufficiently detailed to allow assessment of the other criteria and the data are not duplicated in another publication	Included if method descriptions sufficiently detailed to allow assessment of the other criteria and the data are not duplicated in another publication	Only peer-reviewed publications included, identification and removal of duplicate data not mentioned	Included if method descriptions sufficiently detailed to allow assessment of the other criteria and the data are not duplicated in another publication	Included if method descriptions sufficiently detailed to allow assessment of the other criteria. Also 'mixed foods' and studies where harvest date was not specified were excluded	Only peer-reviewed English language publications included, identification and removal of duplicate data not mentioned	No specific criteria mentioned
Selection criteria for publications regarding analysis	Studies excluded if using analytical methods considered imprecise or not	No specific criteria, other than exclusion of studies that explicitly stated	Studies excluded if methods or statistics were not mentioned	No specific criteria	No specific criteria other than that analytical methods must be clearly stated	Studies excluded if information about variance or	No specific criteria other than that data must be quantitative

methods	defined	the use of inappropriate methods				results of statistical tests was missing	(numeric)
Selection criteria for trial design	Field trials and farm comparisons only	Field trials, farm comparisons and shopping basket studies all included in principle, but all shopping basket studies excluded due to other criteria	Field trials, farm comparisons and shopping basket studies all included	Field trials, farm comparisons and shopping basket studies all included	Field trials, farm comparisons and shopping basket studies all included in principle, unless multiple levels and types of fertilizer were included, but all shopping basket studies excluded due to other criteria	Field trials, farm comparisons and shopping basket studies all included	Field trials, farm comparisons and shopping basket studies all included
Selection criteria for plant genotype comparisons	Same cultivar ^a in organic and conventional treatment	Same cultivar ^a in organic and conventional treatment	Same Group ^a in organic and conventional treatment	Not clearly specified	Same cultivar ^a in organic and conventional treatment.	Not a selection criterion, but ‘evaluated’ the extent to which pairs were of the same Group ^a	Not clearly specified
Percentage of publications included after application of quality criteria	40%	65%	33%	30%	50%	>85% (not clearly specified)	>90% (not clearly specified)
Handling of	Where data on	Where data on	Where data	Not clearly	Not clearly	Not clearly	Not clearly

multiple forms of compounds	multiple forms of a compound were reported, if a 'total' or 'combined' value was present, only this was included in the meta-analysis, if not, only the most abundant form was used, the rest were ignored	multiple forms of a compound were reported in the same study, the contents of the most similar forms were added to form a maximum of 6 aggregate groups of compounds per sample. No data were excluded unless they were clearly duplicated or included in other data. Compounds were grouped into 7 categories (6 types of secondary metabolites + vitamin C) and each group analyzed separately and in combinations	on multiple forms of a compound were reported, every one of them was included separately. Compounds were grouped into 98 categories, and categories represented by at least 10 studies were analyzed separately	specified	specified, however presumably all forms of the same vitamin, mineral or trace elements were considered equivalent, since only these types of compounds were included	specified, however 14 'vitamins and nutrients' were included, some of which presumably include multiple forms	specified
Handling of multiple treatments within a study (e.g. storage duration, harvest times,	Where multiple treatments were reported, only one was chosen for inclusion, e.g. shortest duration of postharvest, or methods representing 'the	Where multiple treatments were reported, the average of all treatments was calculated and used, except for multiple durations of post-	Where multiple treatments were reported, either every one was included as separate data	Not clearly specified	Where multiple treatments were reported, the data from the least processed treatment and/or the maturity stage most representative of	Data for separate treatments were presumably combined using random-effect	Where multiple treatments were reported, as far as possible every one was included as separate data

storage temperature etc.)	most common organic and conventional treatments’ (not clear if they had to be identical).	harvest treatments, where only the shortest duration was used	points, or the average was calculated. Criteria for choosing one or the other option were not specified		commercial maturity were used	models – this is not clearly specified	points
Handling of data from studies containing more than one year/growing season	Where data were reported for multiple years combined, as well as for each year separately, the combined value was used, although separate values for each year were used if no combined value was reported	Where reported in the publication or available from the author, as far as possible data for separate years were included as separate data points	Not clearly specified	Not clearly specified	Not clearly specified, other than that ‘compared foods were grown in the same season’	Data for separate years were combined using random-effect models	As far as possible data for separate years were included as separate data points
Handling of data from studies containing more than one location	Where data were provided for separate locations, each location was included separately	Where data were provided for separate locations, the average for all locations was calculated and used	Where data were provided for separate locations, each location was included separately	Not clearly specified	Not clearly specified, other than that ‘Data reported as an average of multiple farms were considered as a single comparison’	Not clearly specified, but presumably treated as ‘samples’ and therefore combined using random-effect models	Not clearly specified
Handling of	Data from each	Within a Group ^a	Data from	Not clearly	Individual	Not clearly	Not clearly

data from studies containing more than one genotype	cultivar ^a were used separately. Studies where cultivar ^a names were not provided were excluded	the average of data from all cultivars ^a was used. Studies were excluded if cultivar ^a names were not provided	each cultivar ^a were used separately, except in studies (mainly shopping surveys) where only data on the average for a Group ^a was provided	specified	cultivars ^a of the same food were used as separate comparisons	specified, but presumably treated as 'samples'	specified
Handling of data reported on fresh weight or dry matter basis.	Not clearly specified	Data reported on dry matter basis was adjusted to correspond to values on fresh weight basis by multiplying with the average ratio of dry matter content in organic and conventional produce, calculated from those studies where the dry matter content itself was reported	No distinction made	Only data reported on fresh weight basis included	No distinction made	Not clearly specified, but subgroup analyses were done for 3 compounds (calcium, potassium and phosphorus) showing no change in outcome	Not clearly specified
Handling of data reported only in graphical form	Not clearly specified	Approximate values measured manually. Graphical data on a logarithmic	Excluded	Not clearly specified	Not clearly specified	Not clearly specified	Not clearly specified

		scale were excluded					
Variable used for statistical analysis	Response ratio of concentrations in organic and conventional material, also used ‘vote-counting’	Ln-transformed response ratio of concentrations in organic and conventional material (also used normalized difference)	Normalized difference between concentrations in organic and conventional material	‘Vote-counting’	Normalized difference between concentrations in organic and conventional material, also used ‘vote-counting’	Standardized mean difference	Normalized difference between concentrations in organic and conventional material, also used ‘vote-counting’
Statistical test	No test	Re-sampling test ¹⁷ (also t-test)	T-test with ‘robust standard deviation’ (no reference provided)	No test	One-sample t-test or Wilcoxon signed rank tests for differences depending on normality of distributions, ‘sign test’ for ‘vote-counting’	Random-effect models and funnel plots	Wilcoxon signed-rank

^a The terms ‘cultivar’ and ‘Group’ are used here according to the ‘International code of nomenclature for cultivated plants’¹⁸. The cited reviews used either ‘cultivar’ or ‘variety’ when describing the selection process. However none of the reviews (nor individual studies) provided a reference or correct definition of the terms they used, reflecting widespread inconsistency and lack of awareness of this issue. Due to this, the classification here is based on the actual selection outcome, specifically on whether or not a review included studies defining the genotype only at ‘Group’ level as assessed by this criterion.

Table 3. Information to be extracted from the identified publications		
Type of information	Pre-defined categories	Essential information
General information about the publication		
Source of publication	Electronic database searches, hand searches of reference lists of reviews and original publications, direct contact with the authors	No
Author names (at least one)		Yes
Title		No
Year of publication		No
Type of publication	Journal article, conference paper, conference proceedings, book chapter, report, thesis, other	Yes
Name of Journal or Book Series where relevant		No
Country of the first author		No
Research unit of the first author		No
E-mail address of corresponding author		No
Language of publication		No
Availability of English abstract	Yes, no (assumed no if not specified)	N/A
Information if the data presented in the publication were duplicated in other study	Yes, no (assumed no if not specified)	N/A
Potential study quality indicators (for sensitivity analysis)		
Peer-reviewed	Yes, no (assumed no if not specified in publication, publisher's website or other available information)	N/A
Funding source	Public, organic industry, conventional industry, other	No
Type of study	Shopping basket study, farm comparison, controlled experiment	Yes
Number of experimental years		Yes
Location as country, if available also province/state/county and town		Yes
Plant species and, if relevant and available, Group(s) ^a		Yes

Number of plant cultivars ^a included		No
Production system (as defined by author)	Organic, biodynamic, conventional, integrated	Yes
If products are certified (both for organic and conventional treatments)	Yes, no (assumed no if not specified)	N/A
Name of certifying body		No
If inputs are specified	Yes, no	N/A
If specified inputs are compatible with organic standards	Yes ^b , no ^b	N/A
Data recorded for each paired value		
Plant species and, if relevant and available, Group ^a		Yes
Product analyzed		Yes
Plant cultivar ^a name		No
Compound name		Yes
Unit of measurement		No
Name of analytical method or ‘no method provided’		N/A
Value for concentration for each data point		Yes
Value reported on fresh weight basis	Yes, no (assumed yes if not specified)	N/A
Standard error of the mean (SEM) or standard deviation (SD) of the value		No
Number of full replications of each system in terms of growing sites or field plots		No
Number of technical repetitions within each replication		No
If difference between systems was reported by author as significant	Yes, no	N/A
Dry matter %		No
Data source	Numeric, graphical	N/A
Limit of detection (LOD) and/or limit of quantification (LOQ)		No
All ‘treatments’ distinguishing separate sets of paired data found in the publication, which are replicated for at least one organic and one conventional sample, to ensure that all relevant data points from the publication are included in as much detail as available.	E.g. preservation method, sampling/harvesting time/method, year, season, irrigation method, maturity stage at harvest, pre-crop, seed rate, storage conditions etc.	N/A
Data control status	Confirmed or corrected by author, updated or extended by author, confirmed or corrected without input from author	N/A
^a ‘Group’ and ‘cultivar’ defined according to ¹⁸ (as in Table 2); ^b Assessed by study staff based on available input lists, if not explicitly stated in publication. N/A means ‘not applicable’.		

TABLE OF CONTENTS GRAPHICS