Newcastle University ePrints

Kuzmenko AV, Levitskii SA, Vinogradova EN, Atkinson GC, Hauryliuk V, Zenkin N, Kamenski PA. (2013) [Protein biosynthesis in mitochondria](#).
Biochemistry (Moscow), 78(8), 855-866.

Copyright:

This is a preprint of the materials accepted for publication in *Biochemistry (Moscow)*, © 2013, Pleiades Publishing, Ltd.

DOI link for article:

<http://dx.doi.org/10.1134/S0006297913080014>

Always use the definitive version when citing.

Further information on publisher website: www.maik.ru

Date deposited: 10th September 2013

Version of article: Authors' accepted manuscript

This work is licensed under a [Creative Commons Attribution-NonCommercial 3.0 Unported License](http://creativecommons.org/licenses/by-nc/3.0/)

ePrints – Newcastle University ePrints

<http://eprint.ncl.ac.uk>

Protein biosynthesis in mitochondria

Kuzmenko A.^{1,2}, Levitskii S.A.¹, Vinogradova E.¹, Atkinson G.C.², Hauryliuk V.², Zenkin N.³, Kamenski P.^{1*}

¹Molecular Biology Department, Faculty of Biology, M.V. Lomonosov Moscow State University, 1/12 Leninskie Gory, 119991 Moscow, Russia, fax +7 495 9394309, e-mail mail@mail.bio.msu.ru

²Institute of Technology, University of Tartu, Nooruse St 1, 50411 Tartu, Estonia, fax + 372 737 4900, e-mail info@tuit.ut.ee

³Centre for Bacterial Cell Biology, Institute for Cell and Molecular Biosciences, Newcastle University, Newcastle upon Tyne, NE2 4AX, United Kingdom, fax +44 1912083205, e-mail n.zenkin@ncl.ac.uk

* E-mail for correspondence: peter@protein.bio.msu.ru

Abstract

Translation, that is biosynthesis of polypeptides in accordance with information encoded in the genome, is one of the most important processes in the living cell and has been in the spotlight of international research for many years. The mechanisms of protein biosynthesis in bacteria and in the eukaryotic cytoplasm are now understood in great detail. However, much less is known about translation in eukaryotic mitochondria, which is characterized by a number of unusual features. In this review we summarize current knowledge about mitochondrial translation in different organisms while paying special attention to the aspects of this process that differ from cytoplasmic protein biosynthesis.

Keywords

Mitochondria, translation, ribosomes, translation factors.

Introduction

Almost all eukaryotic cells contain mitochondria. These organelles provide energy via the process of oxidative phosphorylation, and are involved in fatty acid, heme and iron-sulfur cluster biosynthesis, as well as programmed cell death. According to the generally accepted concept of endosymbiosis, mitochondria originated in the early stages of eukaryotic evolution from an endosymbiotic α -proteobacterium closely related to the modern *Rickettsia prowazekii* [1]. Over time, as a result of the tight association with the nuclear genome, many predecessors of mitochondrial genes have been transferred to the nucleus, but a small part of them are retained in the organelle DNA. Translation of mitochondrial messenger RNAs (mRNAs) in the mitochondrion is carried out by its own apparatus of protein biosynthesis, and requires fine coordination between the two genomes.

Many aspects of mitochondrial translation remain poorly studied. This is largely due to the fact that mitochondrial translation systems capable of synthesis on natural matrices are still not reconstructed *in vitro*. On the basis of early studies of sedimentation characteristics and RNA composition of mitochondrial ribosomes, as well as their inhibition by antibiotics, it became apparent that mitochondrial translation is similar to that of bacteria [2 - 4]. Recently, it has become clear that the translation process in the mitochondria has a number of distinctive features [5, 6].

A number of high resolution structural studies have aided our understanding of translation at the molecular level. The majority of these were performed on bacterial and eukaryotic cytosolic ribosomes [7]. However, considerable success has also been achieved in the study of structural and functional properties of the mitochondrial translational system. In this review we summarize the current understanding of protein synthesis in mitochondria.

GENERAL FEATURES OF PROTEIN SYNTHESIS IN MITOCHONDRIA

The mitochondrial genome, despite the universality of its function in energy generation shows a high degree of variation in composition, size and gene content across eukaryotes [8]. For example, the smallest mitochondrial genome described to date is the linear mitochondrial DNA (mtDNA) of *Plasmodium falciparum* (5,966 bp). This encodes, besides the large and small ribosomal RNA (rRNA), only three proteins [9]. The human circular mtDNA (16,569 bp) contains 22 genes of transfer RNA (tRNA), 13 protein coding genes and 12S and 16S rRNA genes. Exceeding this by 22 times in size, the *Arabidopsis thaliana* mitochondrial genome (366,924 bp) contains information for only 32 proteins and an incomplete set of tRNAs [10, 11]. Such disparity is explained by an unusually large number (> 80%) of non-coding sequences in the mtDNA of *A. thaliana* compared with the *Leishmania* (<10%) and mammals, whose genes directly follow each

other or are separated by only a few nucleotides [12]. The rate of evolution of the mitochondrial genome is extremely high, partly due to an increased frequency of mutations due to the lack of an effective repair system and its proximity to reactive forms of oxygen produced near the inner mitochondrial membrane [13].

An important feature of mitochondrial translation is a deviation from the standard genetic code. For example, in vertebrate mitochondria one of the three stop codons - UGA - corresponds to tryptophan, and AUA encodes methionine instead of isoleucine. Mitochondria often use alternative triplets as start codons: AUA (*Homo*, *Bos*), AUU (*Homo*, *Mus*), AUC (*Mus*) and GUG (*Coturnix*, *Gallus*). The amount of stop codons is reduced in mitochondrial systems: usually up to two, however 7 of 12 open reading frames of parasitic nematode *Radopholis similis* mitochondria are deprived of canonical stop codons [14]. Additionally, in yeasts *Saccharomyces cerevisiae* and *Torulopsis glabrata* codons starting with CU have been reassigned from leucine to threonine [15].

Transcription in almost all eukaryotic mitochondria is carried out by RNA polymerase homologous to that T3 and T7 phages [16]. The product of this synthesis is a polycistronic primary transcript [16 – 18]. According to the "tRNA punctuation" model, it is further processed by an endonucleolytic excision of tRNA sequences generating the individual protein-encoding mRNAs [19]. There are 11 human mitochondrial mRNAs: 9 monocistronic and two bicistronic with overlapping reading frames. In *S. cerevisiae* mitochondria there are 6 monocistronic mRNAs and one bicistronic mRNA coding for subunits 6 and 8 of F₀F₁-type ATP synthase [20]. Mitochondrial mRNAs lack the cap structure characteristic for the cytoplasmic translation, and are characterized by dramatic lineage-specific differences. Human mitochondrial mRNAs have very short 5'-and 3'-untranslated regions (UTR), and are polyadenylated [20]. On the other hand, in *Saccharomyces cerevisiae* mature mitochondrial transcripts have long UTRs that are important for translation regulation, and are not polyadenylated [21 – 23]. Apparently, this reflects a fundamental difference in the mechanisms of recognition of the start codon and the initiator complex assembly in the mitochondria of yeast and mammals. Mitochondrial mRNAs, unlike those of bacteria, lack Shine-Dalgarno sequences and, accordingly, the complementary segment of the small subunit rRNA. An octanucleotide sequence UAUAAUA was suggested to play the role of a common ribosome recognition site in yeast but this hypothesis has not yet been experimentally confirmed [24]. The degradation of mitochondrial mRNA is controlled by a poly-subunit protein complex – degradosome possessing an RNA helicase, as well as hydrolytic or phospholytic exonuclease activities [25]. In yeast the mitochondrial polynucleotide phosphorylase was not detected, while in mammals, this enzyme was found in the intermembrane space, where, in addition to its main function, it participates in the regulation of RNA import into mitochondria [26].

The number of different tRNA species in mitochondria is usually significantly reduced in

comparison to the cytoplasmic systems (22 and 24 in yeast and human, respectively), which is compensated by each of the tRNAs recognizing several codons. Mitochondrial tRNAs are usually shorter (59-75 nucleotides length in animals) than those of bacteria and the eukaryotic cytosol [27]. They often lack the conservative elements of the tertiary structure such as long-range interactions between D and T arms, and binding sites of magnesium cations, with the latter feature leading to their low thermal stability [28, 29]. Despite the absence of high resolution structural data, the results of chemical modification, nuclease protection and NMR spectroscopy of T7 transcripts of bovine and nematode mitochondrial tRNA suggest that they assume a structure similar to the canonical L-form of the molecule [5]. This is confirmed by the results of cryoelectron microscopy of *Bos taurus* mitochondrial ribosomes: tRNA firmly bound in the peptidyl transferase center has a typical L-conformation with a cavity in the "elbow" [30]. Despite the presence of a complete set of tRNAs in the mitochondrial genome of *S. cerevisiae*, one of the two cytoplasmic isoacceptor lysine tRNAs with CUU anticodon is also imported into the organelle and is involved in translation [31]. In contrast to animals and yeasts, the majority or even all types of mitochondrial tRNAs in plants and trypanosomatids are imported from the cytoplasm [32, 33].

Mitochondrial ribosomes (mitoribosomes) differ from 70S bacterial and 80S eukaryotic particles in several ways. Budding yeast mitoribosomes have a sedimentation coefficient of 74S, mammals - 55S, and *Leishmania tarentolae* - just 50S [34 – 36]. *B. taurus* and *L. tarentolae* mitoribosome structures resolved by cryoelectron microscopy to 9-14 Å are currently available [30, 35]. As in all ribosomes, mitoribosomes have large and small subunits consisting of ribosomal RNA (rRNA) and proteins (see Table 1). The most significant difference between mitoribosomes and cytoplasmic analogues is in the RNA:protein ratio which is 2:1 in bacteria, while in *B. taurus* and *L. tarentolae* mitochondria - 1:2 and 1:3, respectively. At the same time mammalian mitoribosomes (2.71 MDa in mass and ~ 320 Å in diameter) are larger, but *Leishmania* mitoribosomes (2.2 MDa and ~ 245 Å) are smaller than the *E. coli* ribosome [30, 35]. Low values of sedimentation coefficients of mitoribosomes are due to their porous structure, which is a consequence of the absence or severe reduction of some segments of the rRNA [37]. For example, the small subunit 12S rRNA is completely missing the anti-Shine-Dalgarno sequence as well as helices 6, 8-10, 12, 13, 16, 17, 21, 33, 37 and 39-41 in the mammalian mitoribosome. Some of the lost elements are compensated for by new mitoribosome-specific proteins or by increased protein length relative to their bacterial counterparts. However, in mammals, only 19% of missing sequences in small subunit rRNA and 28% in large subunit rRNA are replaced with proteins, while a similar compensation in *L. tarentolae* mitoribosomes reaches 50%. This has led to a number of distinctive morphological features, while maintaining the general organization of the subunits. Such features include, for example, tunnel-like structures that link the intersubunit space with the external environment.

Shielding of rRNA by proteins and the prevalence of protein-protein intersubunit bridges (9 of 15 in animals) are especially the case in mitoribosomes [30]. A unique feature of the small subunit of the mammalian mitoribosome is an extremely dynamic trihedral gate-shaped structure surrounding the mRNA entry site [30]. This is thought to help in attracting mRNAs lacking leader sequences to the small subunit. The large subunit is characterized by a massive central protuberance and a lack of electron density corresponding to 5S rRNA. However, biochemical studies have shown that 5S rRNA is imported into human mitochondria from the cytosol, and is associated with mitoribosomes [38, 39]. The absence of the core 16S rRNA elements constituting the classical E-site – helices 11 and 68, and the loop portion lying between the helices 76 and 77, leads to very weak binding of tRNA to this region [40]. Another distinctive feature of the mitoribosomal large subunit is the presence of finger-like structures in the P-site, formed by the extension of a portion of the mitochondria-specific proteins of the central protuberance. This finger interacts with the T-loop of the tRNA molecule located in the P-site, and presumably exerts a stabilizing effect on the binding. Apparently, this is the reason that mitoribosomes, unlike bacterial ribosomes, are purified with tRNA firmly bound to the P-site. The organization of the tunnel through which the nascent peptide leaves the ribosome deserves special attention [41]. In the case of ribosomes of bacteria and eukaryotic cytoplasm, it leads to an opening referred to as Polypeptide Exit Site (PES). This is a flat structure, located in the lower part of the large subunit formed by elements of domains I and III of 23S rRNA and ribosomal proteins L22, L23, L24 and L29. The situation is different in the case of *L. tarentolae* and *B. taurus* mitoribosomes, where one more hole was discovered at some distance from PES (~ 25 Å in the case of *L. tarentolae*), called PAS (Polypeptide-Accessible Site) [37, 42]. The formation of this structure results from the absence of most of domains I and III in the large subunit rRNA and its incomplete compensation by proteins. Thus, the exit of the nascent polypeptide chain from mitoribosome apparently can occur in two different ways.

Mitochondrial ribosomes almost exclusively synthesize hydrophobic polypeptides that form the functional centers of electron transport chain complexes. In order to achieve efficient integration of nascent polypeptides into the inner mitochondrial membrane, translation and, most likely, assembly of the mitoribosomes occur directly on the membrane [43, 44]. The process of anchoring the mitoribosome on the mitochondrial membrane in *S. cerevisiae* is fairly well understood. It involves a number of proteins (Oxa1p, Mba1p, and Mdm38p) as well as mRNA molecules associated with the membrane through translational activators - a group of proteins involved in the organization and regulation of the translational machinery. Oxa1p interacts with the PES via the C-terminal domain of the mitochondrial homolog of bacterial L23 [45, 46], Mba1p plays the role of mitoribosome receptor [47], and Mdm38p together with Mba1p bind the large subunit. Deletion of all genes encoding these proteins does not lead to complete dissociation of the ribosome from the

membrane, indicating the existence of as yet undiscovered components involved in the process [48, 49].

Several recent studies have described unexpected and intimate functional links between the mitochondrial translation and transcription. In human mitochondria, there is a free pool of the mitoribosomal large subunit protein MRPL12, which selectively binds to mitochondrial RNA polymerase *in vivo* and directly stimulates a transcription system *in vitro* [50]. This is considered to be one of the mechanisms coordinating the assembly of mitochondrial ribosomes. On the other hand, human mitochondrial RNA polymerase (POLRMT) has a role in mitoribosomal biogenesis and/or translation that is completely independent of its function in transcription. Co-immunoprecipitation and fractionation of mitochondrial lysates indicate that POLRMT forms a complex with the 28S ribosomal subunit. This complex also includes a paralog of the mitochondrial transcription factor h-mtTFB2, protein h-mtTFB1, which acts as a 12S rRNA methyl transferase [51]. According to one hypothesis, by stimulating the methyl transferase activity of h-mtTFB1 and preventing the association of immature small subunits with 39S subunits, formation of the POLRMT•mitoribosome complex acts as a control point in the assembly path of the small subunit [51].

Below we describe translation in mammalian mitochondria, with differences in the yeast system noted when necessary.

Translation initiation in mitochondria

COMMON FEATURES

During the initiation of protein synthesis, scanning and positioning of the mRNA start codon occurs in the P-site of the small subunit. This is followed by binding of the initiator tRNA followed by association of the large subunit (subunit joining). Like all stages of translation, initiation is assisted by specific protein factors. There are three universal initiation factors bacteria, namely IF1, IF2 and IF3. In mitochondria, IF2_{mt} is universally present and IF3_{mt} has been identified in the vast majority of eukaryotes [52]. Both human factors have been characterized in detail in *in vitro* translation systems [53]. Using *in vitro* experiments with initiator complexes formed with 55S mitoribosomes from bovine liver on *in vitro*-synthesized 5'-terminal fragments of cytochrome oxidase subunit I (COI) and subunit II of NADH dehydrogenase (ND2) mRNAs in the presence of yeast mitochondrial initiator tRNA (fMet-tRNA_i), the following model of mitochondrial translation initiation has been proposed (Fig. 1) [54]. First, IF3_{mt} catalyzes the dissociation of 55S ribosomes, resulting in formation of free 39S subunits and 28S•IF3_{mt} complexes. For convenience, binding of translational GTPase IF2_{mt}•GTP is shown in the next stage, however the exact timing of its attachment to the complex is unknown. Next, mRNA is threaded through the unique gate-shaped

structure in the 28S subunit, followed by the binding of fMet-tRNA_i. This order of events is dictated by IF3_{mt}, which, unlike its bacterial homologue, destabilizes initiator tRNA binding in the absence of mRNA [55]. Primer extension reaction inhibition assays have shown that upon initiation complex formation, the ribosome is stalled after first 17 nucleotides of mammalian mitochondrial mRNAs. From this it was postulated that the mitoribosome scans the 5'-UTR of mRNA in a search for the start codon, and when it is located in the P-site, translational GTPase IF2_{mt} in a GTP-bound form promotes docking of fMet-tRNA_i, with resultant codon-anticodon interactions stabilizing the initiation complex. If the interactions between the start codon and anticodon are not formed, the 28S subunit continues sliding along the mRNA and eventually dissociates. After initiator tRNA binding, the 39S subunit joins the complex, IF2_{mt} hydrolyzes bound GTP to GDP, and the initiator factors leave the mature 55S initiation complex, which is now ready for the elongation stage of polypeptide synthesis.

The scheme of translation initiation described above is applicable only to mammalian mitochondria. Animal mitochondrial mRNAs lack long 5' UTRs (up to 3 nt in case of COI mRNA). The presence of only three additional nucleotides upstream of the start codon in COI mRNA reduces the efficiency of initiation complex formation by 40%, and 12 additional nucleotides at the 5'-side of the AUG start codon lead to an 80% decrease [54]. A very different situation is observed in yeast, where the long 5'-UTRs precede the start codons, and interact with specific activators of translation (see below). Also in yeast mitochondria, unlike mammals, there are two methionine tRNAs: initiator and elongator. It was originally assumed that similarly to the bacterial case, formylation of methionine of the acylated initiator tRNA is strictly necessary for the initiation of translation. Surprisingly, it was shown that in *S. cerevisiae* translation initiation can occur in the absence of a formyl residue of fMet-tRNA_i. An auxiliary factor Aep3p which promotes the binding of non-formylated tRNA_i to IF2_{mt} was discovered recently [56]. This protein is essential when only non-formylated initiator tRNA is available. Aep3p is a peripheral protein of the mitochondrial inner membrane and contains four RNA-binding PPR (penta-tricopeptide) motifs, and in addition to its role in translation initiation, it stabilizes the bicistronic mitochondrial mRNA encoding subunits 6 and 8 of ATP synthase [57]. In mammals, the participation of a single methioninyl-tRNA in initiation or elongation is determined by competition between transformylase and elongation factor EF-Tu_{mt}: affinity of IF2_{mt} to the formylated form is 50 times higher than to non-formylated, while EF-Tu_{mt} does not show detectable affinity to fMet-tRNA_i [58].

The mitochondria of *Saccharomyces* fungi contain special translation activators, most of which are not found in mammalian mitochondria. In *S. cerevisiae* such specific protein factors are involved in translation of each of the eight polypeptides encoded in the mitochondrial genome [59]. Translational activators are encoded in the nuclear genome, and the proteins are transported to the

mitochondrial inner membrane. Activators also coordinate the rate of synthesis of proteins encoded in the mitochondria with their subsequent assembly into the respiratory chain complexes. A detailed description of the translational activators can be found in a recent comprehensive review [60].

Mitochondrial translational initiation factors

As was mentioned above, no mitochondrial ortholog of bacterial IF1 has been found in eukaryotes [61]. In bacteria this small protein (71 amino acid residues in *E. coli*) together with IF3 is involved in fMet-tRNA_i selection on the 30S subunit, blocking binding of the tRNA to the A site of the ribosome and inhibiting 50S subunit joining in the absence of fMet-tRNA. Until recently, it was thought that the absence of IF1 in mitochondria is generally compensated by a short insertion between domains V and VI (in accordance with the 6-domain scheme of IF2) in IF2_{mt} [62]. This hypothesis is supported by the fact that the expression of bovine IF2_{mt} allows *E. coli* cells to grow with a deletion of the gene encoding IF1 [62]. Cryoelectron microscopy reconstruction 70S ribosomes complexed with bovine IF2_{mt} and fMet-tRNA_i in the presence of mRNA and non-hydrolyzable analogue of GTP, GDPNP, suggested that the insertion in IF2_{mt} occupies the same area on the intersubunit surface of the 30S particle as IF1 in a bacterial system [61]. However, bioinformatic analysis showed that this amino acid sequence is only conserved in vertebrates [52]. Therefore, it appears that this phenomenon is not universal and the insertion has evolved secondarily to the loss of IF1_{mt}.

IF2_{mt} is a highly conserved protein, homologous to IF2 of bacteria and archaea, and to the corresponding eukaryotic cytosolic factor eIF5B. By analogy with the 6-domain organization of IF2 in *E. coli*, 4 domains are identified in IF2_{mt}: III, IV, V and VI; the latter is composed of two subdomains (CI and CII) [63]. Domain IV is a classic G-domain - the binding site of guanidinylnucleotides [53]. In the presence of GDPNP, IF2_{mt} binds to the 28S subunit 3 times more efficiently than in the presence of GDP, and 10 times more efficiently than in the absence of nucleotides, whereas the dissociation constant for the complex of 28S•IF2_{mt}, measured in experiments with biotinylated mitoribosomes in the presence of GDPNP, was found to be 10-20 nM [64]. Domain III forms contacts with the mitoribosome small subunit and the VICII domain is involved in binding of the CCA-end of fMet-tRNA_i. The animal-specific insertion between domains V and VICI, that was suggested to play the role of a lost homolog of IF1, contacts helices h18 and h44 of rRNA and the protein S12.

The situation with the third mitochondrial translation initiation factor IF3_{mt} is much more complicated. The low degree of homology with the bacterial factor (the human factor shares 20.8% identity with IF3 of *E. coli*) was for a long time an obstacle to detection of the respective factor in the mitochondria of eukaryotes. An initial search for IF3_{mt} orthologs in the *Homo sapiens* EST

database using *E. coli* IF3 as the query was not successful [65]. However, candidate genes were successfully identified both in human and mouse using *Mycoplasma sp.* and *Euglena gracilis* chloroplast IF3s as queries [65]. Candidates for the role of this factor in the genomes of invertebrates and lower eukaryotes have proved harder to find. By using a sensitive PSI-Blast algorithm, an IF3_{mt} ortholog in yeast *S. cerevisiae* - Aim23p, was found [52]. The growth of yeast cells with the AIM23 gene deletion on a medium with non-fermentable carbon sources was restored in the presence of the gene encoding *S. pombe* IF3_{mt}, which fully confirms the *in silico* results. The mammalian IF3_{mt} is 278 amino acid residues long, and comprises two domains connected by a linker, as does its bacterial ortholog. According to computer modeling, the N-terminal domain of IF3_{mt} (hereinafter N-domain) is composed of α -helical region located opposite a β -layer composed of 4 chains and a second α -helix joining the linker region [66]. The C-terminal domain (hereinafter C-domain) contains two α -helices (H3 and H4) lying on the β -layer consisting of four chains. One of the chains is divided into two segments by a proline residue. IF3_{mt} has N- and C-terminal "tails", 31 and 33 amino acid residues long, respectively. It also has an N-terminal signal peptide that directs import into the mitochondrion and is cleaved to give the mature form upon the protein reaching the matrix. Both tail segments can not be modeled on the basis of currently available structural data, and presumably acquire a certain conformation only upon binding to the small subunit of the mitoribosome. The linker region is partially α -helical, and biochemical studies of the bacterial homologue indicate that the linker is highly flexible [67].

As is the case with the bacterial factor, IF3_{mt} promotes the assembly of the initiator complex on the 28S subunit if the system contains mRNA and IF2_{mt}•GTP. This is achieved through dissociation of active 55S ribosomes to large and small subunits [65]. Unlike its bacterial counterpart, IF3_{mt} does not promote dissociation of non-cognate initiation complexes and the residues critical for this activity are not conserved in the mitochondrial factor [55].

There are functional differences between the N- and C-domains of bacterial and mitochondrial factors. In the case of the bacterial protein the N-domain serves as an additional anchoring point for binding to the ribosome, while the functional activities reside in the C-domain. The two domains are connected by the linker region. The roles of the individual domains of IF3_{mt} in the factor's interactions with the ribosome were elucidated through a series of mutational experiments [66]. The C-domain plus linker region combined has the greatest affinity to the small subunit (K_d of 60 nM vs. 30 nM for the full-length protein). The binding of the N-domain to the 28S subunit is characterized by a larger K_d value of 240 nM. Removal of the linker leads to a decrease in affinity (K_d is 90 and 390 nM, in the case of the full-length protein and N-domain, respectively). Therefore, it was proposed that the linker region modulates IF3_{mt}-mediated dissociation of the ribosome into subunits. Replacements of C-terminal domain amino acid residues

at positions 170, 171 and 175 to alanine lead to a virtually complete loss of the factor's activity in dissociation of the mitoribosome and stimulation of the initiation complex formation, while having little effect the protein's affinity to the small subunit. Most likely, these residues play an important role in blocking the intersubunit bridge B2b.

The roles of N- and C-terminal 'tails' in the IF3_{mt} functional cycle was analyzed in another series of mutational experiments [68]. Surprisingly, removal of these regions leads to a tenfold increase in the affinity of the factor to the large subunit, with the majority of this effect determined by the N-terminal extension. As was mentioned in the previous section, IF3_{mt} has another property which differs from its bacterial homologues, namely dissociation of fMet-tRNA_i from the complex with the 28S subunit in the absence of mRNA. This activity requires the presence of the C-terminal 'tail' and the linker region, with the amino acid residues at positions 247 and 248 playing a key role. According to the available three-dimensional model, the C-terminal extension is directed toward the linker region, which allows these sections of the protein to act together.

There are considerable differences between the IF3 and the IF3_{mt} binding sites on the small subunit, as was determined using chemical cross-linking followed by mass spectrometry [69]. The full-length IF3_{mt} binds to ribosomal proteins MRPS5, MRPS9, MRPS10, MRPS18, which have corresponding homologs in the 70S ribosome, as well as with a group of mitochondria-specific proteins of the small subunit: MRPS29, MRPS32, MRPS36 and PTCD3. The only ribosomal protein cross-linked to the N-domain of IF3_{mt} is MRPS10, located in the head of 28S subunit, with the rest of the detected proteins being cross-linked to the C-domain and linker.

Translation elongation in mitochondria

General features

Comparison among all known translation systems shows that the elongation stage of translation has been particularly well conserved during evolution. The mitochondrial elongation factor GTPase EF-Tu_{mt} (Elongation Factor Thermounstable) forms a ternary complex with GTP and aminoacyl-tRNA, and delivers the latter to the A-site of mitoribosome [70]. Formation of the cognate codon-anticodon interaction promotes GTP hydrolysis by EF-Tu leading to the release of the factor from the ribosome. The guanine nucleotide exchange factor EF-Ts_{mt} (Elongation Factor Thermostable) promotes replacement of GDP with GTP on EF-Tu_{mt}. Next, the ribosome catalyzes the transpeptidation reaction. This results in a pre-translocation complex with the P-site occupied by the deacylated tRNA and the A-site harboring the peptidyl-tRNA that has been extended by one amino acid residue. In the final step of elongation, the mitochondrial GTPase EF-G1 (Elongation Factor G 1) catalyzes the movement of mRNA by one codon in the 5'-3' direction, resulting in deacylated tRNA leaving the ribosome, and the peptidyl-tRNA moving in the P-site, completing the

cycle. The E (Exit) site that transiently binds the deacylated tRNA dissociating from the ribosome is universal for bacterial and eukaryotic cytosolic ribosomes. However, its existence is still being questioned in the case of the mitochondrial ribosomes [30].

Mitochondrial translation elongation factors

EF-Tu_{mt} is extracted from the mitochondrial fraction of bovine liver in a tight complex with EF-Ts_{mt} (Elongation Factor Thermostable) and remains associated in the presence of GDP concentrations up to 1 mM [71]. This is due to the balance of EF-Tu_{mt} affinities to G nucleotides and EF-Ts_{mt}: the Kds for binding of GTP and GDP are 1 and 18 mM, respectively, while binding to EF-Ts_{mt} is considerably tighter with Kd of 5.5 nM [72]. Both factors are present in mitochondria *in vivo* in a stoichiometric ratio of 1:1, with around 400 copies per mitochondrion. This is different from the pattern in bacteria, where the ratio is 8:1 and $\approx$ 200,000 molecules, respectively [73]. A mitochondrial localization signal was determined for both proteins by sequencing the N-terminus in purified preparations of isolated mitochondria [74].

X-ray crystallography of EF-Tu_{mt}•GDP complex shows a three-domain organization scheme, typical of bacterial homologs [75]. Domain I consists of 8 α -helices and 8 β -strands that form the binding pocket for guanyl nucleotides. A flexible linker of 11 amino acid residues joins the G-domain with domain II, which is formed by 11 β -strands folded in a "Greek key" motif. Domain III folds into a "jellyroll" structure, formed by a β -layer of 6 sheets. A unique feature of mitochondrial EF-Tu is the presence of a C-terminal extension. In humans and *Bos taurus*, its length is 11, and in *C. elegans* - 57 amino acid residues. It is thought that this extension in nematodes is required for the formation of the ternary complex with tRNAs, which typically have unusually short T-stems and a complete absence of the D-stem in these organisms. However, the function of the extension in mammals is associated with the binding of aminoacyl-tRNA to the small ribosomal subunit [75].

The binding pocket for the acceptor arm of aminoacyl-tRNA is formed at the junction between domains I and II / III of EF-Tu_{mt}. Interestingly, the mitochondrial factor is active in a system with bacterial tRNAs, while EF-Tu from *E. coli* is not able to participate in elongation using mitochondrial tRNAs [76, 77]. The formation of a correct codon-anticodon duplex in the A site sends a signal through the tRNA molecule to the G-domain of EF-Tu that leads to hydrolysis of GTP [78]. Presumably, the shorter mitochondrial tRNA molecules in complex with bacterial EF-Tu are not correctly positioned on the ribosome and can not effectively stimulate nucleotide hydrolysis.

In addition to participating in translation elongation, EF-Tu_{mt} acts as a chaperone in the process of quality control of mitochondrial proteins [79]. EF-Tu_{mt} binds to unfolded or misfolded newly synthesized polypeptides and directs them to the protease complex, where they are degraded.

The low degree of sequence similarity of EF-Ts_{mt} with its bacterial homologs (25-35%) has prevented reliable predictions from being made about its three-dimensional structure. Bacterial EF-Ts consists of four domains: the N-terminal, a core domain (divided by N- and C-subdomains), a dimerization domain with a «coiled-coil» fold, and a C-terminal module [80]. The X-ray structure of the EF-Tu_{mt}•EF-Ts_{mt} complex has revealed dramatic differences from the organization outlined above [81]. The most conserved part of the protein - the N-domain - is assembled from three α -helices similarly to the bacterial homolog. The core domains of the two proteins share the "β-sandwich" fold, although the number the organization of the β-strands varies. The most dramatic difference between the two proteins is a reduction of the dimerization domain to large loop folded opposite to the C-subdomain and the complete absence of the C-terminal module in the mitochondrial counterpart [81].

Interactions between EF-Ts_{mt} and EF-Tu_{mt}•GDP lead to large-scale conformational changes in the latter, which is not the case for the corresponding *E. coli* and *T. thermophilus* complexes. GDP to GTP exchange is accompanied by significant conformational rearrangements in EF-Tu_{mt} leading to the disruption of the G-nucleotide binding site, which presumably occurs in three steps [82 – 84]. First, the Mg²⁺ ion binding site is disrupted, followed by the destabilization of EF-Tu_{mt} interactions with the oxygen atoms of the GDP β-phosphate moiety, and, lastly, contacts with the ribose and purine ring are lost.

The equivalent of bacterial EF-Ts is absent in budding yeast *S. cerevisiae*, however, is found in fission yeast *Schizosaccharomyces pombe*. In this respect at least, the fission yeast mitochondrial system is more similar to that of humans than is budding yeast. EF-Tu_{mt} in *S. cerevisiae* as profoundly different affinities to for GTP and GDP in comparison to that of *E. coli* EF-Tu (5 and 25 μM vs 300 and 3 nM, respectively) and, therefore, does not require stabilization by GDP [85]. The functional link between the absence of EF-Ts_{mt} and low affinities to G nucleotides of *S. cerevisiae* EF-Tu_{mt} was demonstrated by mutational analysis [86]. Single point mutations in *S. pombe* EF-Tu_{mt} lowering its affinities to G nucleotides also lead to its independence from the EF-Ts_{mt} rendering *S. pombe* EF-Tu_{mt} functionally similar to its *S. cerevisiae* counterpart.

EF-G catalyses the translocation of peptidyl-tRNA from the A- to the P-site of the ribosome. In contrast to bacteria in which a single EF-G acts both during elongation and ribosomal recycling, in most eukaryotes the two functions are split between two mitochondrial factors, EF-G1_{mt} and EF-G2_{mt} [87, 88]. These two factors have amino acid sequence identity of 35% in average [87]. EF-G1_{mt} has translocation activity with both 55S mitoribosomes and 70S bacterial monosomes. Bacterial EF-G, however, is incompatible with mitochondrial ribosomes. Experiments on hybrid ribosomes with heterologous ribosomal proteins L7/L12 demonstrated that the specificity lies in the interactions with these proteins [89].

Despite the fact that EF-G1_{mt} was isolated from bovine liver many years ago, it is still one of the most poorly characterized proteins of the mitochondrial translation apparatus [90]. As with the bacterial protein, EF-G1_{mt} can be structurally divided into 5 domains, and its shape is strikingly similar to that of the ternary complex EF-Tu•GDPNP•Phe-tRNA^{Phe} [91]. Despite the fact that the amino acid residues involved in EF-G's interaction with antibiotic fusidic acid are present in the mitochondrial factor, the antibiotic does not significantly inhibit EF-G1_{mt} both in the presence of bacterial and mitochondrial ribosomes [90].

Translation termination and recycling in the mitochondria

The third and fourth stages of protein synthesis in mitochondria occur when the stop codon of mRNA is present in the ribosomal A-site and is decoded by a termination, or Release Factor, RF. The stop codon is recognized by RF1_{mt} (Release Factor 1), which promotes hydrolysis of the ester bond between P-site tRNA and the newly synthesized polypeptide, releasing the completed protein [92]. The cleavage occurs in the peptidyl transferase center of the large subunit. The mechanism of RF1_{mt} release from its complex with mitoribosomes is still unknown. Next, two factors – RRF1_{mt} (Ribosome Recycling Factor) and EF-G2_{mt} in complex with GTP – jointly promote the dissociation of tRNA and mRNA from the ribosome and split the ribosomal particles into subunits (Fig. 3), setting the stage for the re-use of the ribosome in the next round of translation. GTP hydrolysis by EF-G2_{mt} is required for the release of RRF1_{mt} and EF-G2_{mt} from the ribosome [92].

Until recently, it was widely believed that human mitochondria have an extended repertoire of termination codons in comparison with the bacterial system [93]. Two open reading frames are terminated by non-standard triplets: AGA and AGG in genes MTCOI and MTND6, respectively. However, recent investigations suggest that the 55S mitoribosome undergoes a -1 frameshift, which results in a standard UAG stop codon that is recognized by RF1_{mt} in the A site [94].

There are two classes of termination factors in bacteria, archaea and the eukaryotic cytosol: class-I and class-II [95]. The class-I factors are involved in direct recognition of the stop codon and release of the peptide, whereas class-II factors facilitate the release of the class-I factors from the ribosome. Class-II factors have not been found in mitochondria. In archaea and in the cytoplasm of eukaryotes all three stop codons are recognized by a single class-I release factor – aRF1 and eRF1, respectively; and in bacteria – by two proteins, RF1 and RF2. The UAA codon is recognized by both bacterial factors; RF1 specifically recognizes UAG, and RF2 – UGA [96, 97]. Bioinformatic analyses have showed that the mitochondrial factors have greatest sequence similarity to the bacterial RF1 [98].

The three-dimensional structure of human RF1_{mt} has been modeled using the available structure of *Thermus thermophilus* RF1 [92]. It was found that all protein domains and motifs

typical for this class are present in the mitochondrial factor: the PXT motif of domain 2 (PKT in the case of the mitochondrial factor) participating in stop codon recognition, the 5 α -helix tip of domain 4, as well as the highly conserved GGQ motif of domain 3 that catalyzes the hydrolysis of an ester bond between the tRNA and the newly synthesized peptide [99]. Moreover, the arrangement of functional sites is compatible with the "open" conformation of the factor that is required for its functional activity on the ribosome [100]. Two additional proteins possessing the RF-specific GGQ motif are present in human mitochondria in addition to RF1_{mt}, namely ICT1 and C12 [101, 102]. ICT1 has ribosome-dependent, A-site-codon-independent peptidyl hydrolase activity. This protein is thought to promote the release of peptides from ribosomes "stuck" on mRNAs lacking a stop codon, thus performing a function analogous to tmRNA in bacteria [103]. Co-immunoprecipitation experiments show that ICT1 is tightly associated with mitoribosomal large subunits, and functionally active in a heterologous system with bacterial 70S ribosomes.

As mentioned in the previous chapter, there are two forms of the protein EF-G, EF-G1_{mt} and EF-G2_{mt}, in the mitochondria of most eukaryotes. The former appears to be responsible for translocation, whereas the latter is involved in ribosomal recycling. This view is supported by recent *in vivo* results using an *E. coli* strain carrying a temperature-sensitive RRF variant. In this strain, EF-G2_{mt} in combination with RRF_{mt} is able to restore normal growth at non-permissive temperatures [103]. Molecular phylogenetic analysis of the EF-G protein family suggests an asymmetric separation of functional activities between the two EF-Gs: EF-G2_{mt} is highly divergent, primarily conserving sites that maintain an EF-G-like architecture and ribosome-binding capabilities, while EF-G1_{mt} has experienced substantial and well conserved changes in the G-domain that suggest a modified mechanism of interaction with nucleotides [104].

The ribosomal recycling stage in mitochondria differs from the bacterial system in another important aspect: GTP hydrolysis by EF-G2_{mt} appears to occur after mitoribosome dissociation into subunits and is required for dissociation of EF-G2_{mt} and RRF_{mt} from the large subunit, allowing the subunits to associate again in the absence of IF3_{mt} [103].

Conclusions and perspectives

In this review we have attempted to compile the available data on protein synthesis in the mitochondria of eukaryotic cells. Our current knowledge of protein synthesis in mitochondria is still rather limited in comparison to that of bacterial and cytosolic translation in eukaryotes. Atomic resolution structures of the mitochondrial ribosome are lacking, and our biochemical understanding of the mitochondrial translation lacks kinetic details. Although mitochondrial translation is generally organized similarly to that in bacteria it is characterized by a number of significant differences. In recent years numerous scientific reports have elucidated these differences, reigniting

the interest of the scientific community.

Acknowledgements

The authors are grateful to the members of P. Kamenski, V. Hauryliuk and N. Zenkin laboratories for their assistance in writing this review. Research in the Kamenski lab was supported by the funding from the International Associated Laboratory «RNA-mitocure», Russian Foundation for Basic Research and Russian Ministry of Education and Science (contract 8817); in the Hauryliuk lab – by Estonian Science Foundation grants (ETF9012 and PUT37) and European Regional Development Fund (RLOTITIPP), and in the N. Zenkin lab – by the UK Biotechnology and Biosciences Research Council and by European Research Council (ERC-2007-StG 202994-MTP). A. Kuzmenko is supported by Archimedes Foundation, and by the “U.M.N.I.K” program. G. C. Atkinson is supported by Estonian Science Foundation grant ETF9020 and European Social Fund grant «Mobilitas» MJD99.

References

1. Yang, D., Oyaizu, Y., Oyaizu, H., Olsen, G. J., and Woese, C. R. (1985) *Proc Natl Acad Sci U S A* **82**, 4443-4447
2. Leon, S. A., and Mahler, H. R. (1968) *Arch Biochem Biophys* **126**, 305-319
3. Linnane, A. W., Lamb, A. J., Christodoulou, C., and Lukins, H. B. (1968) *Proc Natl Acad Sci U S A* **59**, 1288-1293
4. Wagner, R. P. (1969) *Science* **163**, 1026-1031
5. Watanabe, K. (2010) *Proc Jpn Acad Ser B Phys Biol Sci* **86**, 11-39
6. Christian, B. E., and Spremulli, L. L. (2012) *Biochim Biophys Acta* **1819**, 1035-1054
7. Melnikov, S., Ben-Shem, A., Garreau de Loubresse, N., Jenner, L., Yusupova, G., and Yusupov, M. (2012) *Nat Struct Mol Biol* **19**, 560-567
8. Gray, M. W. (2012) *Cold Spring Harb Perspect Biol* **4**, a011403
9. Wilson, R. J., and Williamson, D. H. (1997) *Microbiol Mol Biol Rev* **61**, 1-16
10. Anderson, S., Bankier, A. T., Barrell, B. G., de Bruijn, M. H., Coulson, A. R., Drouin, J., Eperon, I. C., Nierlich, D. P., Roe, B. A., Sanger, F., Schreier, P. H., Smith, A. J., Staden, R., and Young, I. G. (1981) *Nature* **290**, 457-465
11. Unseld, M., Marienfeld, J. R., Brandt, P., and Brennicke, A. (1997) *Nat Genet* **15**, 57-61
12. Gray, M. W., Lang, B. F., Cedergren, R., Golding, G. B., Lemieux, C., Sankoff, D., Turmel, M., Brossard, N., Delage, E., Littlejohn, T. G., Plante, I., Rioux, P., Saint-Louis, D., Zhu, Y., and Burger, G. (1998) *Nucleic Acids Res* **26**, 865-878
13. Brown, W. M., George, M., Jr., and Wilson, A. C. (1979) *Proc Natl Acad Sci U S A* **76**,

1967-1971

14. Jacob, J. E., Vanholme, B., Van Leeuwen, T., and Gheysen, G. (2009) *BMC Res Notes* **2**, 192
15. Osawa, S., Jukes, T. H., Watanabe, K., and Muto, A. (1992) *Microbiol Rev* **56**, 229-264
16. Masters, B. S., Stohl, L. L., and Clayton, D. A. (1987) *Cell* **51**, 89-99
17. Tiranti, V., Savoia, A., Forti, F., D'Apolito, M. F., Centra, M., Rocchi, M., and Zeviani, M. (1997) *Hum Mol Genet* **6**, 615-625
18. Ringel, R., Sologub, M., Morozov, Y. I., Litonin, D., Cramer, P., and Temiakov, D. (2011) *Nature* **478**, 269-273
19. Ojala, D., Montoya, J., and Attardi, G. (1981) *Nature* **290**, 470-474
20. Temperley, R. J., Wydro, M., Lightowlers, R. N., and Chrzanowska-Lightowlers, Z. M. (2010) *Biochim Biophys Acta* **1797**, 1081-1085
21. Montoya, J., Ojala, D., and Attardi, G. (1981) *Nature* **290**, 465-470
22. Groot, G. S., Flavell, R. A., Van Ommen, G. J., and Grivell, L. A. (1974) *Nature* **252**, 167-169
23. Grohmann, K., Amairic, F., Crews, S., and Attardi, G. (1978) *Nucleic Acids Res* **5**, 637-651
24. Dunstan, H. M., Green-Willms, N. S., and Fox, T. D. (1997) *Genetics* **147**, 87-100
25. Dziembowski, A., Piwowarski, J., Hoser, R., Minczuk, M., Dmochowska, A., Siep, M., van der Spek, H., Grivell, L., and Stepien, P. P. (2003) *J Biol Chem* **278**, 1603-1611
26. Wang, G., Chen, H. W., Oktay, Y., Zhang, J., Allen, E. L., Smith, G. M., Fan, K. C., Hong, J. S., French, S. W., McCaffery, J. M., Lightowlers, R. N., Morse, H. C., 3rd, Koehler, C. M., and Teitell, M. A. (2010) *Cell* **142**, 456-467
27. Suzuki, T., and Nagao, A. (2011) *Annu Rev Genet* **45**, 299-329
28. Helm, M., Brule, H., Friede, D., Giege, R., Putz, D., and Florentz, C. (2000) *RNA* **6**, 1356-1379
29. Yokogawa, T., Watanabe, Y., Kumazawa, Y., Ueda, T., Hirao, I., Miura, K., and Watanabe, K. (1991) *Nucleic Acids Res* **19**, 6101-6105
30. Sharma, M. R., Koc, E. C., Datta, P. P., Booth, T. M., Spremulli, L. L., and Agrawal, R. K. (2003) *Cell* **115**, 97-108
31. Kamenski, P., Kolesnikova, O., Jubenot, V., Entelis, N., Krashenninnikov, I. A., Martin, R. P., and Tarassov, I. (2007) *Mol Cell* **26**, 625-637
32. Hancock, K., and Hajduk, S. L. (1990) *J Biol Chem* **265**, 19208-19215
33. Schneider, A. (2011) *Annu Rev Biochem* **80**, 1033-1053
34. O'Brien, T. W. (1971) *J Biol Chem* **246**, 3409-3417
35. Maslov, D. A., Sharma, M. R., Butler, E., Falick, A. M., Gingery, M., Agrawal, R. K.,

- Spremulli, L. L., and Simpson, L. (2006) *Mol Biochem Parasitol* **148**, 69-78
36. Borst, P., and Grivell, L. A. (1971) *FEBS Lett* **13**, 73-88
37. Agrawal, R. K., and Sharma, M. R. (2012) *Curr Opin Struct Biol* **22**, 797-803
38. Smirnov, A., Entelis, N., Martin, R. P., and Tarassov, I. (2011) *Genes Dev* **25**, 1289-1305
39. Entelis, N. S., Kolesnikova, O. A., Dogan, S., Martin, R. P., and Tarassov, I. A. (2001) *J Biol Chem* **276**, 45642-45653
40. Agrawal RK, S. M., Yassin A, Lahiri I, Spremulli LL. (2011) *Structure and function of organellar ribosomes as revealed by cryo-EM*, Springer Wien, New York
41. Gruschke, S., and Ott, M. (2010) *Bioessays* **32**, 1050-1057
42. Sharma, M. R., Booth, T. M., Simpson, L., Maslov, D. A., and Agrawal, R. K. (2009) *Proc Natl Acad Sci U S A* **106**, 9637-9642
43. Watson, K. (1972) *J Cell Biol* **55**, 721-726
44. Fiori, A., Mason, T. L., and Fox, T. D. (2003) *Eukaryot Cell* **2**, 651-653
45. Szyrach, G., Ott, M., Bonnefoy, N., Neupert, W., and Herrmann, J. M. (2003) *EMBO J* **22**, 6448-6457
46. Jia, L., Dienhart, M., Schramp, M., McCauley, M., Hell, K., and Stuart, R. A. (2003) *EMBO J* **22**, 6438-6447
47. Ott, M., Prestele, M., Bauerschmitt, H., Funes, S., Bonnefoy, N., and Herrmann, J. M. (2006) *EMBO J* **25**, 1603-1610
48. Bauerschmitt, H., Mick, D. U., Deckers, M., Vollmer, C., Funes, S., Kehrein, K., Ott, M., Rehling, P., and Herrmann, J. M. (2010) *Mol Biol Cell* **21**, 1937-1944
49. Frazier, A. E., Taylor, R. D., Mick, D. U., Warscheid, B., Stoepel, N., Meyer, H. E., Ryan, M. T., Guiard, B., and Rehling, P. (2006) *J Cell Biol* **172**, 553-564
50. Surovtseva, Y. V., Shutt, T. E., Cotney, J., Cimen, H., Chen, S. Y., Koc, E. C., and Shadel, G. S. (2011) *Proc Natl Acad Sci U S A* **108**, 17921-17926
51. Surovtseva, Y. V., and Shadel, G. S. (2013) *Nucleic Acids Res* **41**, 2479-2488
52. Atkinson, G. C., Kuzmenko, A., Kamenski, P., Vysokikh, M. Y., Lakunina, V., Tankov, S., Smirnova, E., Soosaar, A., Tenson, T., and Hauryliuk, V. (2012) *Nucleic Acids Res* **40**, 6122-6134
53. Spremulli, L. L., Coursey, A., Navratil, T., and Hunter, S. E. (2004) *Prog Nucleic Acid Res Mol Biol* **77**, 211-261
54. Christian, B. E., and Spremulli, L. L. (2010) *J Biol Chem* **285**, 28379-28386
55. Bhargava, K., and Spremulli, L. L. (2005) *Nucleic Acids Res* **33**, 7011-7018
56. Lee, C., Tibbetts, A. S., Kramer, G., and Appling, D. R. (2009) *J Biol Chem* **284**, 34116-34125

57. Ellis, T. P., Helfenbein, K. G., Tzagoloff, A., and Dieckmann, C. L. (2004) *J Biol Chem* **279**, 15728-15733
58. Spencer, A. C., and Spremulli, L. L. (2004) *Nucleic Acids Res* **32**, 5464-5470
59. Towpik, J. (2005) *Cell Mol Biol Lett* **10**, 571-594
60. Herrmann, J. M., Woellhaf, M. W., and Bonnefoy, N. (2013) *Biochim Biophys Acta* **1833**, 286-294
61. Yassin, A. S., Haque, M. E., Datta, P. P., Elmore, K., Banavali, N. K., Spremulli, L. L., and Agrawal, R. K. (2011) *Proc Natl Acad Sci U S A* **108**, 3918-3923
62. Gaur, R., Grasso, D., Datta, P. P., Krishna, P. D., Das, G., Spencer, A., Agrawal, R. K., Spremulli, L., and Varshney, U. (2008) *Mol Cell* **29**, 180-190
63. Laalami, S., Sacerdot, C., Vachon, G., Mortensen, K., Sperling-Petersen, H. U., Cenatiempo, Y., and Grunberg-Manago, M. (1991) *Biochimie* **73**, 1557-1566
64. Spencer, A. C., and Spremulli, L. L. (2005) *Biochim Biophys Acta* **1750**, 69-81
65. Koc, E. C., and Spremulli, L. L. (2002) *J Biol Chem* **277**, 35541-35549
66. Christian, B. E., and Spremulli, L. L. (2009) *Biochemistry* **48**, 3269-3278
67. Hua, Y., and Raleigh, D. P. (1998) *J Mol Biol* **278**, 871-878
68. Haque, M. E., Grasso, D., and Spremulli, L. L. (2008) *Nucleic Acids Res* **36**, 589-597
69. Haque, M. E., Koc, H., Cimen, H., Koc, E. C., and Spremulli, L. L. (2011) *Biochim Biophys Acta* **1814**, 1779-1784
70. Smits, P., Smeitink, J., and van den Heuvel, L. (2010) *J Biomed Biotechnol* **2010**, 737385
71. Schwartzbach, C. J., and Spremulli, L. L. (1989) *J Biol Chem* **264**, 19125-19131
72. Cai, Y. C., Bullard, J. M., Thompson, N. L., and Spremulli, L. L. (2000) *Protein Sci* **9**, 1791-1800
73. Worlax, V. L., Bullard, J. M., Ma, L., Yokogawa, T., and Spremulli, L. L. (1997) *Biochim Biophys Acta* **1352**, 91-101
74. Worlax, V. L., Burkhardt, W., and Spremulli, L. L. (1995) *Biochim Biophys Acta* **1264**, 347-356
75. Andersen, G. R., Thirup, S., Spremulli, L. L., and Nyborg, J. (2000) *J Mol Biol* **297**, 421-436
76. Hunter, S. E., and Spremulli, L. L. (2004) *Mitochondrion* **4**, 21-29
77. Bullard, J. M., Cai, Y. C., Zhang, Y., and Spremulli, L. L. (1999) *Biochim Biophys Acta* **1446**, 102-114
78. Piepenburg, O., Pape, T., Pleiss, J. A., Wintermeyer, W., Uhlenbeck, O. C., and Rodnina, M. V. (2000) *Biochemistry* **39**, 1734-1738
79. Suzuki, H., Ueda, T., Taguchi, H., and Takeuchi, N. (2007) *J Biol Chem* **282**, 4076-4084

80. Kawashima, T., Berthet-Colominas, C., Wulff, M., Cusack, S., and Leberman, R. (1996) *Nature* **379**, 511-518
81. Jeppesen, M.G., Navratil, T., Spremulli, L.L., and Nyborg, J. (2005) *J Biol Chem* **280**, 5071-5081.
82. Zhang, Y., Li, X., and Spremulli, L. L. (1996) *FEBS Lett* **391**, 330-332
83. Zhang, Y., Sun, V., and Spremulli, L. L. (1997) *J Biol Chem* **272**, 21956-21963
84. Zhang, Y., and Spremulli, L. L. (1998) *J Biol Chem* **273**, 28142-28148
85. Rosenthal, L. P., and Bodley, J. W. (1987) *J Biol Chem* **262**, 10955-10959
86. Chiron, S., Suleau, A., and Bonnefoy, N. (2005) *Genetics* **169**, 1891-1901
87. Hammarsund, M., Wilson, W., Corcoran, M., Merup, M., Einhorn, S., Grander, D., and Sangfelt, O. (2001) *Hum Genet* **109**, 542-550
88. Tsuboi, M., Morita, H., Nozaki, Y., Akama, K., Ueda, T., Ito, K., Nierhaus, K. H., and Takeuchi, N. (2009) *Mol Cell* **35**, 502-510
89. Terasaki, M., Suzuki, T., Hanada, T., and Watanabe, K. (2004) *J Mol Biol* **336**, 331-342
90. Chung, H. K., and Spremulli, L. L. (1990) *J Biol Chem* **265**, 21000-21004
91. al-Karadaghi, S., Aevansson, A., Garber, M., Zheltonosova, J., and Liljas, A. (1996) *Structure* **4**, 555-565
92. Chrzanowska-Lightowlers, Z. M., Pajak, A., and Lightowlers, R. N. (2011) *J Biol Chem* **286**, 34479-34485
93. Richter, R., Pajak, A., Dennerlein, S., Rozanska, A., Lightowlers, R. N., and Chrzanowska-Lightowlers, Z. M. (2010) *Biochem Soc Trans* **38**, 1523-1526
94. Temperley, R., Richter, R., Dennerlein, S., Lightowlers, R. N., and Chrzanowska-Lightowlers, Z. M. (2010) *Science* **327**, 301
95. Loh, P. G., and Song, H. (2010) *Curr Opin Struct Biol* **20**, 98-103
96. Dontsova, M., Frolova, L., Vassilieva, J., Piendl, W., Kisselev, L., and Garber, M. (2000) *FEBS Lett* **472**, 213-216
97. Ito, K., Frolova, L., Seit-Nebi, A., Karamyshev, A., Kisselev, L., and Nakamura, Y. (2002) *Proc Natl Acad Sci U S A* **99**, 8494-8499
98. Soleimanpour-Lichaei, H. R., Kuhl, I., Gaisne, M., Passos, J. F., Wydro, M., Rorbach, J., Temperley, R., Bonnefoy, N., Tate, W., Lightowlers, R., and Chrzanowska-Lightowlers, Z. (2007) *Mol Cell* **27**, 745-757
99. Trobro, S., and Aqvist, J. (2009) *Biochemistry* **48**, 11296-11303
100. Laurberg, M., Asahara, H., Korostelev, A., Zhu, J., Trakhanov, S., and Noller, H. F. (2008) *Nature* **454**, 852-857
101. Richter, R., Rorbach, J., Pajak, A., Smith, P. M., Wessels, H. J., Huynen, M. A., Smeitink, J.

- A., Lightowlers, R. N., and Chrzanowska-Lightowlers, Z. M. (2010) *EMBO J* **29**, 1116-1125
102. Antonicka, H., Ostergaard, E., Sasarman, F., Weraarpachai, W., Wibrand, F., Pedersen, A. M., Rodenburg, R. J., van der Knaap, M. S., Smeitink, J. A., Chrzanowska-Lightowlers, Z. M., and Shoubridge, E. A. (2010) *Am J Hum Genet* **87**, 115-122
103. Haque, M. E., and Spremulli, L. L. (2010) *EMBO J* **29**, 1019-1020
104. Atkinson, G. C., and Baldauf, S. L. (2011) *Mol Biol Evol* **28**, 1281-1292

	Bacteria	Bovine mitochondria	<i>Leishmania</i> mitochondria
Molecular weight, MDa	2.3	2.7	2.2
Diameter, Å	≈ 260	≈ 320	≈ 245
Molar ratio RNA:protein	≈ 2:1	≈ 1:2	≈ 1:3
Small subunit rRNA and its size	16S (1542 nt)	12S (950 nt)	9S (610 nt)
Number of proteins in small subunit	21	≈ 29	≈ 56
Large subunit rRNA and its size	23S rRNA (2904 nt) 5SrRNA (120 nt)	16SrRNA (1560 nt)	12S rRNA (1173 nt)
Number of proteins in large subunit	34	≈ 50	≈ 77

Table 1. General features of mitochondrial and bacterial ribosomes.

Figure 1. Schematic representation of translation initiation in mitochondria. SRS – small ribosome subunit, LRS – large ribosomal subunit. See text for detailed information.

Figure 2. Schematic representation of translation elongation in mitochondria. SRS – small ribosome subunit, LRS – large ribosomal subunit. See text for detailed information.

Figure 3. Schematic representation of translation termination in mitochondria. SRS – small ribosome subunit, LRS – large ribosomal subunit. See text for detailed information.