Gao G, Clare AS, Rose C, Caldwell GS.

[Reproductive sterility increases the capacity to exploit the green seaweed](#)

[Ulva rigida for commercial applications.](#)

Algal Research 2017, 24(Part A), 64-71.

Copyright:

© 2017. This manuscript version is made available under the [CC-BY-NC-ND 4.0 license](#)

DOI link to article:

<https://doi.org/10.1016/j.algal.2017.03.008>

Date deposited:

16/03/2017

Embargo release date:

22 March 2018

This work is licensed under a

[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International licence](#)

Reproductive sterility increases the capacity to exploit the green seaweed *Ulva rigida* for commercial applications

Guang Gao^{a,#}, Anthony S. Clare^a, Craig Rose^b, and Gary S. Caldwell^{a,*}

^aSchool of Marine Science and Technology, Ridley Building, Newcastle University, Newcastle upon Tyne, NE1 7RU, England, UK

^bSeaweed & Co. Ltd, Office 2.3 North Tyneside Business Centre, 54a Saville Street, North Tyneside, NE30 1NT, England, UK

[#]Current address: Marine Resources Development Institute of Jiangsu, Huaihai Institute of Technology, Lianyungang 222005, China

*Correspondence: Gary Caldwell, tel. +44 (0)191 208 6660, e-mail: gary.caldwell@ncl.ac.uk

Abstract

Periodic biomass losses caused by reproduction have greatly limited the expansion of the land based cultivation of the seaweed *Ulva*. In an attempt to resolve this, we obtained a sterile mutant (SM) of *U. rigida* by mutating a wild type (WT) strain using ultraviolet radiation (UVC). The SM grew five times faster than the WT with 40.0% and 30.9% higher nitrate and phosphate uptake rates. Crucially, the SM remained in the vegetative state throughout the 27 day trial, unlike the WT. The lipid content was more than double that of the WT with more monounsaturated, polyunsaturated, omega-3 and omega-6 fatty acids but with a 26.3% lower protein content. The swelling, water holding, and oil holding capacities were all lower than the WT. The rapidly growing SM showed distinct promise for application to carbon capture and wastewater bioremediation and for more conventional aquaculture practices. Despite some modified nutritional and functional properties, the SM still retained a desirable nutritional profile.

Keywords: aquaculture, mutation, eutrophication, green tide, UV-radiation

1. Introduction

The green seaweed *Ulva* is an increasingly important food (Ortiz et al., 2006), feed (Ahmed et al., 2015; Cyrus et al., 2015), and biofuel feedstock (Bruhn et al., 2011), as well playing a role in the delivery of crucial wastewater and CO₂ remediation services (Chung et al., 2011; Al-Hafedh et al., 2015). Interest in *Ulva* as a source of bioproducts and bioprocesses has driven concomitant efforts to optimize *Ulva* cultivation (Hiraoka & Oka, 2008; Bolton et al., 2009; Carl et al., 2016). *Ulva* species should, in theory, be ideal cultivation candidates given their cosmopolitan distribution (Kirkendale et al., 2013), very high growth rates (Bruhn et al., 2011), and broad environmental tolerances (Luo & Liu, 2011). However, previous studies (for example, Oza & Sreenivasa Rao, 1977; DeBusk et al., 1986; Ale et al., 2011; Castelar et al., 2014) have experienced difficulty in maintaining *Ulva* species in the vegetative state. The formation and release of reproductive cells effectively terminates *Ulva*

growth and leads to a disintegration of part or all of the thallus, dramatically reducing *Ulva* productivity (Oza & Sreenivasa Rao, 1977).

There is thus an imperative to develop robust *Ulva* strains exhibiting traits of either reduced or absent reproduction. Several strain/trait optimization methods are currently available for seaweeds, the simplest being the selection of strains that preferentially express somatic growth over gamete production from the existing wild populations (Van Der Meer, 1983). For example, fast growing clones of *Eucheuma* (Doty & Alvarez, 1975) and *Gigartina exasperata* (Waaland, 1978) have been discovered and distributed to farmers. In terms of *Ulva* species, a sterile *U. pertusa* mutant was discovered at Omura Bay, Japan in 1973, which underwent vegetative development without sexual reproduction and could be maintained under axenic laboratory conditions (Migita, 1985; Murase et al., 1994; Kakinuma et al., 2006). The sterile strain also had a higher photosynthetic efficiency and growth rate, and also had a higher total free amino acid and total carbon and nitrogen content.

Although it is straightforward to look for robust strains from wild populations, the process can be time and labour intensive as it requires the collection of a wide range of ecotypes. Instead, mutagenesis and mutant strain screening can be used to avoid the process. Mutagenesis has successfully been employed for seaweeds (Niwa et al., 2011; Fu et al., 2014), with physical mutagens such as ultraviolet light (UV) widely used (Charrier et al. 2015). These studies demonstrate that developing sterile, or at least low level reproductive strains is technically achievable and would be beneficial for long-term cultivation.

To date, sterile strains of *U. rigida* (a ubiquitous green seaweed implicated in green tide events) have not been reported. This study aimed to develop a sterile strain that would demonstrate high growth potential. To achieve that, varying doses of UV radiation were employed to mutate a wild strain of *U. rigida*. In addition, patterns in growth, nutrient uptake, and chemical composition were examined to shed light on whether the sterile strain can be used for bioremediation and food purposes.

2. Materials and methods

2.1. Plant collection

Vegetative *U. rigida* fronds of 50–60 mm in length were collected from the low intertidal zone of Cullercoats Bay, Tyne and Wear, UK (55.03° N, 1.43° W) after a spring tide in August, 2013. The fronds were placed in a plastic bag and transported to the laboratory within one hour where they were gently rinsed with one micrometer filtered natural seawater to remove any sediment, epiphytes or small grazers.

2.2. Mutation of *Ulva gametes* and isolation of sterile mutants

Gametes were obtained by inducing reproduction from 2.5 mm discs of *U. rigida* thallus (Hiraoka and Enomoto, 1998). One hundred discs were rinsed with autoclaved seawater and transferred to a flask containing 200 mL of autoclaved seawater. The flask was held at 4 °C in darkness for six hours and then transferred to an 18 °C incubator with 80 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ light intensity and 16L: 8D photoperiod. Gametes were released three days later, attracted to a point source of white light (gametes being phototactic) and collected by pipette. Gametes were further concentrated by centrifugation (1000 g, five minutes) and then transferred to 85 mm diameter Petri dishes with the final density of 2×10^7 gametes per dish. Penicillin G and streptomycin sulfate (final concentrations of 1000 and 250 mg L⁻¹ respectively) were added to restrict bacterial growth. Six dishes were exposed to UV lamps (40 W, 254 nm) for 18, 36, 54, 72, 90 and 108 seconds (one dish per treatment). These exposures were chosen as no gametes survived when exposed for more than 120 seconds in preliminary experiments (data not shown). Afterwards, the dishes were placed in darkness at 18 °C for 24 hours to allow gamete settlement. Germlings from each dish were detached (112, 486 individuals in total) and transferred to six tanks (13 L) when they reached the length of 1 cm. The cultures were aerated with a flow rate of 100 L min⁻¹ to suspend the germlings. The seawater was enriched with 1000 $\mu\text{mol L}^{-1}$ nitrate and 50 $\mu\text{mol L}^{-1}$ phosphate and was exchanged every 3–4 days to avoid nutrient limitation. The culture temperature was 18 ± 1 °C and light intensity was 80 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ with a 16L: 8D photoperiod. Over the following four months

any thalli showing reproductive features—the colour of thalli changing from green to yellowish—were removed and those that maintained the vegetative state were cultivated further. Thalli that failed to demonstrate any reproductive tendencies were identified as sterile. Two sterile individuals were obtained, one of which was used in this study.

2.3. Growth measurement

Five wild type (WT) and three sterile mutant (SM) thalli (one thallus per 500 ml conical flask) were grown at 18 ± 1 °C with a 16L: 8D photoperiod and light intensity of $80 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ using a combination of warm and cold fluorescent tubes. The temperature and light conditions used here were previously determined as optimum for *U. rigida* growth (Gao, 2016). The cultures were aerated with a flow rate of 100 L min^{-1} . The seawater was enriched with $500 \mu\text{mol L}^{-1}$ nitrate and $25 \mu\text{mol L}^{-1}$ phosphate and was renewed every day to avoid nutrient limitation. The initial fresh weight for each thallus was $\sim 0.3 \text{ g}$, equal to a stocking density of 0.6 g L^{-1} . The lower quantity of the SM thalli was due to the limited availability of the sterile material. Fresh masses of WT and SM thalli were recorded every three days for 18 or 27 days. The 27-day culture only applied to the SM strain as the WT thalli were lost after day 18 due to reproduction events. The SM was transferred to a 13 L tank on day 9 and then reduced to the stocking density of 1.8 g L^{-1} on day 18 to avoid/reduce self-shading. The stocking density for the WT strain was not reduced given the lower growth rate. The fresh mass was determined after gently blotting each thallus with tissue paper. Specific growth rates (SGR) were calculated using the formula: $\text{SGR (\%)} = [\ln (M_2/M_1)]/t \times 100$, where M_2 is the final mass, M_1 is the initial mass and t is the number of culture days. The mean SGR over 18 days was based on the initial biomass and the biomass reached by day 18.

2.4. Reproduction assessment

Discs of 7 mm diameter from WT and SM strains were placed in 500 ml flasks (three flasks for each strain, each flask containing 25 discs) and cultured under the same conditions as per the growth trial. This experiment was conducted for 27 days.

Reproductive discs were recognized by their colour change; formation of reproductive cells in *Ulva* is accompanied by a change in colour from green (vegetative state) to yellowish (reproductive state) and then to white (after gamete release). This was verified microscopically. Sporulation formation in more than half of the disc area was considered equivalent to complete sporulation. Reproduction rate was expressed as the ratio of reproductive discs to all discs in a flask.

2.5. Nutrient uptake rate measurement

Fresh thalli of the WT and SM strains were cultivated in 13 L tanks with a stocking density of 0.5 g L⁻¹ for 12 days. The culture conditions were the same as per the growth experiment. The thalli were cultured in seawater enriched with 500 µmol N and 25 µmol P for three days prior to the experiment. The change in nutrient levels in the seawater was monitored daily and the daily consumed quantities of nitrate and phosphate were replenished after the measurements. The seawater was renewed every three days. Nitrate and phosphate uptake rates by thalli on days 3, 6, 9, and 12 were estimated from the decrease of NO₃⁻ and PO₄³⁻ in the culture medium over a given time interval (24 hours). Only the data on day 3 were presented as both strains were fully vegetative by day 3. Nitrate was measured by a rapid spectrophotometer method (Collos et al., 1999) and phosphate by the phosphomolybdenum blue colorimetry method (Murphy & Riley, 1962).

2.6. Dry mass

The thalli in the cultures of section 2.5 were harvested at the end of the experiment, and then oven dried at 50 °C until consistent mass (for 24 hours). As shown in Fig. 1, the WT growth rate decreased culture time and was close to zero by day 12, suggesting that biomass would not increase after day 12. To obtain a maximum biomass for biochemical composition analysis, we harvested the thalli on day 12. The tissues were ground to a powder, sieved and placed into tubes within a desiccator pending analyses of chemical composition and functional properties.

2.7. Biochemical composition

Triplicate samples were run for all biochemical and functional analyses for

both strains.

2.7.1. Protein content

Total protein was determined by the Kjeldahl method. The crude protein was calculated using nitrogen content multiplied by 5.45 based on the mean value of three *Ulva* species (Shuuluka et al., 2013), and total nitrogen was measured using an Elementar Vario Macro Cube (Elementar, Hanau Germany). Data were expressed as percentage of dry mass.

2.7.2. Amino acids analysis

Amino acid composition was determined by high-performance liquid chromatography (Ultimate 3000, USA)-tandem mass spectrometry (API 3200 Q-TRAP, USA) (HPLC-MS/MS) (Shimbo et al., 2010). *Ulva* powder (100 mg) was transferred to an ampoule and hydrolysed with 4 ml of 6 N hydrochloric acid at 110 °C for 24 hours. Fifty microliters of this solution was dried under nitrogen and then re-dissolved in 200 µl of deionized water. Eighty microliters of borate buffer (0.1 M, pH 8.8) was added to 40 µl of sample solution and mixed. The solution was derivatised with 40 µl of diethyl ethoxymethylene-malonate at 55 °C for 15 minutes. Samples were transferred to insert vials and 20 µl were injected into the HPLC. The HPLC system consisted of a SRD-3600 Solvent Rack with analytical 6-channel vacuum degasser, a DGP-3600A pump, WPS-3000TSL analytical autosampler, and a tcc-3200 column compartment. Chromatographic separations were performed on an MSLab HP-C18 column (150 × 4.6 mm, 5 µm). The mobile phase consisted of water (A) and acetonitrile (B). The solvent was delivered to the column at a flow rate of 0.8 ml minute⁻¹ as follows: 0–0.1 minute from A-B (95:5) to A-B (95:5); 1–8 minutes from A-B (95:5) to A-B (40:60); 8–8.1 minutes from A-B (40:60) to A-B (0:100); 8.1–10 minutes from A-B (0:100) to A-B (0:100); 10–10.1 minutes from A-B (0:100) to A-B (95:5); 10.1–15 minutes from A-B (95:5) to A-B (95:5).

The conditions for MS-MS detection were optimized to obtain the highest signal intensity and were as follows: mode: positive-ion mode; ion spray voltage:

5,500 V; nebulizer gas pressure: 55 psi; curtain gas pressure: 20 psi; collision gas pressure: medium; turbo gas temperature: 500 °C; entrance potential: 10 V; collision cell exit potential: 2 V. Nitrogen gas was used as the collision gas in the multiple reaction monitoring (SRM) mode. The data were obtained using Analyst software version 1.5.1 (Applied Biosystems). Amino acid standard solution (Aladdin, China) was injected for the system's calibration and amino acid quantification.

2.7.3. Lipid extraction

Lipid was extracted by a modified Folch method (Folch et al., 1957). Homogenized samples (3 g) were extracted with 60 ml chloroform: methanol (2:1) solution. After vortexing for 20 minutes at room temperature, 12 ml of NaCl (0.88%) was added to aid stratification. Samples were centrifuged for five minutes at 1000 g and the upper phase removed. Sixty milliliters of methanol: water (1:1) was used to rinse the tubes. After centrifuging and removal of the upper phase once more, the bottom phase was dried under a steady nitrogen stream. When the chloroform was completely evaporated, the crude lipid was weighed. Results were expressed as percentage of dry mass.

2.7.4. Fatty acid analysis

Fatty acid methyl esters (FAME) were prepared based on extracted lipid from section 2.7.3 according to Joseph and Ackman (1992). A mixture of 100 mg of fat matter and 1.5 ml 0.5 N methanolic NaOH (2 g NaOH in 100 ml methanol) was heated at 100 °C for five minutes. After reaching room temperature, 2 ml of boron trifluoride 12% in methanol was added to the solution, which was then heated at 100 °C for 30 minutes. After cooling for five minutes at room temperature, 1 ml of isooctane and 5 ml of saturated NaCl solution were added. The solution was centrifuged for five minutes at 859 g and the upper layer was transferred to a new tube. One milliliter of isooctane was added to the first solution and the procedure was repeated. The two layers of isooctane were dried under a steady stream of nitrogen and the residue was re-dissolved in 1 ml of hexane. FAMEs were analysed using a gas chromatograph (Shimadzu, GC-2014, Kyoto, Japan) fitted with an Agilent CP-SIL 88

fused silica capillary column (100 m × 0.25 mm ID × 0.2 µm film thickness). Purified helium was used as a carrier gas with a head pressure of 210 kPa and a column flow of 1 ml minute⁻¹. A split injection system was used with an auto injector (Shimadzu, AOC-20i) with a split ratio of 50.0 and an injector temperature of 255 °C. FAME peaks were detected by flame ionisation detection at 260 °C. Samples of 1 µl were injected at an initial column temperature of 70 °C, held for one minute. The temperature was then raised at 5 °C minute⁻¹ to 100 °C, held for two minutes, and then increased at 10 °C minute⁻¹ to 160 °C, and held for 61 minutes. Finally, the temperature was increased to 240 °C at a rate of 5 °C minute⁻¹, and held for 21 minutes, thus giving a final gradient of 113 minutes total runtime. Peaks were identified by comparing the chromatograms to a commercial 52 FAME standard (GLC463, Nu-Chek Prep Inc., Elysian, MN, USA). Quantification was based on peak areas of individual identified and non-identified FAME, expressed as a percentage of the total peak areas for quantified FAME.

2.7.5. *Ash content*

Ash was determined by incinerating samples in a muffle furnace at 550 °C for 24 hours. The total ash content was expressed as percentage of dry mass.

2.7.6. *Carbohydrate*

Carbohydrate was estimated by subtracting the total protein, lipid, and ash content from the total dry mass (Ortiz et al., 2006).

2.8. *Functional properties*

2.8.1. *Swelling capacity*

The swelling capacity (SWC) was calculated using a slightly modified method of bed volume after equilibrating in excess solvent (Yaich et al., 2011). Triplicate samples of *Ulva* powder (0.2 g) were placed into 10 ml measuring cylinders. Ten milliliters of distilled water was added, immediately vigorously mixed and incubated for 18 hours at 37 °C. The swelling volume was recorded and SWC was expressed as ml of swollen sample per gram of sample.

2.8.2. Water holding capacity

The water holding capacity (WHC) was analysed by the centrifugation method (Yaich et al., 2011). Triplicate samples of *Ulva* powder (0.2 g) were placed into pre-weighed centrifuge tubes. Five milliliters of distilled water was added and mixed vigorously. The dispersion was centrifuged for 25 minutes at 3000 g after incubating at 37 °C for one hour. The supernatant was removed and the residue was oven dried at 50 °C for 25 minutes. The WHC was expressed as grams of water bound per gram of sample.

2.8.3. Oil holding capacity

The oil holding capacity (OHC) was measured according to Yaich et al. (2011). Triplicate samples of *Ulva* powder (0.5 g) were dispersed in 5 ml of food grade corn oil in centrifuge tubes. The mixtures were stirred and incubated at 37 °C for one hour, followed by centrifugation at 3000 g for 25 minutes. The oil supernatant was transferred to a 10 ml measuring cylinder and measured. The OHC was expressed as grams of oil held by one gram of sample (dry weight). The density of the corn oil was 0.92 g ml⁻¹.

2.9. Statistical analysis

Results were expressed as means of replicates $\pm$ standard deviation. All data were analysed using the software SPSS v.21.

The SGRs of both the wild and mutant strains at each culture time were normally distributed (Shapiro-Wilk, $P > 0.05$) and had equal variances (wild: Levene's test, $F = 0.743$, $P > 0.05$; mutant: Levene's test, $F = 2.257$, $P > 0.05$). The effect of culture time on SGR was assessed using repeated measures ANOVA with Least-Significant Difference (LSD) for *post hoc* investigation. The mean SGR of each strain over the first 18-day culture was also normally distributed with equal variances (Levene's test, $F = 2.208$, $P > 0.05$); therefore a t-test was conducted.

The reproduction rate of the WT strain at all culture times was also normally distributed with equal variances (Levene's test, $F = 3.300$, $P > 0.05$); a repeated

ANOVA with LSD was conducted.

Data on nutrient uptake, protein, lipid, ash, carbohydrate, and functional properties of both strains were normally distributed (Shapiro-Wilk, $P > 0.05$) with equal variances (Levene's test, $F < 6.750$, $P > 0.05$); t-tests were conducted. A confidence interval of 95% was set for all tests.

All amino acids in the WT strain and total amino acids in the SM strain were normally distributed apart from the total essential amino acids (Shapiro-Wilk, $P = 0.010$). The variances of the two samples for all amino acids could be considered equal (Levene's test, $F < 6.750$, $P > 0.05$) except for arginine (Levene's test, $F = 11.144$, $P = 0.029$) and aspartic acid (Levene's test, $F = 11.046$, $P = 0.029$). A one-way multivariate ANOVA (MANOVA) was used to compare the differences in amino acid content as MANOVA is reasonably robust to violations of normal distribution and homogeneity of variance, particularly when the group sizes are equal (Pallant, 2010). Bonferroni adjustment was used to reduce the chance of Type I error. A new alpha level of 0.0024 (0.05 divided by 21) was set since there were 21 dependent variables. All fatty acids in the WT strain and polyunsaturated fatty acids (PUFA) and n6-fatty acids in the SM strain were normally distributed except for C18:2n6 (Shapiro-Wilk, $P = 0.011$) and C18:2n3 (Shapiro-Wilk, $P = 0.040$). The variances for all fatty acids could be considered equal (Levene's test, $F < 6.750$, $P > 0.05$) except for C20 (Levene's test, $F = 13.398$, $P = 0.022$) and C20:4n6 (Levene's test, $F = 13.408$, $P = 0.022$). A one-way MANOVA was used to compare the differences in fatty acid content between the two strains; a Bonferroni adjusted alpha level of 0.0014 (0.05 divided by 37) was set since there were 37 dependent variables.

3. Results and Discussion

3.1. Growth and reproduction

The WT SGR differed significantly over culture times (Fig. 1; $F = 197.212$, $df = 5, 20$, $P < 0.001$), reaching $22.8 \pm 1.7\%$ during the first three days but declining to

almost zero by day 12 (LSD, $P < 0.001$; Fig. 1A) and subsequently losing biomass with the SGR dropping to $-15.6 \pm 2.4\%$ during the last three days (Fig. 1A). The SM SGR also differed significantly over culture times ($F = 9.872$, $df = 8, 16$, $P < 0.001$). In contrast to the WT, the SM maintained a positive SGR over the course of the 27-day culture, albeit at a reduced rate with time (LSD, $P < 0.01$), with the SGR on the last three days still relatively high ($11.4 \pm 2.3\%$; Fig. 1B). There were significant differences in mean SGR between the WT and SM strains over the first 18 days (Fig. 1C), with the SM mean SGR five times faster than the WT (Independent samples t-test, $t = -48.173$, $df = 6$, $P < 0.001$).

The WT growth decline with time was consistent with other published *Ulva* research (Ale et al., 2011). The decrease in growth may be due to a switch to reproduction, which effectively terminates vegetative growth and leads to a pronounced disintegration of part, or all of the thallus (Bruhn et al., 2011; Gao et al. 2017). The WT reproduction rate differed significantly over culture times ($F = 545.701$, $df = 8, 16$, $P < 0.001$). There was no reproductive activity during the first six days whereas $18.7 \pm 2.3\%$ of discs became fertile by day nine (Fig. 2). The reproduction rate increased with time (LSD, $P < 0.01$) reaching 100% by day 21. In contrast, all SM discs remained in the vegetative state throughout the 27-day trial. As the SM discs did not experience reproductive biomass loss, the resulting fronds grew to more than 50 cm in length (Fig. 3). In addition, the decline of growth in both strains with culture time could be due to decreased light availability due to self-shading. The biomass was only harvested at the end of the experiment with a possibility of increased self-shading and thus decreased light availability. A positive relationship between light and biomass productivity in *U. ohnoi* has been demonstrated (Mata et al., 2016).

High and sustained growth rates are desired in aquaculture. The SM mean SGR over 18 days was $20.0 \pm 0.6\%$ – five times faster than the WT. The mean SGR over the 27-day trial was $17.4 \pm 0.3\%$, exceeding values reported by Gordillo et al. (2001) ($1\text{--}6\%$ in a 10-day lab culture of *U. rigida*) and Viaroli et al. (1996) ($2.5\text{--}8.1\%$

in a two-week outdoor culture of *U. rigida*).

3.2. Nutrient uptake

Apart from the beneficial effects of sterility, this quick growth (Fig. 1&S1) could also be related to an enhanced capacity for nutrient uptake, which if true, would indicate that SM could be an effective strain for nutrient capture. This assumption was supported by the nutrient uptake data. Nitrate uptake by the SM was $171.6 \pm 12.1 \mu\text{mol g FW}^{-1} \text{ day}^{-1}$, which was 40.0% higher than the WT (Fig. 4A) (Independent samples t-test, $t = -5.337$, $df = 4$, $P = 0.006$). The phosphorus uptake of the SM was $5.7 \pm 0.4 \mu\text{mol g FW}^{-1} \text{ day}^{-1}$, significantly higher (30.9%) than the WT ($4.4 \pm 0.5 \mu\text{mol g FW}^{-1} \text{ day}^{-1}$; Independent samples t-test, $t = -3.506$, $df = 4$, $P = 0.025$, Fig. 4B).

Nitrogen and phosphorus are essential nutrients that support growth, with rapidly proliferating cells having a large nutrient demand. The nitrate uptake capacity of the SM was greater when compared with other *Ulva* species, e.g. 133.7 and 109.7 $\mu\text{mol g FW}^{-1} \text{ day}^{-1}$ in *U. fenestrata* and *U. intestinalis* respectively (Björnsäter & Wheeler, 1990), and $\sim 70 \mu\text{mol g DW}^{-1} \text{ day}^{-1}$ in *U. lactuca* (Ale et al., 2011), making the SM a potentially efficient biofilter for wastewater bioremediation, as a biofilter to combat eutrophication, or as an extractive component within an integrated multi-trophic aquaculture system. The phosphate uptake capacity of the SM exceeded the 0.637 ± 0.266 and $0.15 \mu\text{mol g FW}^{-1} \text{ day}^{-1}$ of *U. rotundata* (Hernández et al., 2005) and *U. pertusa* (Seob et al., 2010) respectively, the $1.4\text{--}1.6 \mu\text{mol g}^{-1} \text{ FW day}^{-1}$ (based on the ratio of FW:DW = 5:1) for *U. intestinalis* (Martínez-Aragón et al., 2002), and was comparable to the $1.01\text{--}5.04$ and $3.42\text{--}6.54 \mu\text{mol g FW}^{-1} \text{ day}^{-1}$ for *U. fenestrata* and *U. intestinalis* respectively (Björnsäter & Wheeler, 1990).

The N: P uptake ratio (29.9) in the SM was higher than the 3.7–4.4 for *U. linza*, 4.5–5.1 for *U. prolifera* (Luo et al., 2012), 9.4 for *U. rotundata* (Hernández et al., 2005), and 16.6–16.8 for *U. intestinalis* (Björnsäter & Wheeler, 1990), and was

comparable to the 11.8–36.0 for *U. fenestrata* (Björnsäter & Wheeler, 1990).

3.3. Chemical composition and functional properties

The protein content of the WT accounted for $24.1 \pm 0.8\%$ of the biomass, which was 58.8% higher than the SM (Figure 5; Independent samples t-test, $t = 8.224$, $df = 4$, $P = 0.001$). Although the SM had a faster nitrate uptake rate, the higher growth rate (Fig. S1) could dilute the assimilated nitrogen, resulting in a lower protein content. In addition, nitrate limitation occurred in the SM but not in the WT towards the end of the culture (Fig. S2), which could also contribute to the lower protein content. Løvlie (1969) also reported a lower protein content in a fast-growing mutant of *U. mutabilis*. In contrast, the total lipid content of the SM was more than double that of the WT with the latter only $5.4 \pm 0.4\%$ of dry mass (Independent samples t-test, $t = -19.052$, $df = 4$, $P < 0.001$). The SM ash content ($34.0 \pm 1.1\%$) was also higher than the WT ($31.4 \pm 1.2\%$) (Independent samples t-test, $t = -2.812$, $df = 4$, $P = 0.048$). There were no significant differences between strains for carbohydrate content.

The 13.8% lipid content of the SM was high compared with the WT and higher still when compared with other *Ulva* species; 2.5–3.5% in *U. clathrata* (Peña-Rodríguez et al., 2011) and 7.9% in *U. lactuca* (Yaich et al., 2011). The most abundant fatty acid in both strains was C16:0 (Table 1). There was also a high proportion of C18 unsaturated fatty acids. The SM had significantly more C12:0, C14:0, C17:1 (cis-9), C18:0, C18:2n6, C18:3n3, C18:3n6, C20:3n6, C20:4n6, C22:5n3 and C22:4n6 fatty acids than the WT, which translated to significantly more monounsaturated fatty acids (MUFA), polyunsaturated fatty acids (PUFA), omega-3 fatty acids (n3-FA), and omega-6 fatty acids (n6-FA), which could have greater benefits for health and nutrition applications (Koletzko et al., 2004; Ruxton et al., 2014). The ratio of n6 to n3-FA was higher in the SM but the value was still below the maximum World Health Organization recommended value of 10, indicating that the SM strain is an efficient food resource to reduce n6-FA/n3-FA.

Among the amino acids, glutamic acid and alanine were the most abundant in both strains. Aspartic acid and arginine were the second highest group for the WT while valine and leucine were the next highest for the SM. There was more histidine, serine, glutamic acid, isoleucine, total amino acids, and non-essential amino acids (NEAA) in the WT compared to the SM strain (Table 2). Angell et al. (2014) defined three internal nitrogen states in *U. ohnoi*: nitrogen-limited (<1.2% N), metabolic (1.2-2.6% N), and luxury (>2.6%). All amino acids and growth increase with rising nitrogen content in the nitrogen-limited state. Growth rate does not change and all amino acids still increase with rising nitrogen content in the metabolic state. The luxury state was in which growth rate decreases and only specific amino acids (glutamic acid, glutamine, and arginine) have a large increase. Meanwhile, the nitrogen-limited content for *U. intestinalis* and *U. fenestrata* were 2.5% and 3.2% respectively (Björnsäter and Wheeler 1990), suggesting species specific differences. In the present study, the internal nitrogen content in the WT and SM strains were $4.34 \pm 0.14\%$ and $2.79 \pm 0.32\%$. Based on the nutrient consumption (Fig. S2) and growth data (Fig. S1), the SM experienced nitrogen limitation while the WT may have been in the state of luxury. The high content of glutamic acid in the WT would also support its nitrogen state (Angell et al., 2014).

Although there was less protein in the SM, there was a relatively high ratio of essential amino acids (EAA) to NEAA (1.06 ± 0.06), making the SM preferable in terms of EAA sources as the EAA: NEAA ratio of other *Ulva* species is usually below 1 (Taboada et al., 2010; Peña-Rodríguez et al., 2011).

The faster growing SM will have had a higher demand for phospholipid membrane synthesis due to cell division. Cell membranes have a high PUFA content to maintain membrane fluidity (Yang et al., 2011). This would explain why the SM had both a higher overall lipid content (relative to protein) and an increased concentration of PUFA, as well as the higher metabolic demand for phosphorous. In addition, more energy will have been needed by the SM to maintain the higher rate of cell division, requiring a higher rate of transformation of fixed carbon into biomass.

However, there was no statistical difference in carbohydrate content between strains, a crude proxy for photosynthetic performance. The reason could be that more of the ATP and NADPH produced during photosynthesis were used to synthesize lipid rather than carbohydrate and protein.

The rapid vegetative growth of the SM may have contributed to inadvertent nitrogen limitation towards the end of the culture trial (Fig. S2). When nitrogen is limited, the fixed carbons are channeled towards the synthesis of lipid rather than protein, as there are no chemical building blocks for making more (nitrogen containing) protein (Roessler, 1990; Floreto et al., 1996; Malta and de Nys, 2015).

The swelling, and water- and oil-holding capacities of the SM were all significantly lower than the WT (SWC = 27.7% lower, Independent samples t-test, $t(4) = 30.946$, $P < 0.001$; WHC = 25.3% lower, $t = 15.399$, $df = 4$, $P < 0.001$; OHC = 11.5% lower; $t = 4.461$, $df = 4$, $P = 0.011$; Fig. 6A). Intake of food with a higher swelling capacity can enhance satiety and thereby lead to a reduced calorie intake (Dettmar et al., 2011). Foods with a high SWC therefore have applications in adjunctive therapies for the treatment of obesity (Dettmar et al., 2011). The WT SWC (11.2 ± 0.2 ml g⁻¹ DW) was slightly lower than that of *U. lactuca* (13.0 ± 0.70 ml g⁻¹ DW; Wong and Cheung, 2000), but higher than that reported elsewhere for *U. lactuca* (0.3 ml g⁻¹ DW; Yaich et al., 2011) and also higher than commercial fibres from oat, wheat, apple, and bamboo (4.98–7.60 ml g⁻¹ DW; Rosell et al., 2009). The SWC is mainly influenced by the protein and fibre content as well as the particle size of fibre. In addition, the conditions of measurement, such as pH, temperature, and ionic strength and types of ions in solution, can also affect the SWC (Fleury & Lahaye, 1991; Yaich et al., 2011). The water-holding capacity can modify the viscosity and texture of formulated food and a higher WHC can assist in avoiding syneresis. The WT WHC (6.98 ± 0.17 g g⁻¹ DW) was lower than that reported by Wong and Cheung (2000) (9.71 ± 0.11 g g⁻¹ DW), comparable to Yaich et al. (2011) (6.66 to 7.00 g g⁻¹ DW), and higher than commercial fibres from oat, wheat, apple, and bamboo (3.69–6.89 g g⁻¹ DW; Rosell et al., 2009). According to Robertson and Eastwood (1981),

water can be associated with fibre either as bound or trapped water. The amount of bound water depends on chemical composition whereas trapped water refers to the fibre structure. The centrifugation method used here incorporates both fibre-water states. Protein conformations and the nature of the water binding sites in the protein molecules also play a role in water holding (Chou & Morr, 1979). The total protein and carbohydrate content of the SM ($52.1 \pm 1.7\%$) was 17.5% lower than the WT (Figure 6B), contributing to the SM's lower SWC and WHC. The WT ($1.75 \pm 0.07 \text{ g g}^{-1} \text{ DW}$) and SM OHCs ($1.54 \pm 0.04 \text{ g g}^{-1} \text{ DW}$) were higher than reported by Wong and Cheung (2000) ($0.65 \pm 0.03 \text{ g g}^{-1} \text{ DW}$). This may be due to different treatment temperatures. The OHC in the present study was determined at 37 °C, whereas Wong and Cheung (2000) used 25 °C. Our results were comparable to Yaich et al. (2011) (around $1.60 \text{ g g}^{-1} \text{ DW}$ at 40 °C). In general, proteins play a major role in fat absorption by hydrophobic bonding. This may result in a higher OHC in WT that has a higher protein content compared to the SM. Furthermore, the OHC of seaweeds is also related to particle size, overall charge density and the hydrophilic nature of fibre (Kinsella & Melachouris, 1976; Voutsinas & Nakai, 1983). Due to their OHC, *Ulva* species can be used as stabilizers in formulated food products. Moreover, the OHC of both *Ulva* strains in the present study was significantly higher than commercial cellulose ($0.98 \pm 0.09 \text{ g g}^{-1} \text{ DW}$; Hassan et al., 2011), making them useful in reducing blood lipid level, obesity and coronary heart disease risk.

4. Conclusion

The results clearly demonstrate that this new sterile strain is a positive step towards increasing the long-term productivity of *Ulva* by preventing reproductive events. Thus, this provides a platform to serve multiple market applications. However, the caveat should be added that using clonal strains does increase the risks of disease outbreaks. The rapid and sustained growth rate combined with high rates of nitrate and phosphate uptake suggest a strong carbon capture and wastewater bioremediation capacity. In addition, the enhanced MUFA, PUFA, and omega-3 fatty acids offer

benefits for health food applications. More targeted molecular studies combined with further genetic optimization will likely deliver benefits beyond that achieved by physical mutagenesis; however, the complexities of modifying eukaryotic genomes prevent these approaches from making an immediate impact.

Acknowledgements

This research was funded by a School of Marine Science and Technology doctoral scholarship awarded to GG. We thank David Whitaker, Eleni Charzidimitriou, and Xude Zhang for their technical assistance over the duration of this project. CR is Director of Seaweed & Co. Ltd which advises on the supply and quality control of seaweeds into the UK marketplace. Finally, we recognize the detailed and constructive comments from four anonymous reviewers that have greatly improved the clarity of our work.

References

- Ahmed, E.-W., Ahmed, A.-H., Aly, O., Emad Samara, K.A. 2015. Effect of dietary seaweed (*Ulva lactuca*) supplementation on growth performance of sheep and on in vitro gas production kinetics. Turk. J. Vet. Anim. Sci. 39, 81-86.
- Al-Hafedh, Y.S., Alam, A., Buschmann, A.H. 2015. Bioremediation potential, growth and biomass yield of the green seaweed, *Ulva lactuca* in an integrated marine aquaculture system at the Red Sea coast of Saudi Arabia at different stocking densities and effluent flow rates. Rev. Aquacult. 7, 161-171.
- Ale, M.T., Mikkelsen, J.D., Meyer, A.S. 2011. Differential growth response of *Ulva lactuca* to ammonium and nitrate assimilation. J. Appl. Phycol. 23, 345-351.
- Björnsäter, B.R., Wheeler, P.A. 1990. Effect of nitrogen and phosphorus supply on growth and tissue composition of *Ulva fenestrata* and *Enteromorpha intestinalis* (Ulvales, Chlorophyta). J. Phycol. 26, 603-611.

487 Bolton, J.J., Robertson-Andersson, D.V., Shuuluka, D., Kandjengo, L. 2009. Growing
 488 *Ulva* (Chlorophyta) in integrated systems as a commercial crop for abalone
 489 feed in South Africa: a SWOT analysis. J. Appl. Phycol. 21, 575-583.

490 Bruhn, A., Dahl, J., Nielsen, H.B., Nikolaisen, L., Rasmussen, M.B., Markager, S.,
 491 Olesen, B., Arias, C., Jensen, P.D. 2011. Bioenergy potential of *Ulva lactuca*:
 492 Biomass yield, methane production and combustion. Bioresource Technol.
 493 102, 2595-2604.

494 Carl, C., Lawton, R.J., Paul, N.A., De Nys, R. 2016. Reproductive output and
 495 productivity of filamentous tropical *Ulva* over time. J. Appl. Phycol. 28,
 496 429-438.

497 Castelar, B., Reis, R.P., dos Santos Calheiros, A.C. 2014. *Ulva lactuca* and *U.*
 498 *flexuosa* (Chlorophyta, Ulvophyceae) cultivation in Brazilian tropical waters:
 499 recruitment, growth, and ulvan yield. J. Appl. Phycol. 26, 1989-1999.

500 Charrier, B., Rolland, E., Gupta, V., Reddy, C.R.K. 2015. Production of genetically
 501 and developmentally modified seaweeds: exploiting the potential of artificial
 502 selection techniques. Front. Plant Sci. 6, 127.

503 Chou, D.H., Morr, C.V. 1979. Protein-water interactions and functional properties. J.
 504 Am. Oil Chem. Soc. 56, A53-A62.

505 Chung, I.K., Beardall, J., Mehta, S., Sahoo, D., Stojkovic, S. 2011. Using marine
 506 macroalgae for carbon sequestration: a critical appraisal. J. Appl. Phycol. 23,
 507 877-886.

508 Collos, Y., Mornet, F., Sciandra, A., Waser, N., Larson, A., Harrison, P.J. 1999. An
 509 optical method for the rapid measurement of micromolar concentrations of
 510 nitrate in marine phytoplankton cultures. J. Appl. Phycol., 11, 179-184.

511 Cyrus, M.D., Bolton, J.J., Scholtz, R., Macey, B.M. 2015. The advantages of *Ulva*
 512 (Chlorophyta) as an additive in sea urchin formulated feeds: effects on
 513 palatability, consumption and digestibility. Aquacult. Nutr. 21, 578-591.

514 DeBusk, T., Blakeslee, M., Ryther, J. 1986. Studies on the outdoor cultivation of *Ulva*
 515 *lactuca* L. Bot. Mar. 29, 381-386.

516 Dettmar, P. W., Strugala, V., Richardson, J. C. 2011. The key role alginates play in
517 health. *Food Hydrocolloid.*, 25, 263–266.

518 Doty, M.S., Alvarez, V.B. 1975. Status, problem, advances and economics of
519 *Eucheuma* farms. *Mar. Technol. Soc. J.* 9, 30-35.

520 Fleury, N., Lahaye, M. 1991. Chemical and physico-chemical characterisation of
521 fibres from *Laminaria digitata* (kombu breton): A physiological approach. *J.*
522 *Sci. Food Agr.* 55, 389-400.

523 Floreto, E.A.T., Teshima, S., Ishikawa, M. 1996. Effects of nitrogen and phosphorus
524 on the growth and fatty acid composition of *Ulva pertusa* kjellman
525 (Chlorophyta). *Bot. Mar.* 39, 69-74.

526 Folch, J., Lees, M., Sloane-Stanley, G.H. 1957. A simple method for the isolation and
527 purification of total lipids from animal tissues. *J. Biol. Chem.* 226, 497-509.

528 Fu, F., Sui, Z.H., Zhou, W., Wang, J.G., Chang, L.P., Ci, S.F. 2014. UV-irradiation
529 mutation of tetraspores of *Gracilariopsis lemaneiformis* and screening of
530 thermotolerant strains. *J. Appl. Phycol.* 26, 647-656.

531 Gao, G. 2016. Developing Systems for the Commercial Culture of *Ulva* species in the
532 UK. Ph. D. thesis, Newcastle University, UK.

533 Gao, G., Clare, A.S., Rose, C., Caldwell, G.S. 2017. Eutrophication and
534 warming-driven green tides (*Ulva rigida*) are predicted to increase under
535 future climate change scenarios. *Mar. Pollut. Bull.* 114, 439-447.

536 Hassan, F.A., Ismail, A., Hamid, A.A., Azlana, A., Al-sheraji S.H. 2011.
537 Characterisation of fibre-rich powder and antioxidant capacity of *Mangifera*
538 *pajang* K. fruit peels. *Food Chem.* 126, 283-288.

539 Hernández, I., Fernández-Engo, M.A., Pérez-Lloréns, J.L., Vergara, J.J. 2005.
540 Integrated outdoor culture of two estuarine macroalgae as biofilters for
541 dissolved nutrients from *Sparus auratus* waste waters. *J. Appl. Phycol.* 17,
542 557-567.

543 Hernández, I., Martínez-Aragón, J.F., Tovar, A., Pérez-Lloréns, J.L., Vergara, J.J.
544 2002. Biofiltering efficiency in removal of dissolved nutrients by three species

545 of estuarine macroalgae cultivated with sea bass (*Dicentrarchus labrax*) waste
 546 waters 2. Ammonium. J. Appl. Phycol. 14, 375-384.
 547 Hiraoka, M., Enomoto, S., 1998. The induction of reproductive cell formation of *Ulva*
 548 *pertusa* Kjellman (Ulvales, Ulvophyceae). Phycol. Res. 46, 199-203.
 549 Hiraoka, M., Oka, N. 2008. Tank cultivation of *Ulva prolifera* in deep seawater using
 550 a new “germling cluster” method. J. Appl. Phycol. 20, 97-102.
 551 Joseph, J.D., Ackman, R.G. 1992. Capillary column gas chromatographic method for
 552 analysis of encapsulated fish oils and fish oil ethyl esters: collaborative study.
 553 J. AOAC Int. 75, 488-506.
 554 Kakinuma, M., Coury, D., Kuno, Y., Itoh, S., Kozawa, Y., Inagaki, E., Yoshiura, Y.,
 555 Amano, H. 2006. Physiological and biochemical responses to thermal and
 556 salinity stresses in a sterile mutant of *Ulva pertusa* (Ulvales, Chlorophyta).
 557 Mar. Biol. 149, 97-106.
 558 Kinsella, J.E., Melachouris, N. 1976. Functional properties of proteins in foods: a
 559 survey. Crit. Rev. Food Sci. Nutr. 7, 219-280.
 560 Kirkendale, L., Saunders, G.W., Winberg, P. 2013. A molecular survey of *Ulva*
 561 (Chlorophyta) in temperate Australia reveals enhanced levels of
 562 cosmopolitanism. J. Phycol. 49, 69-81.
 563 Koletzko, B., Boey, C.C.M., Campoy, C., Carlson, S.E., Chang, N.,
 564 Guillermo-Tuazon, M.A., Joshi, S., Prell, C., Quak, S.H., Sjarif, D.R. 2014.
 565 Current information and Asian perspectives on long-chain polyunsaturated
 566 fatty acids in pregnancy, lactation, and infancy: systematic review and practice
 567 recommendations from an early nutrition academy workshop. Ann. Nutr.
 568 Metab. 65, 49-80.
 569 Lawton R J, Mata L, de Nys R, Paul N A. Algal bioremediation of waste waters from
 570 land-based aquaculture using *Ulva*: selecting target species and strains. PLoS
 571 ONE, 2013, 8(10): e77344.
 572 Løvlie, A. 1969. Cell size, nucleic acids, and synthetic efficiency in the wild type and
 573 a growth mutant of the multicellular alga *Ulva mutabilis* Føyn. Dev. Biol. 20,
 574 349-367.

575 Luo, M.B., Liu, F. 2011. Salinity-induced oxidative stress and regulation of
 576 antioxidant defense system in the marine macroalga *Ulva prolifera*. J. Exp.
 577 Mar. Biol. Ecol. 409, 223-228.

578 Luo, M.B., Liu, F., Xu, Z.L. 2012. Growth and nutrient uptake capacity of two
 579 co-occurring species, *Ulva prolifera* and *Ulva linza*. Aquat. Bot. 100, 18-24.

580 Malta, E., de Nys, R. 2016. The effect of short-term preharvest strategies on the
 581 carbon constituents of *Ulva ohnoi* M. Hiraoka et S. Shimada. J. Appl. Phycol.
 582 28, 555-565 Markou, G., Angelidaki, I., Georgakakis, D. 2012. Microalgal
 583 carbohydrates: an overview of the factors influencing carbohydrates
 584 production, and of main bioconversion technologies for production of biofuels.
 585 Appl. Microbiol. Biotechnol. 96, 631-645.

586 Martínez-Aragón, J. F., Hernández, I., Pérez-Lloréns, J. L., Vázquez, R., Vergara, J. J.
 587 2002. Biofiltering efficiency in removal of dissolved nutrients by three species
 588 of estuarine macroalgae cultivated with sea bass (*Dicentrarchus labrax*) waste
 589 waters 1. Phosphate. J. Appl. Phycol. 14, 365–374.

590 Mata L, Magnusson M, Paul N A, de Nys R. The intensive land-based production of
 591 the green seaweeds *Derbesia tenuissima* and *Ulva ohnoi*: biomass and
 592 bioproducts. J. Appl. Phycol. 2016, 28: 365-375.

593 Migita, S. 1985. The sterile mutant of *Ulva pertusa* Kjellman from Oumra Bay. Bull.
 594 Fac. Fisheries Nagasaki Univ. 57, 33-37.

595 Murase, N., Maegawa, M., Matsui, T., Ohgai, M., Katayama, N., Saitoh, M.,
 596 Yokohama, Y. 1994. Growth and photosynthesis temperature characteristics
 597 of the sterile *Ulva pertusa*. Nippon Suisan Gakk. 60, 625-630.

598 Murphy, J., Riley, J.P. 1962. A modified single solution method for the determination
 599 of phosphate in natural waters. Anal. Chim. Acta 27, 31-36.

600 Niwa, K., Yamamoto, T., Furuita, H., Abe, T. 2011. Mutation breeding in the marine
 601 crop *Porphyra yezoensis* (Bangiales, Rhodophyta): cultivation experiment of
 602 the artificial red mutant isolated by heavy-ion beam mutagenesis. Aquaculture
 603 314, 182-187.

604 Ortiz, J., Romero, N., Robert, P., Araya, J., Lopez-Hernandez, J., Bozzo, C.,
 605 Navarrete, E., Osorio, A., Rios, A. 2006. Dietary fiber, amino acid, fatty acid
 606 and tocopherol contents of the edible seaweeds *Ulva lactuca* and *Durvillaea*
 607 *antarctica*. Food Chem. 99, 98-104.

608 Oza, R.M., Sreenivasa Rao, P. 1977. Effect of different culture media on growth and
 609 sporulation of laboratory raised germlings of *Ulva fasciata* Delile. Bot. Mar.
 610 20, 427-432.

611 Pallant, J. 2010. SPSS Survival Manual: A Step by Step Guide to Data Analysis
 612 Using SPSS for Windows Version 10. Open University Press, Buckingham
 613 Philadelphia.

614 Peña-Rodríguez, A., Mawhinney, T.P., Ricque-Marie, D., Cruz-Suárez, L.E. 2011.
 615 Chemical composition of cultivated seaweed *Ulva clathrata* (Roth) C. Agardh.
 616 Food Chem. 129, 491-498.

617 Robertson, J.A., Eastwood, M.A. 1981. An examination of factors which may affect
 618 the water holding capacity of dietary fibre. Brit. J. Nutr. 45, 83-88.

619 Rosell, C.M., Santos, E., Collar, C. 2009. Physico-chemical properties of commercial
 620 fibres from different sources: A comparative approach. Food Res. Int. 42,
 621 176-184.

622 Roessler, P.G. 1990. Environmental control of glycerolipid metabolism in microalgae:
 623 commercial implications and future research directions. J. Phycol. 26,
 624 393-399.

625 Ruxton, C.H.S., Reed, S.C., Simpson, M.J.A., Millington, K.J. 2004. The health
 626 benefits of omega-3 polyunsaturated fatty acids: a review of the evidence. J.
 627 Hum. Nutr. Diet., 17, 449-459.

628 Seob, C.T., Ju, K.E., Young, K.K. 2010. Effect of salinity on growth and nutrient
 629 uptake of *Ulva pertusa* (Chlorophyta) from an eelgrass bed. Algae 25, 17-26.

630 Shimbo, K., Kubo, S., Harada, Y., Oonuki, T., Yokokura, T., Yoshida, H., Amao, M.,
 631 Nokomura, M., Kageyama, N. and Yamazaki, J. 2010. Automated precolumn
 632 derivatization system for analyzing physiological amino acids by liquid

633 chromatography/mass spectrometry. Biomed. Chromatogr. 24, 683-691.

634 Shuuluka, D., Bolton, J.J., Anderson, R.J. 2013. Protein content, amino acid
 635 composition and nitrogen-to-protein conversion factors of and from natural
 636 populations and from an aquaculture system, in South Africa. J. Appl. Phycol.
 637 25, 677-685.

638 Taboada, C., Millán, R., Míguez, I. 2010. Composition, nutritional aspects and effect
 639 on serum parameters of marine algae *Ulva rigida*. J. Sci. Food Agr. 90,
 640 445-449.

641 Van Der Meer, J.P. 1983. The domestication of seaweeds. Bioscience 33, 172-176.

642 Voutsinas, L.P., Nakai, S. 1983. A simple turbidimetric method for determining the
 643 fat binding capacity of proteins. J. Agr. Food Chem. 31, 58-63.

644 Waaland, J.R. 1978. Growth and strain selection in *Gigartina exasperata*
 645 (Florideophyceae). Proceedings of the International Seaweed Symposium, 9,
 646 241-248.

647 Wong, K.H., Cheung, P.C.K. 2000. Nutritional evaluation of some subtropical red and
 648 green seaweeds: Part I—proximate composition, amino acid profiles and some
 649 physico-chemical properties. Food Chem. 71, 475-482.

650 Yaich, H., Garna, H., Besbes, S., Paquot, M., Blecker, C., Attia, H. 2011. Chemical
 651 composition and functional properties of *Ulva lactuca* seaweed collected in
 652 Tunisia. Food Chem. 128, 895-901.

653 Yang X, Sheng W, Sun G Y, Lee J.C.M. 2011. Effects of fatty acid unsaturation
 654 numbers on membrane fluidity and α -secretase-dependent amyloid precursor
 655 protein processing. Neurochem. Int., 58, 321-329.

Figure legends

Fig. 1. The specific growth rates of wild type (WT) and sterile mutant (SM) strains of *Ulva rigida*: A) WT, B) SM, and C) mean specific growth rates of WT and SM over an 18-day cultivation. The error bars indicate the standard deviations ($n = 3$ or 5 , 3 for the sterile strain and 5 for the wild strain). Different superscript letters indicate significant differences over culture periods (A, B) or strains (C).

Fig. 2. Reproduction rates of the wild type (WT) and sterile mutant (SM) strains of *Ulva rigida* over a 27-day cultivation. The error bars indicate the standard deviations ($n = 3$).

Fig. 3. The sterile mutant strain of *Ulva rigida* obtained by mutating a wild type strain. Scale bar = 10 cm.

Fig. 4. Nutrient uptake of the wild type (WT) and sterile mutant (SM) strains of *Ulva rigida* on day 3 over a 12 days of cultivation; A) nitrate, and B) phosphate. The error bars indicate the standard deviations ($n = 3$). Different superscript letters indicate significant differences between strains.

Fig. 5. Bulk chemical composition of the wild type (WT) and sterile mutant (SM) strains of *Ulva rigida*. The error bars indicate the standard deviations ($n = 3$). Different superscript letters indicate significant differences between strains.

Fig. 6 Functional properties (A) and content of protein plus carbohydrate (B) in wild type (WT) and sterile mutant (SM) strains. The error bars indicate the standard deviations ($n = 3$). SWC = swelling capacity, WHC = water holding capacity, and OHC = oil holding capacity. Different superscript letters indicate significant differences between strains.

Table 1. Fatty acid profiles (% of total fatty acids) of wild type (WT) and sterile mutant (SM) strains of *Ulva rigida*. Data are the means $\pm$ standard deviation (n = 3). DF, F, and *P* values were from the MANOVA. Bonferroni adjustment with new alpha level of 0.0014 was used to reduce the chance of Type I error. The fatty acids with significant differences between the strains are in bold. SFA = saturated fatty acids, MUFA = monounsaturated fatty acids, PUFA = polyunsaturated fatty acids. n6-FA = total omega-6 fatty acids, n3-FA = total omega-3 fatty acids.

Fatty acid	Wild Type	Sterile Mutant	DF	F	<i>P</i> value
C10:0	0.02 $\pm$ 0.00	0.02 $\pm$ 0.00	1	1.000	0.374
C12:0	0.04$\pm$0.00	0.01$\pm$0.00	1	519.501	<0.001
C14:0	0.87$\pm$0.03	0.29$\pm$0.03	1	1056.034	<0.001
C14:1 (<i>cis</i> 9)	0.06 $\pm$ 0.05	0.06 $\pm$ 0.02	1	0.045	0.842
C15:0	0.07 $\pm$ 0.00	0.04 $\pm$ 0.00	1	40.500	0.003
C16:0	20.99 $\pm$ 1.58	15.93 $\pm$ 1.70	1	14.231	0.020
C16:1 (<i>trans</i> 9)	0.01 $\pm$ 0.00	0.02 $\pm$ 0.01	1	.500	0.519
C16:1 (<i>cis</i> 9)	1.21 $\pm$ 0.13	0.67 $\pm$ 0.10	1	32.564	0.005
C17:0	0.04 $\pm$ 0.01	0.02 $\pm$ 0.00	1	12.000	0.026
C17:1 (<i>cis</i> 9)	0.13$\pm$0.02	0.05$\pm$0.00	1	84.500	0.001
C18:0	0.31$\pm$0.01	0.19$\pm$0.01	1	342.250	<0.001
C18:1 (<i>cis</i> 9)	0.44 $\pm$ 0.08	0.33 $\pm$ 0.06	1	4.070	0.114
C18:1 (<i>cis</i> 11)	10.21 $\pm$ 0.43	8.18 $\pm$ 0.29	1	46.061	0.002
C18:1 (<i>cis</i> 12)	0.04 $\pm$ 0.01	0.02 $\pm$ 0.00	1	12.000	0.026
C18:2n6	1.63$\pm$0.13	9.78$\pm$0.51	1	717.533	<0.001
C18:2n3	0.01 $\pm$ 0.00	0.02 $\pm$ 0.01	1	12.500	0.024
C20:0	0.06 $\pm$ 0.01	0.04 $\pm$ 0.00	1	25.000	0.007
C18:3n6	0.26$\pm$0.01	1.17$\pm$0.12	1	166.156	<0.001
C18:3n3	12.42$\pm$0.17	16.75$\pm$0.77	1	89.938	0.001
C20:2n6	0.01 $\pm$ 0.00	0.05 $\pm$ 0.01	1	32.000	0.005

C22:0	1.05±0.29	0.47±0.05	1	12.101	0.025
C20:3n6	0.14±0.01	0.51±0.06	1	136.800	<0.001
C22:1 (<i>cis</i> 13)	0.09±0.01	0.07±0.01	1	16.200	0.016
C20:4n6	0.24±0.01	2.09±0.17	1	361.140	<0.001
C22:2n6	0.02±0.01	0.00±0.00	1	2.000	0.230
C20:5n3	1.04±0.02	1.05±0.05	1	.211	0.670
C24:0	0.09±0.03	0.02±0.00	1	16.531	0.015
C24:1 (<i>cis</i> 15)	0.01±0.00	0.01±0.00	1	4.500	0.101
C22:4n6	0.06±0.00	0.44±0.02	1	627.200	<0.001
C22:5n3	2.57±0.15	0.86±0.11	1	255.001	<0.001
Unidentified	45.78±2.11	40.78±1.88	1	9.544	0.037
SFA	23.55±1.76	17.04±1.76	1	20.358	0.011
MUFA	12.22±0.65	9.39±0.19	1	52.852	0.002
PUFA	18.44±0.36	32.79±0.22	1	3506.722	<0.001
n6-FA	2.38±0.14	14.08±0.60	1	26.430	0.007
n3-FA	16.03±0.35	18.69±0.82	1	1078.298	<0.001
n6/n3	0.15±0.01	0.76±0.06	1	240.890	<0.001

689 **Table 2.** Amino acids profile (g 100 g⁻¹ DW) in wild type (WT) and sterile mutant
690 (SM) strains of *Ulva rigida*. Data are the means $\pm$ standard deviation (n = 3). DF, F,
691 and *P* values were from the MANOVA. Bonferroni adjustment with a new alpha level
692 of 0.0024 was used to reduce the chance of Type I error. The amino acids with
693 significant difference between two strains are in bold. EAA = essential amino acids,
694 NEAA = non-essential amino acids.

Amino acids	Wild Type	Sterile Mutant	DF	F	<i>P</i> value
Histidine	0.36$\pm$0.01	0.22$\pm$0.02	1	83.248	0.001
Serine	1.27$\pm$0.08	0.85$\pm$0.04	1	64.233	0.001
Arginine	1.57 $\pm$ 0.20	0.93 $\pm$ 0.03	1	29.797	0.005
Glycine	0.83 $\pm$ 0.10	0.77 $\pm$ 0.06	1	0.918	0.392
Aspartic acid	1.94 $\pm$ 0.24	1.01 $\pm$ 0.04	1	42.525	0.003
Alanine	1.98 $\pm$ 0.11	1.47 $\pm$ 0.11	1	30.452	0.005
Glutamic	2.21$\pm$0.05	1.41$\pm$0.11	1	120.476	<0.001
Threonine	1.15 $\pm$ 0.15	0.89 $\pm$ 0.09	1	6.305	0.066
Proline	0.96 $\pm$ 0.10	0.66 $\pm$ 0.04	1	24.223	0.008
Lysine	1.24 $\pm$ 0.11	0.79 $\pm$ 0.09	1	28.358	0.006
Tyrosine	0.58 $\pm$ 0.05	0.46 $\pm$ 0.03	1	12.000	0.026
Methionine	0.42 $\pm$ 0.04	0.38 $\pm$ 0.04	1	1.194	0.336
Valine	1.39 $\pm$ 0.05	1.12 $\pm$ 0.10	1	18.229	0.013
Isoleucine	0.81$\pm$0.02	0.65$\pm$0.03	1	52.285	0.002
Phenylalanine	1.17 $\pm$ 0.13	1.09 $\pm$ 0.09	1	0.726	0.442
Cysteine	0.18 $\pm$ 0.02	0.15 $\pm$ 0.01	1	6.487	0.064
Leucine	1.12 $\pm$ 0.06	1.10 $\pm$ 0.09	1	0.135	0.732
Total	19.18$\pm$0.54	13.93$\pm$0.55	1	3790.801	<0.001
EAA	9.22 $\pm$ 0.49	7.17 $\pm$ 0.49	1	18.732	0.007
NEAA	9.95$\pm$0.17	6.76$\pm$0.05	1	869.035	<0.001
EAA: NEAA	0.93 $\pm$ 0.05	1.06 $\pm$ 0.06	1	8.176	0.046

695
696
697
698

699
700

Fig. 1

701
702

703
704

Fig. 2

705

706

Fig. 3

707
708

709
710

Fig. 4

711
712

713
714

Fig. 5

715
716

717
718

Fig. 6