Manning DAC, Baptista J, Sanchez-Limon M, Brandt K.

[Testing the ability of plants to access potassium from framework silicate minerals.](#)

Science of the Total Environment 2016, 574, 476-481.

Copyright:

© 2016. This manuscript version is made available under the [CC-BY-NC-ND 4.0 license](#)

DOI link to article:

<http://dx.doi.org/10.1016/j.scitotenv.2016.09.086>

Date deposited:

11/09/2016

Embargo release date:

14 October 2017

This work is licensed under a

[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International licence](#)

1 **Testing the ability of plants to access potassium from framework silicate**
2 **minerals**

3

4 ¹David A C Manning*, ¹Joana Baptista, ²Mallely Sanchez Limon and ³Kirsten
5 Brandt

6

7 ¹School of Civil Engineering & Geosciences, Newcastle University, Newcastle
8 upon Tyne, UK, NE1 7RU

9 ²University Claude Bernard Lyon 1, 43 Boulevard du 11 Novembre 1918, 69100
10 Villeurbanne, France

11 ³School of Agriculture, Food and Rural Development, Newcastle University,
12 Newcastle upon Tyne, UK, NE1 7RU

13

14 *Corresponding author <David.Manning@newcastle.ac.uk>

15

16

Testing the ability of plants to access potassium from framework silicate minerals

ABSTRACT: The availability of K, essential for plant growth, from syenite (a silicate rock in which potassium feldspar is the dominant mineral; >90 wt%), and phlogopite mica has been demonstrated using carefully designed plant growth pot experiments in which the only added source of K was the mineral of interest, with no loss of nutrients through drainage. Using pure quartz sand as a soil, both growth (increase in diameter) of leek plants and K-content of the plant material showed a dose-dependent positive response to the application (114-43000 mg K/pot) of milled syenite with increases in plant diameter of 0.5-0.7 mm/week, increasing with application rate. Phlogopite mica (114-6000 mg K/pot) supported the highest observed increase in diameter (approx. 1 mm/week) and plant K-content, both similar to that observed for a positive control (KCl). These experiments demonstrate that plants can obtain K for growth from milled syenite, in which feldspar is the dominant K-bearing mineral, and confirm previous observations that micas can be an effective source of K.

Keywords: Potassium, soil, fertiliser, mineral, silicate, feldspar

1. Introduction

For healthy growth, all plants require a range of major and trace nutrients. With the exception of nitrogen, the ultimate source of all nutrients is the geological mineral assemblage that occurs within a soil. The function of soil

42 minerals varies, with (for example) congruent dissolution of apatite providing P,
43 and cation exchange on clay surfaces controlling K and N (as ammonium) within
44 the soil solution.

45 This paper focuses on the availability of K for plant growth, and
46 demonstrates for the first time that K is available for plant growth when
47 potassium feldspar (KAlSi_3O_8) is the dominant source of K in a soil. Potassium
48 feldspar occurs widely in (especially temperate) soils, yet is regarded as inert on
49 the timescale of crop production given its slow rate of dissolution as determined
50 in the laboratory and low K yields in conventional laboratory tests of K
51 availability (Harley and Gilkes, 2000; Manning, 2010a; White and Brantley,
52 1995). Other widely occurring potassium silicate minerals include micas, which
53 have long been known to act as sources of K through cation exchange and
54 weathering reactions (Sparks and Huang, 1985; Mohammed et al., 2014).
55 Similarly, there is long-standing evidence that nepheline-bearing rocks
56 (including nepheline syenites; nepheline is also an aluminosilicate mineral, a
57 feldspathoid) may be effective as sources of K for plant growth (Ciceri et al.,
58 2015). Conventionally, potassium fertilisers are based on soluble salts, derived
59 from the natural mineral sylvite (KCl) or sulfates such as polyhalite
60 ($\text{K}_2\text{SO}_4 \cdot 2\text{CaSO}_4 \cdot \text{MgSO}_4 \cdot 2\text{H}_2\text{O}$) and carnallite ($\text{MgCl}_2 \cdot \text{KCl} \cdot 6\text{H}_2\text{O}$).

61 Globally, inputs of K to soils from all sources (including crop residues) are
62 much less than the amount removed through the harvesting of crops (Sheldrick
63 et al., 2002). Nutrient balance studies show that if this deficit was an accurate
64 representation of the need for K to support crop growth, then the requirement
65 for supplementation with K would be far greater than the need for P or N, which
66 more often are in balance (Sheldrick et al., 2002). Indeed, if all K removed

67 annually by crops had to be replaced to fill the 'potash gap' (Manning, 2015),
68 world mined production of potash would need to double. The ability of many
69 soils to sustain reasonable yields without annual supply of K thus indicates that
70 their content of plant available K may be replenished from sources already
71 present in the soil (Mohammed et al. 2014)

72 At present, global production of K fertilizers is about 33.5 million tonnes
73 K₂O equivalent annually (Jasinski, 2014), typically mined from deposits of
74 natural K chloride and sulfate salts, some of which contain other nutrients. 99%
75 of world potash is mined from just 12 countries, dominated by Canada
76 (Rittenhouse, 1979), and whole continental regions have insignificant (e.g.
77 Africa, Australasia), or low production (e.g. South America). The price of K
78 fertilizers reached US\$1000/tonne in 2009 (Manning, 2010a), reducing to
79 US\$350-400/tonne more recently. In addition to paying the commodity price
80 and for value added during processing, farmers must pay high associated
81 transport and distribution costs. Additionally, K from sources such as KCl is
82 highly soluble and can be rapidly lost through leaching, so (depending on the
83 soil) it may need to be replaced for every growing season.

84 To supply the K required for crop growth it is therefore important to
85 consider novel inputs in addition to conventional sources, given constraints on
86 price and availability. K silicate minerals including feldspar and mica contain up
87 to 16 wt% and 11 wt% K₂O respectively, and have been used in a number of
88 studies (Ciceri et al., 2015). In particular, Leonardos et al. (1987) present a
89 convincing case for the use of silicate minerals as sources of K for deeply
90 weathered tropical soils, where leaching of conventional salt-based fertilisers
91 often removes nutrients even before they can be used by the growing plants. But

although some trials have given positive results (e.g. Bakken et al., 1997; 2000; Sanz-Scovino and Rowell, 1988; Mohammed et al, 2014; Manning, 2010b), other trials have failed to do so (e.g. Bolland and Baker, 2000; Harley and Gilkes, 2000; Hinsinger et al., 1996), and the generally low price, historically, of conventional K fertilisers has reduced the need for alternatives.

A critical aspect of the use of novel sources of K is the unambiguous demonstration that K within a plant is derived from a known source. In this paper, we report the design and results of experiments in which we show that K can be derived from (a) milled syenite rock in which potassium feldspar is present as the dominant K-bearing mineral, or (b) mica (the mineral phlogopite). We believe that this is the first time that K availability from a feldspathic rock to plants has been demonstrated experimentally, and our results support the future design and use of silicate minerals as fertilisers particularly in deeply-leached tropical soils in which oxide minerals predominate.

2. Materials and Methods

Plant growth experiments were carried out using closed system pot trials in which there was no through drainage that would permit unknown loss of nutrients from the pot, and in which the dominant source of K was the silicate mineral of interest. An artificial soil was used, based on a high purity silica sand ('Sibelco RHT': >99% quartz and >99% 125-710 µm grain size, www.sibelco.eu), which was mixed with 20% organic matter (Irish moss peat – fine, Recipe no 822, East Riding Horticulture, www.eastridinghorticultureltd.co.uk) by volume, corresponding to approximately 3% by weight and adding potentially 4-8 mg K (based on supplier's specification) equally to each pot. The K sources were

syenite (12.8% K₂O) supplied by Terrativa SA (Belo Horizonte, Brazil) and phlogopite mica (10.19% K₂O) supplied by LKAB minerals (product PD900; www.lkabminerals.com). The syenite was supplied milled, and the phlogopite was supplied as flakes (approx. 5 mm or less) and was milled using a TEMA mill prior to use. Both were sieved to <0.150 μ m. The chemical compositions of the materials used (Table 1) were determined by X-ray fluorescence spectroscopy using a Panalytical PW2404 wavelength-dispersive sequential X-ray spectrometer (School of Geosciences, Edinburgh University). Major elements were determined on fused glass discs, 40mm-diameter, prepared from approximately 0.9g of sample powder mixed with 4.5 g lithium metaborate flux and fused in Pt-5%Au crucibles at 1100°C. Mineralogically (based on petrography), the syenite is composed of >90% K-feldspar (microcline), with <10 wt% each of pyroxene, amphibole and accessory minerals, all of which are K-free, and biotite. X-ray diffraction (Supplementary Material) did not detect biotite (or any other mica) in the syenite used in these experiments; this technique would normally detect the (001) reflection of micas at >1 wt%.

Wt %	phlogopite	syenite 1	syenite 2	syenite 3	sand
SiO ₂	41.97	62.55	62.60	62.77	98.89
Al ₂ O ₃	9.54	16.26	16.30	16.40	0.13
Fe ₂ O ₃	9.05	2.81	2.81	2.80	0.26
MgO	24.16	0.69	0.70	0.72	0.23
CaO	0.62	2.11	2.16	2.15	0.03
Na ₂ O	0.02	1.50	1.52	1.52	n.d
K ₂ O	10.2	12.78	12.79	12.70	0.019
TiO ₂	0.18	0.31	0.31	0.33	0.031
MnO	0.04	0.07	0.07	0.07	n.d
P ₂ O ₅	0.12	0.19	0.20	0.20	0.012
LOI	3.82	0.50	0.35	0.33	0.13
Total	99.85	99.75	99.81	99.98	99.71

Table 1. Chemical composition of mineral materials used in the experiments. The three syenite samples are replicates of the material used.

In addition, the particle size distribution of the materials used was determined using a Malvern Mastersizer 2000 (Malvern Instruments Ltd, Malvern, UK); the results show 90% of the milled syenite was <150 µm, and 90% of the milled phlogopite was < 60 µm (Supplementary material; Figure S2). Particle surfaces were observed using a Carl Zeiss SIGMA HD VP Field Emission scanning electron microscope combined with an Oxford AZtec ED X-ray analysis system to confirm mineral identification (School of Geosciences, Edinburgh University).

Characterised by ourselves, these commercial materials were chosen to ensure (as far as possible) use of tightly specified materials that are available from commercial sources for future experiments or commercial use.

There were ten different K treatments (Table 2): four different doses of milled syenite, 3 doses of phlogopite, 2 doses of KCl and a K-free control treatment, each with ten replicates (pots). Each 13 cm pot with 1 kg capacity was labelled with a 3-digit code obtained from a random number generator, arranged in a randomized design, giving a total of one hundred pots.

Syenite (12.8 % K ₂ O)	
g mineral/pot	mg K/pot
406	43000
56.8	6000
7.87	830
1.08	114
Phlogopite (10.2 % K ₂ O)	
g mineral/pot	mg K/pot
71.0	6000

9.84	830
1.35	114
KCl (63 % K ₂ O)	
g mineral/pot	mg K/pot
0.22	114
0.11	57.0

Table 2. Application rates of K fertilisers used in the pot trial experiment.

Leek (*Allium ampeloprasum* L. var, 'Oarsman' F1 (A.L. Tozer Ltd., www.tozerseeds.com)) was chosen as the trial plant. Its growth as a sheaf of concentric leaves (Hay and Kemp, 1992; Tsouvaltzis et al., 2010) allows periodic measurement of the increase in diameter, facilitating measurement of growth without killing the plants to determine biomass. Leek seeds were sown in moist compost in a greenhouse three months before starting the trials, and were watered by hand twice a week to deplete the stores of potassium and other nutrients. The seedlings were carefully rinsed with tap water to remove as much compost as possible from the roots, and then wrapped in paper towel to absorb excess water. Plants were sorted according to size in 2 groups (small and large), and two seedlings were selected for each pot, one of each size, pairing the largest plants with the smallest plants to minimise variation in combined size. The diameter, height, and weight were recorded before being planted in the pot. Each pot was placed on an individual saucer to prevent transfer of water or nutrients from one pot to the other, and the pot experiment was carried out in an unheated greenhouse for a period of 10 weeks, between April and July, providing a soil temperature regime representative of subtropical or tropical regions.

Once a week, each pot was provided with 200ml of a K-free Hoagland's nutrient solution containing all required nutrients except K (modified from Johnson et al., 1957; Supplementary Information Table S1).

A constant water tension was applied to each pot by supplying deionized water through a capillary irrigation system (Hydrospike HS-300; www.hydrospike.com), which ensured that the water availability would be appropriate for plant growth under field conditions irrespective of the composition of the soil (Thorup-Kristensen 1994). This irrigation system involved a ceramic spike positioned into the soil in each pot, attached by a watering tube to a water reservoir placed approx. 60cm below the pots. The surface of the reservoir was allowed to recede by no more than 10cm by refilling to initial height at least once every week. The water in the reservoir was replaced once every month to prevent growth of algae, which may clog the tip of the watering tube. With this system the moisture condition was identical for all the treatments, irrespective of differences in water use by plants of different sizes (Thorup-Kristensen 1994), and there was no through drainage.

The diameter of each plant 1 cm above the soil surface was measured weekly, using a digital caliper (Mohammed et al., 2014). The increase in plant diameter since planting was calculated by subtracting the initial diameter (T_0) from each of the subsequent measurements (T_n). Signs of nutrient deficiency in the leaves, such as yellow leaves and chlorosis, were recorded.

After 10 weeks, the whole plants were harvested and the roots were washed gently to remove soil particles, excess water was removed using paper towels. The shoots and roots were placed in separate aluminium trays, dried in an oven at 65°C for 4 days and the dry biomass recorded.

After 2 hours additional drying (to facilitate milling), the dried samples were ground in an electric coffee mill. Dried shoots from the two plants from each pot were combined into one sample per pot. Approximately 2g of ground plant material per sample was then analysed for K, Na, Ca and Mg. The dried samples were digested in a mixture of concentrated HNO₃ and HCl (1:2). Element concentration in the acid digest was determined after suitable dilution for each sample by atomic absorption spectrometry (Varian SpectraAA-400). Chive (*Allium schoenoprasum* L.; WEPAL-IPE sample 111, Wageningen University) was used as a plant reference material, to verify the accuracy of the analysis. The K content of the reference was found to be 21.1 g/kg (n = 3; standard deviation = 0.7 g/kg) compared to the declared consensus value of 22.6 g/kg (n = 128; standard deviation = 1.36 g/kg).

For statistical analysis, the trial had one factor: fertiliser. There were ten fertiliser treatments, including the controls, each with 10 replicates and ten sampling times. The plants in 11 pots (0-3 in each treatment) appeared to be affected by disease, which resulted in very low biomass (<30% of the average within the treatment), the data from these pots were treated as outliers and omitted from the statistical analysis. Growth rate was calculated for each plant as the slope of the change in diameter across all 10 weeks if R²>0.85 (this was fulfilled for one or both plants in all pots with healthy plants). The offtake was calculated as the K concentration in foliage multiplied by the foliage dry mass per pot (since the amount of root material was insufficient for measurement of K). The results were tested by analysis of variance (ANOVA) using General Linear Models in Minitab 17 Statistical software, assessing for normal distribution and considering p-values <0.05 significant.

3. Results and Discussion

The observed increase in leek diameter depended on the mineral treatment. It was greatest for the positive control containing KCl, and for the treatment containing phlogopite mica 6000 mg pot⁻¹ (Figure 1). Treatment with different amounts of syenite gave results between the negative control and the positive control, with the change in diameter increasing with greater application rates (R² 0.23, P<0.0008).

Figure 1. Variation in the plant diameter over a period of 10 weeks in the pot trial for the nepheline syenite and phlogopite mica. Data are the mean value of 10 replicates. Error bars represent 1 x SEM

For all treatments an increase in the diameter was observed, reaching a maximum at the end of the experiment (Figure 1). Plants were able to mobilise K from both syenite and phlogopite, showing increased rates of growth compared with the negative control for all but the lowest applications (Figure 2). The phlogopite treatment gave results similar to the KCl control, as observed in other studies using micas (Mohammed et al., 2014). Importantly, the growth rates for the treatments with syenite were significantly ($P < 0.00001$) correlated ($R^2 = 0.3179$) with the log of the syenite dose, and the rates for the treatments with 830, 6000 and 43000 mg K/pot were statistically different from the negative control ($p < 0.007$, $p < 0.001$ and $p < 0.001$ respectively), demonstrating that some of the K from the syenite was available for uptake by plants.

Figure 2. Effect of crushed rock treatments on rate of diameter increase for leeks over ten weeks. Errors bars represent 1x SEM; Tukey Pairwise Grouping indicated by a, b, c.

When considering the K concentration in the plant, significant differences were seen between treatments. K concentration in the treatment with maximum dose of syenite was higher than the negative control ($p < 0.001$), and proportional with syenite dose across all treatments ($R^2 0.797$, $P < 0.000001$). The phlogopite mica treatments (with the exception of the lowest dose treatment; $p > 0.6$) differed from the negative control. The K concentrations for the maximum dose of syenite and the KCl treatment were not statistically different ($p < 0.9$). In the maximum dose for the phlogopite mica, the K content was much greater than all other treatments of syenite and phlogopite mica, and greater than observed for the KCl treatment ($p < 0.01$).

266

267 Figure 3. K concentration in leek plants for different treatments with silicate
 268 minerals. Error Bars x 1SEM; *Tukey Pairwise Grouping indicated by a, b.*

269

270

271 Calculation of offtake (plant K concentration x dry mass; Figure 4) again shows
 272 clearly that K was taken up from syenite treatments; it also shows for all
 273 treatments that no K was released from other sources than the added minerals,
 274 as applications of K were never exceeded by offtake. A comparison of Figures 2
 275 and 4 demonstrates that growth was completely inhibited for treatments that
 276 provided less than 8 mg plant-available K per pot (negative control, and the
 277 lowest dose of syenite and phlogopite, respectively), while treatments providing
 278 8-29mg per pot showed partial K-limitation, and the growth in remaining
 279 treatments was unaffected by K-availability within the timescale of the

experiment. Sufficient K was present in the negative control and lowest dose applications of syenite and phlogopite to give offtake of approximately 8 mg K/pot; this is consistent with the availability of K from the peat compost added in identical quantity to each pot, but may also reflect the K present within the plant following initial growth in compost prior to planting in the trial pots.

Figure 4. Amount of K removed from the soil by plant growth (offtake); *Tukey Pairwise Grouping indicated by a, b, c, d, e and f.*

The results described above were obtained from experiments that lasted only 10 weeks, so while they demonstrate that the plants were able to access some K from milled syenite to sustain growth, and equally clearly demonstrated that large amounts of syenite had no toxic effects on the plants, they did not allow assessment of the rate of degradation of the syenite beyond what may be the contribution from an accessible surface layer or the fine particles that coat

grain surfaces, both created by mechanical impact during the milling process. Figure 5 shows SEM images of feldspar surfaces before and after the 10-week growth experiment with 43000 mg K/pot added as syenite. In these, cleavage steps are clearly visible as are micropores naturally present within the mineral. The feldspar surface is coated with particles $<1\mu\text{m}$, and overall there is little visible change in appearance (apart from reduction in the coverage by fine particles), consistent with the short time for reaction. The amount of K removed by plants from the soil as offtake was equivalent to 0.1-0.2% of the added syenite, which is consistent with the observation of little change in grain surface appearance.

A) feldspar surface before plant growth B) feldspar surface after plant growth

Figure 5. Surfaces of potassium feldspar grains from leek growth experiments, 10 weeks duration; 43000 mg K/pot added as syenite.

The process of release of K from feldspar, the dominant mineral within the syenite, and mica differs. In micas, the crystal structure consists of layers of

negatively-charged aluminosilicate sheets with cations, such as K, in the interlayer site. When in soils, the micas weather with the release of K as a consequence of cation exchange reactions (Hinsinger et al., 1993; Singh and Goulding, 1997; Sparks and Huang, 1985). Vermiculite is produced as a consequence of weathering in soils of magnesium-bearing micas. The feldspars have a structure with a 3-dimensional network of Si and Al coordinated tetrahedrally with oxygen. This structure has to be destroyed to permit release of K to the soil solution.

In comparison with these experiments, on a longer timescale feldspars that have been exposed to soil processes become corroded, as is evident from grains sampled in the field (White and Brantley, 2003; Mol et al., 2003). Such observations confirm the instability of feldspar within the natural soil system, and thus the potential for sustained release of K, which would then become available for uptake by plants.

It is important to bear in mind that the mineral application rates were chosen for the explicit purpose of experimentally testing the availability of K from a silicate-mineral source within a relatively short timescale, and are not intended to lead to recommendations for rates of field application. However, the implications of this research for fertilizer design are significant. We believe that this is the first time that an experiment has been carried out where care was taken to ensure that the soil was free of potassium minerals, or other sources of K, other than what was added as a treatment. Normally, feldspars and potassic clays/micas occur widely within soils developed on a silicate parent rock,

334 especially in temperate climates where physical weathering dominates over
335 chemical weathering.

336 For silicate minerals to be used as a source of K (and perhaps other
337 nutrients), it is important that the mineralogical composition of the host soil is
338 known, and that this is combined with an understanding of the weathering
339 regime. Addition of a K-feldspar to a temperate soil that already contains K-
340 feldspar, as is common in soils derived from glacial till or developed on granite in
341 northern Eurasia or North America, may not give a response. In contrast,
342 addition of K-feldspar (in syenite or other silicate rocks) to a deeply weathered
343 tropical soil that is dominated by oxy-hydroxides and clays such as kaolinite, is
344 more likely to succeed. In such soils feldspar has already been consumed by
345 weathering. Clearly, experiments of longer duration than 10 weeks and that use
346 appropriate soils and crops would be needed to test this further. Additionally,
347 the response observed in these experiments, which used an artificial sandy soil,
348 suggests that feldspar and mica-bearing rocks may be useful as sources of K in
349 temperate sandy soils that have very low K contents, consistent with results
350 reported by Mohammed et al. (2014).

351 The research confirms the long-established finding that biotite or
352 phlogopite micas are able to deliver K to plants. Under the conditions and
353 duration of the present experiment, the plant response from phlogopite was
354 approximately 10% of that of treatments with KCl, since for the KCl treatments
355 the offtake was approx. 40% of the total added K, while 4% of the 830mg K per
356 pot added as phlogopite was taken up into the plant tissue (other phlogopite
357 treatments providing either too little or too much to include in the comparison).
358 This is consistent with the ability of such micas, as they weather, to influence the

K content of the soil solution by cation exchange. Our experiments emphasise how future longer-term experiments must be designed to address the importance of a conceptual model (Sparks and Huang, 1985, Holmqvist et al. 2003) for the availability of K, with fixed K in feldspar, exchangeable K in mica (and vermiculite), and available K in solution (as provided by the dissolution of KCl and other K salts), and to apply this model to deeply weathered tropical soils.

4. Conclusions

Carefully designed pot experiments have been carried out to define the conditions for testing the ability of milled syenite, dominated by the mineral K-feldspar, and milled phlogopite to provide the K needed for plant growth. In these experiments, the only source of K was the treatment. The response of the experimental plant, leek, was proportional to the application rates, both for growth rate and K content of the leaves. These results show that phlogopite is predicted to be a suitable source of K in soils in which K-containing silicates are absent, a condition that holds widely in deeply weathered soils in tropical regions. This may also be the case for syenite and other feldspar-bearing silicate rocks although with a lower initial plant response per g K supplied, however with the expectation that remaining feldspar will continue to release K during a much longer period. In tropical regions with deeply weathered soils, and in temperate soils where K is rapidly removed by through drainage, fertilisers based on silicate minerals should be considered as sources of K.

Acknowledgements

We thank the following companies for supplying the minerals used in this research: Terrativa SA (syenite), LKAB (phlogopite) and Sibelco (silica sand). Mallely Sanchez Limon carried out the work as part of an ERASMUS-MUNDUS project placement. The authors gratefully acknowledge technical and instrument support from Mr Michael Botha, Mr Phil Green and Dr Nicola Cayzer as well as helpful advice from Mr Othman Qadir.

References

- Bakken A. K.; Gautneb H.; Myhr K. 1997. The potential of crushed rocks and mine tailings as slow-releasing J fertilizers assessed by intensive cropping with Italian rygrass in different soil types, *Nutrient Cycling in Agroecosystems*. 47: 41-48.
- Bakken A. K.; Gautneb H.; Sveistrup T.; Myhr K. 2000. Crushed rocks and mine tailings applied as K fertilisers on grassland, *Nutrient Cycling in Agroecosystems*, 56: 53-57.
- Bolland M.D.A.; Baker M.J. 2000. Powdered granite is not an effective fertilizer for clover and wheat in sandy soils from Western Australia. *Nutrient Cycling in Agroecosystems*. 56: 59-68.
- Ciceri, D.; Manning, D.A.C.; Allanore, A. 2015. Historical and technical developments of potassium resources. *Science of the Total Environment*. 502: 590-601.
- FAO 2010. *Current world fertilizer trends and outlook to 2013/14*. Food and Agriculture Organization of the United Nations
<ftp://ftp.fao.org/ag/agp/docs/cwfto14.pdf>.

407 Harley A.; Gilkes, R. 2000. Factors influencing the release of plant nutrient
 408 elements from silicate rock powders: a geochemical overview. *Nutrient*
 409 *Cycling in Agroecosystems*. 56: 11-36.

410 Hay, R.K.M.; Kemp, D.R. 1992. The prediction of leaf canopy expansion in the leek
 411 from a simple model dependent on primordial development. *Annals of*
 412 *Applied Biology*. 120: 537–545.

413 Hinsinger P., Bolland M.D.A.; Gilkes R.J. 1996. Silicate rock powder: effect on
 414 selected chemical properties of a range of soils from Western Australia and on
 415 plant growth as assessed in a glasshouse experiment. *Fertilizer Research*.
 416 45: 69-79.

417 Hinsinger, P.; Elsass, F.; Jaillard, B.; Robert, M. 1993. Root-induced irreversible
 418 transformation of a trioctahedral mica in the rhizosphere of rape. *Journal of*
 419 *Soil Science*. 44: 535–545.

420 Holmqvist, J., Øgaard, A.F., Öborn, I., Edwards, A.C., Mattsson, L., Sverdrup, H.
 421 2003. Application of the PROFILE model to estimate potassium release from
 422 mineral weathering in Northern European agricultural soils. *European Journal*
 423 *of Agronomy* 20: 149-163.

424 Jasinski, S. M., 2014. Potash. United States Geological Survey Minerals Yearbook,
 425 <http://minerals.usgs.gov/minerals/pubs/commodity/potash>.

426 Johnson, C.M.; Stout, P.R.; Broyer, T.C.; Carlton, A.B. 1957. Comparative chlorine
 427 requirement of different plant species. *Plant and Soil*. 8, 337-353.

428 Leonardos O.H.; Fyfe W.S.; Kronberg B.I. 1987. The use of ground rocks in laterite
 429 systems: an improvement to the use of conventional soluble fertilisers?
 430 *Chemical Geology*. 60: 361-370.

431 Manning, D.A.C. 2010a. Mineral sources of potassium for plant nutrition: a
 432 review. *Agronomy for Sustainable Development*, 30: 281-294.

433 Manning, D.A.C. 2010b. Stone meal as a source of plant nutrients, especially
 434 potash: a mineralogical approach. In: *Anais I Congresso Brasileiro de*
 435 *Rochagem, Brasilia, 21-24 de Setembro de 2009*, Eder de Souza Martins and
 436 Suzi Huff Theodoro (editors), Embrapa Cerrados Planaltina, DF, 322pp.

437 Manning, D.A.C. 2015. How minerals will feed the world in 2050. *Proceedings of*
 438 *the Geologists' Association*. 126: 14-17.

439 Mohammed, S.O.; Brandt, K.; Gray, N.D.; White, M.L.; Manning, D.A.C. 2014.
 440 Comparison of silicate minerals as sources of K for plant nutrition in sandy
 441 soil. *European Journal of Soil Science*. 65: 653-662.

442 Mol, G.; Vriend, S.P.; van Gaans, P.F.M. 2003. Feldspar weathering as the key to
 443 understanding soil acidification monitoring data; a study of acid sandy soils in
 444 the Netherlands. *Chemical Geology*. 202: 417-441.

445 Rittenhouse, P.A. 1979. Potash and politics. *Economic Geology*. 74: 353-357.

446 Sanz-Scovino, J.I.; Rowell, D.L. 1988. The use of feldspars as potassium fertilisers
 447 in the savannah of Columbia. *Fertilizer Research*. 17: 71-83.

448 Sheldrick, W.F.; Syers, J.K.; Lingard J. 2002. A conceptual model for conducting
 449 nutrient audits at national, regional and global scales. *Nutrient Cycling in*
 450 *Agroecosystems*. 62: 61-67.

451 Singh, B.; Goulding, K.W.T. 1997. Changes with time in the potassium content and
 452 phyllosilicates in the soil of the Broadbalk continuous wheat experiment at
 453 Rothamsted. *European Journal of Soil Science*. 48: 651-659.

454 Sparks D.L.; Huang P.M. 1985. Release of soil potassium by weathering reactions.
 455 In Minson R.D. (ed) Potassium in Agriculture, pp 202-276. American Society of
 456 Agronomy, Madison WI.

457 Thorup-Kristensen, K. 1994. An easy pot incubation method for measuring
 458 nitrogen mineralization from easily decomposable organic material under
 459 well defined conditions. Fertilizer Research, 38, 239–247.

460 Tsouvaltzis, P.; Siomos, A.S.; Gerasopoulos, D.; Bosabalidis, A.M. 2010. Extension,
 461 anatomy and metabolic activity of leaves in minimally processed leek stalks.
 462 Postharvest Biology and Technology, 2010. 57: 149-154.

463 White, A.F.; Brantley, S.L. 1995. *Chemical weathering rates of silicate minerals*.
 464 *Reviews in Mineralogy* , **31**, Mineralogical Society of America, Washington,
 465 USA.

466 White A.F.; Brantley S.L. 2003. The effect of time on the weathering of silicate
 467 minerals: Why do weathering rates differ in the laboratory and field? Chemical
 468 Geology 202, 479–506.

469

470

Supplementary Information

Table S1. Composition of K-free Hoagland Nutrient Solution

Compound	Molecular weight	Conc. of stock solution (M)	Conc. of stock solution g / liter	Volume of stock solution per liter of final solution (ml)	Element
NH ₄ NO ₃	80.05	1.00	80.05	3.0	N
Ca(NO ₃) ₂ · 4 H ₂ O	236.16	1.00	236.16	4.0	Ca+N
(NH ₄) ₂ PO ₄	115.08	1.00	115.08	2.0	N+P
MgSO ₄ · 7 H ₂ O	246.49	1.00	246.49	1.0	Mg+S
		mM			
MgCl ₂ · 6 H ₂ O	203.31	25	5.082	1.0 ¹	Cl
H ₃ BO ₃	61.84	25	1.516		B
MnSO ₄ · H ₂ O	169.01	2.0	0.338		Mn
ZnSO ₄ · 7H ₂ O	289.55	2	0.575		Zn
CuSO ₄ · 5H ₂ O	249.71	0.5	0.125		Cu
H ₂ MoO ₄ (85% MoO ₃)	161.97	0.5	0.081		Mo
Fe-EDTA ²	346.08	20	6.922	1.0	Fe

1. A combined stock solution is made up containing all micronutrients except iron.
2. Sodium ferric ethylenediamine tetraacetate.

X-ray diffraction

X-ray diffraction analysis was carried out using a PANalytical X'Pert Pro Multipurpose Diffractometer (MPD) with an X'Celerator detector and a secondary monochromator (PANalytical Ltd., Cambridge, UK). Scans were made over the range 2° – 70° , using Cu-K α radiation ($\lambda = 1.54180 \text{ \AA}$). Samples were prepared by packing approximately 500 mg of dry milled sample into 16-mm diameter steel sample wells, and were rotated during analysis. Phase identification was carried out using HighScore Plus software with reference to the ICDD Powder Diffraction File 2 database (1999) and the Crystallography Open Database (February 2012).

Figure S1. X-ray diffraction pattern for syenite.

In this figure, the XRD pattern (Cu K α radiation, $1.541 \text{ \AA}$) for syenite (top trace) is compared with intermediate microcline (green), orthoclase (blue) and phlogopite (red), using CrystalDiffract libraries (www.crystallmaker.co.uk). Note

the absence from the syenite sample of the characteristic (001) phlogopite peak at around $9^{\circ} 2\theta$.

Figure S2. Particle size distribution of minerals used in this study (Malvern Mastersizer 2000; Malvern Instruments Ltd, Malvern, UK)

Table S2. Surface area (geometrical) of minerals used in this study, determined using Malvern Mastersizer 2000.

Material	Surface area (m ² /g)
Phlogopite	1.03
Syenite 1	0.663
Syenite 2	0.661
Syenite 3	0.652