Ersine D, Thomas AJ, Taylor JP, Savage MA, Attems J, McKeith IG, Morris CM, Khundakar AA.

[Neuronal Loss and A-Synuclein Pathology in the Superior Colliculus and Its Relationship to Visual Hallucinations in Dementia with Lewy Bodies.](#)

The American Journal of Geriatric Psychiatry 2017,

<https://doi.org/10.1016/j.jagp.2017.01.005>

Copyright:

© 2017. This manuscript version is made available under the [CC-BY-NC-ND 4.0 license](#)

DOI link to article:

<https://doi.org/10.1016/j.jagp.2017.01.005>

Date deposited:

21/03/2017

Embargo release date:

10 January 2018

This work is licensed under a

[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International licence](#)

Neuronal Loss and A-Synuclein Pathology in the Superior Colliculus and Its Relationship to Visual Hallucinations in Dementia with Lewy Bodies

Daniel Erskine, M.Res., Alan J. Thomas, Ph.D., M.R.C.Psych.,
John-Paul Taylor, Ph.D., M.R.C.Psych., Michael A. Savage, M.Res.,
Johannes Attems, M.D., Ian G. McKeith, Ph.D., F.Med.Sci.,
Christopher M. Morris, Ph.D., Ahmad A. Khundakar, Ph.D.

Objective: Patients with dementia with Lewy bodies (DLB) often experience visual hallucinations, which are related to decreased quality of life for patients and increased caregiver distress. The pathologic changes that contribute to visual hallucinations are not known, but several hypotheses implicate deficient attentional processing. The superior colliculus has a role in visual attention and planning eye movements and has been directly implicated in several models of visual hallucinations. Therefore, the present study sought to identify neurodegenerative changes that may contribute to hallucinations in DLB. **Methods:** Postmortem superior colliculus tissue from 13 comparison, 10 DLB, and 10 Alzheimer disease (AD) cases was evaluated using quantitative neuropathologic methods. **Results:** α -Synuclein and tau deposition were more severe in deeper layers of the superior colliculus. DLB cases had neuronal density reductions in the stratum griseum intermedium, an important structure in directing attention toward visual targets. In contrast, neuronal density was reduced in all laminae of the superior colliculus in AD. **Conclusion:** These findings suggest that regions involved in directing attention toward visual targets are subject to neurodegenerative changes in DLB. Considering several hypotheses of visual hallucinations implicating dysfunctional attention toward external stimuli, these findings may provide evidence of pathologic changes that contribute to the manifestation of visual hallucinations in DLB. (Am J Geriatr Psychiatry 2017; ■■■:■■■-■■■)

Key Words: Dementia, Lewy, visual, superior colliculus, neuropathology, neuropsychiatry

Received November 1, 2016; revised January 5, 2017; accepted January 6, 2017. From the Ageing Research Laboratories (DE, JA, CMM, AAK), Newcastle University, Newcastle upon Tyne, United Kingdom; Biomedical Research Building (AJT, J-PT, IGMK), Campus for Ageing and Vitality, Newcastle University, Newcastle upon Tyne, United Kingdom; Medical Toxicology Centre (DE, CMM), Newcastle University, Newcastle upon Tyne, United Kingdom; and Institute of Neuroscience (MAS), Newcastle University, Newcastle upon Tyne, United Kingdom. Send correspondence and reprint requests to Ahmad Khundakar, Institute of Neuroscience, Ageing Research Laboratories, Edmondson Building, Campus for Ageing and Vitality, Newcastle upon Tyne NE4 5PL, United Kingdom. e-mail: ahmad.khundakar@newcastle.ac.uk

© 2017 American Association for Geriatric Psychiatry. Published by Elsevier Inc. All rights reserved.

<http://dx.doi.org/10.1016/j.jagp.2017.01.005>

Pathology of Visual Hallucinations in DLB

Dementia with Lewy bodies (DLB) is the second most common progressive neurodegenerative dementia after Alzheimer disease (AD), accounting for approximately 4.2% of all dementia cases.^{1,2} DLB is clinically characterized by three core symptoms of fluctuating cognition, parkinsonism, and complex visual hallucinations.³ The defining histologic feature of DLB, along with Parkinson disease and Parkinson disease dementia, is the presence of intracytoplasmic α -synuclein-containing inclusions termed Lewy bodies.⁴ The topographic distribution of Lewy body-related pathology in the respective disorders is believed to underlie the patterns of clinical manifestations.⁵ Lewy body pathology affects brainstem, limbic, and neocortical regions in DLB,⁶ often in combination with the two hallmark histologic features of AD, β -amyloid and tau pathology.³

Visual attention is more abnormal in DLB than in AD or Parkinson disease,⁷ and DLB patients show greater impairment in filtering distracting visual stimuli compared with AD or comparison patients.⁸ Deficits in visual attention have been suggested to contribute to the manifestation of complex visual hallucinations in DLB.^{9,10} Complex visual hallucinations are common in DLB,¹¹ occurring in 60%–80% of cases,¹² and typically consist of objects, such as animals, people, and faces.¹³ Visual hallucinations are related to impaired quality of life for DLB patients,^{14,15} contribute to caregiver distress,¹⁶ and thus represent an important therapeutic target. However, their pathologic etiology is poorly understood.

The superior colliculus has a distinctive laminar structure consisting of seven layers in three major gray matter strata, the stratum griseum superficiale (SGS), the stratum griseum intermedium (SGI) and the stratum griseum profundum (SGP), interspersed with predominantly fiber-rich layers.¹⁷ The superior colliculus may be subject to neurodegenerative changes in DLB because increased saccadic latency, which has been reported in DLB,¹⁸ has also been observed in nonhuman primates after chemical inactivation of the superior colliculus.¹⁹ The superior colliculus is also involved in attentional functions that are impaired in DLB, such as target selection and the filtering of distracting stimuli.¹⁷ One model has directly implicated dysfunction of the dorsal attentional network, with which the superior colliculus is believed to interact, in the manifestation of visual hallucinations in DLB.²⁰ The pathway from the retina to the inferior pulvinar, routed through

the SGS, has also been directly implicated in the manifestation of hallucinations in Lewy body disease.²¹

The aim of the present study was to identify whether the superior colliculus is subject to neurodegenerative changes in DLB using stereologic measures of cell density and densitometric analysis of neuropathologic lesions. Cell density and neuropathologic data were then compared with neuropsychological data obtained during life, where available, and the known function of the individual laminae of the superior colliculus to determine the relationship between neurodegenerative changes and clinical features. DLB cases were compared with aged comparison cases and “disease comparison” AD cases to elucidate whether degenerative changes occur to the superior colliculus in DLB and their relationship to its distinct clinical manifestation.

METHODS

Tissue Preparation

Human postmortem tissue was obtained by a convenience sample from the Newcastle Brain Tissue Resource, and ethical approval was granted by the Newcastle University ethics board and the Joint Ethics Committee of Newcastle and North Tyneside Health Authority (reference no. 08/H0906/136). DLB and AD subjects had been part of several prospective clinical studies and had received detailed clinical assessments according to international consensus guidelines^{3,22} and case note review after death. Cases with psychiatric or neurodegenerative comorbidities were not included in the present study. Neuropathologic assessment was performed according to standardized neuropathologic diagnostic procedures.^{3,23–26} Clinical and pathologic data were collated to establish a clinico-pathologic consensus diagnosis. Three groups of cases were included in the present study: 10 DLB cases, 10 AD cases, and 13 clinically confirmed aged comparison cases that showed no, or only low, age-associated neurodegenerative pathology at *postmortem* examination. Case details are presented in Table 1.

At autopsy, the upper midbrain was dissected from the cerebrum at the level of the third cranial nerve, along a line from the junction of the mammillary body running posteriorly to the upper part of the superior colliculus (Figure 1). For stereologic analysis, 3 × 30 μ m adjacent

TABLE 1. Demographic Information

Case Identification	Gender	Age at Death (yr)	PM Interval (hr)	Braak NFT	McKeith LB	Clinicopathologic Diagnosis	NPI (hall)	NPI Interval to Death (mo)
C5	M	73	25	0	None	Control	—	—
C22	F	80	25	2	None	Control	—	—
C23	F	90	63	2	None	Control	—	—
C24	M	91	72	1	None	Control	—	—
C25	M	83	112	2	None	Control	—	—
C26	M	71	25	1	None	Control	—	—
C27	M	79	39	2	None	Control	—	—
C28	F	93	12	3	None	Control	—	—
C29	M	94	25	2	None	Control	—	—
C30	F	81	75	2	None	Control	—	—
C31	M	84	45	2	None	Control	—	—
C32	M	78	64	2	Brainstem	Control	—	—
C33	F	63	16	2	None	Control	—	—
Mean $\pm$ standard deviation		81.5 $\pm$ 9	46.0 $\pm$ 29	1.8 $\pm$ 1			N/A	N/A
D13	M	75	18	2	Limbic	DLB	—	—
D24	M	77	29	2	Neocortical	DLB	1	2
D6	F	78	96	3	Neocortical	DLB	6	1
D25	F	91	84	5	Limbic	DLB	9	5
D21	M	76	13	2	Neocortical	DLB	0	2
D26	M	74	42	4	Neocortical	DLB	4	9
D27	F	73	99	3	Neocortical	DLB	0	27
D28	M	71	22	3	Neocortical	DLB	2	10
D29	F	87	90	2	Neocortical	DLB	3	—
D30	M	74	60	2	Neocortical	DLB	—	—
Mean $\pm$ standard deviation		77.6 $\pm$ 6	55.3 $\pm$ 35	2.8 $\pm$ 1			3.1 $\pm$ 3	7.8 $\pm$ 9
A2	M	91	22	5	None	AD	0	16
A9	M	68	24	6	None	AD	—	—
A13	M	80	39	6	None	AD	—	—
A14	M	87	21	6	None	AD	0	11
A15	F	80	10	6	Amygdala	AD	0	8
A16	M	82	12	6	Amygdala	AD	0	—
A10	F	86	123	6	Brainstem	AD	—	—
A17	F	58	70	6	None	AD	—	—
A18	M	84	40	6	None	AD	2	10
A19	F	89	11	6	None	AD	—	—
Mean $\pm$ standard deviation		80.5 $\pm$ 10	37.2 $\pm$ 35	5.9 $\pm$ 0			0.4 $\pm$ 1	11.1 $\pm$ 3

Note: N/A stands for not applicable. PM interval represents the interval from death to fixation; Braak NFT represents Braak neurofibrillary pathology stage as outlined in Braak et al.;²⁷ McKeith LB represents Lewy body pathology stage as outlined in McKeith et al.;³ clinicopathologic diagnosis represents the consensus diagnosis made by senior clinicians and a neuropathologist, following case note review and autopsy findings; NPI (hall) represents the score on the hallucinations subscale of the Neuropsychiatric Inventory, as detailed by Cummings et al.²⁸

FIGURE 1. Upper midbrain section stained with Loyez's hematoxylin. The superior colliculus is circled.

sections were cut and stained with cresyl violet. Then, 6- μ m sections were stained with antibodies (KM51 anti- α -synuclein [Leica Biosystems, Milton Keynes, UK], 1:250; AT8 anti-phosphorylated tau [Autogen, Holliston, MA], 1:4,000; 4G8 anti- β -amyloid [Covance, Princeton, NJ], 1:15,000) using Menapath polymer detection kits (Menarini Pharma, Berkshire, UK).

Stereology

Stereologic estimates of neuronal density were made in each of the three prominent gray matter laminae of the superior colliculus (SGS, SGI, and SGP) based on their cytoarchitecture and laminar organization (Figure 2).^{29,30} In cresyl violet sections, neurons were differentiated from glia by the presence of Nissl substance within cytoplasm, a pale nucleus, and a single identifiable nucleolus.³¹

Stereologic analysis was conducted using an AxioVision Z.1 microscope equipped with a motorized stage (Zeiss, Oberkochen, Germany), coupled to a computer with Stereologer software (Bethesda, MA). The rater (DE), working together with the senior investigator trained in stereologic methodology (AAK),

traced an outline around the region of interest (i.e., SGS, SGI, or SGP) using a 2.5 $\times$ objective. Dissector frames were placed in a uniform, random arrangement to calculate the density of cells within a defined region (as described previously).³¹ Neuronal counts were conducted at 63 $\times$ oil-immersion objective using the optical dissector probe. Glial cell counts were calculated in all laminae in dissector frames of 3,500 μ m², with neuron counts calculated in dissector frames of 1,900 μ m². Section thickness did not vary across disease groups in any layer. The mean coefficients of error for neuronal and glial cell estimates was calculated using the Gundersen-Jensen method.³² The mean coefficient of error values for all stereologically obtained data showed acceptable levels of accuracy (<0.15).³³

To evaluate whether neuronal density changes occur as a result of increasing age, postmortem delay, or length of time fixed in formalin, correlational analyses were conducted between these variables and neuronal density in each layer of the superior colliculus across all cases. To assess whether neuronal density is altered based on the duration of disease, correlation analyses were conducted between neuronal density in each layer and duration of disease in DLB and AD cases.

Neuropathology

To quantify neuropathologic lesions, images of each gray matter stratum of the superior colliculus (Figure 2) were taken on a Zeiss AxioVision Z.1 microscope using a DsFi1 camera (Nikon, Tokyo, Japan). Stereologer software was used to delineate a region of interest with a 2.5 $\times$ objective, before placement of dissector frames in a uniform, random arrangement. This method prevented the introduction of bias by giving every area of the region of interest an equal probability of being sampled for analysis. In all cases β -amyloid, tau, and α -synuclein were measured using a 20 $\times$ objective. Approximately eight images were taken per lamina per case within the dissector frames and analyzed using ImagePro Plus v.4.1 image analysis system (Media Cybernetics, Bethesda, MA). Using previously published techniques,³⁴ the mean percentage area of immunopositivity was determined by standardizing red-green-blue thresholds per antibody and applying to all sections per case. Each case thus had a mean value generated per antibody for each stratum.

The percentage area immunoreactive for α -synuclein, tau, and β -amyloid was compared across disease groups

FIGURE 2. The delineation of the laminae of the superior colliculus. Laminae were delineated as reported previously.^{29,30} The stratum zonale (SZ) consisted primarily of glia in the most superficial regions, with sparse, small spindle, or triangular-shaped neurons in regions proximal to the succeeding layer. The SGS contained a more densely distributed population of spindle-shaped or triangular neurons that are slightly larger and more deeply stained than those in the SZ. The stratum opticum (SO) predominantly consisted of glial cells, but some sparse triangular and spindle-shaped neurons, similar to those of the SGS, are also observed. The SGI contained larger, deeply stained, multipolar neurons, which are more frequently encountered in deeper regions of this lamina. The stratum album intermedium (SAI) contained some large and deep-staining neurons reminiscent of the SGI but more sparsely distributed and accompanied by numerous glia and occasional small triangular and spindle-shaped neurons. The SGP had a densely distributed neuronal population, predominantly consisting of medium cells with triangular or oval shapes and no large neurons, as observed in the previous laminae. The stratum album profundum (SAP) was relatively acellular, mostly consisting of fibers, although a sparse population of neurons similar to those in the SGP was observed. The border of the periaqueductal gray (PAG) was noted because of the presence of large, deeply staining neurons on a clearly defined border.

to evaluate the vulnerability of the superior colliculus with neurodegenerative pathology in DLB, AD, and comparison cases. Correlational analyses were conducted between neuropathologic and stereologic data to evaluate the relationship between neuropathology and neuronal density in each disease group.

Clinicopathologic Correlation

DLB patients had been recruited for clinical research studies, during which some had received serial assessments of visual hallucination severity and frequency at frequent intervals until death.^{13,35} Because of the hypothesized role of the superior colliculus in the manifestation of visual hallucinations in Lewy body disease,²¹ neuronal density and pathologic burden were correlated with final Neuropsychiatric Inventory Hallucinations Subscale score (NPI [hall]) in all layers of the superior colliculus²⁸ in DLB cases, where available. Final NPI (hall) scores were available for 8 of 10 DLB cases. The mean interval from final NPI assess-

ment to death was 7.81 ± 3.45 months. The relationship between visual hallucinations and the superior colliculus was not explored in AD because visual hallucinations are not a core feature of AD and were less frequently assessed. Additionally, as a separate disease entity, it is possible that visual hallucinations in AD, when present, have a different underlying pathogenesis to that in DLB.

Statistics

Inspection of Q-Q plots and Shapiro-Wilk tests suggested that demographic, stereologic, and neuropathologic data were either not normally distributed or did not have homogeneity of variance. Therefore, Kruskal-Wallis (KW) tests with post-hoc Mann-Whitney U tests were used for these data. Because of the relatively small sample size, corrections for multiple comparisons were not applied. However, effect sizes were reported using Cohen's *d*. For comparisons

FIGURE 3. Stereologic analysis of neuronal density. Neuronal and glial cell density across experimental groups in each lamina. * $p < 0.05$.

between different laminae, Friedman tests with post-hoc Wilcoxon signed-ranks tests were performed.

RESULTS

Demographics

No significant differences were found between groups in age at death (KW $\chi^2 = 2.461$, $df = 2$, $p = 0.292$), post-mortem delay (KW $\chi^2 = 2.708$, $df = 2$, $p = 0.258$), or length of time fixed in formalin (KW $\chi^2 = 0.472$, $df = 2$, $p = 0.790$; Table 1).

Stereology

In the SGS there was a significant main effect of diagnosis on neuronal density (KW $\chi^2 = 6.579$, $df = 2$, $p = 0.037$). No significant difference was found between comparison cases and DLB. AD cases (mean[$\times 100$] = 0.0054 [95% confidence interval {CI}: 0.0047–0.0061]) had reduced neuronal density compared with comparison cases (mean[$\times 100$] = 0.0067 [95% CI: 0.0059–0.0076]; $U = 27.000$, $p = 0.018$, $d = 1.014$). There was no significant difference in neuronal density between DLB and AD in the SGS, although a trend toward a significant reduction in AD (mean[$\times 100$] = 0.0054 [95% CI:

0.0047–0.0061]) compared with DLB (mean[$\times 100$] = 0.0062 [95% CI: 0.0053–0.0071]) was observed ($U = 26.000$, $df = 1$, $p = 0.07$, $d = 0.60$). There was no significant main effect of diagnosis on glial density in the SGS (Figure 3).

In the SGI there was a significant main effect of diagnosis on neuronal density (KW $\chi^2 = 13.046$, $df = 2$, $p = 0.001$). DLB (mean[$\times 100$] = 0.0030 [95% CI: 0.0025–0.0034]; $U = 18.500$, $p = 0.003$, $d = 1.08$) and AD (mean[$\times 100$] = 0.0029 [95% CI: 0.0026–0.0032]; $U = 13.5$, $p = 0.001$, $d = 1.24$) had reduced neuronal density compared with comparison cases (mean[$\times 100$] = 0.0036 [95% CI: 0.0033–0.0038]). There was no significant difference in neuronal density between DLB and AD in the SGI. There was no significant main effect of diagnosis on glial density in the SGI (Figure 3).

In the SGP there was a significant main effect of diagnosis on neuronal density (KW $\chi^2 = 6.600$, $df = 2$, $p = 0.037$). No significant difference was found between comparison cases and DLB. AD cases (mean[$\times 100$] = 0.0028 [95% CI: 0.0025–0.0031]) had reduced neuronal density compared with comparison cases (mean[$\times 100$] = 0.0036 [95% CI: 0.0031–0.0040]; $U = 27.000$, $p = 0.018$, $d = 1.12$). There was no significant difference in neuronal density between DLB and AD in the SGP. There was no significant main effect of diagnosis on glial density in the SGP (Figure 3).

TABLE 2. Neuropathologic Data Demonstrating Comparisons Between Experimental Groups

Lamina	Protein Target	Control	DLB	AD
SGS	α -Synuclein	0.0005 $\pm$ 0.0013	0.0090 $\pm$ 0.0091 ^a	0.0000 $\pm$ 0.0001
	Tau	0.0071 $\pm$ 0.0219	0.0153 $\pm$ 0.0230	0.0419 $\pm$ 0.0723 ^a
	β -Amyloid	0.0065 $\pm$ 0.0234	0.1315 $\pm$ 0.1514 ^b	0.1770 $\pm$ 0.1617 ^b
SGI	α -Synuclein	0.0010 $\pm$ 0.0026	0.0268 $\pm$ 0.0123 ^a	0.0000 $\pm$ 0.0001
	Tau	0.0219 $\pm$ 0.0598	0.0421 $\pm$ 0.0684 ^b	0.2215 $\pm$ 0.3804 ^b
	β -Amyloid	0.0315 $\pm$ 0.1062	0.1368 $\pm$ 0.1590	0.4177 $\pm$ 0.5238 ^b
SGP	α -Synuclein	0.0039 $\pm$ 0.0112	0.0317 $\pm$ 0.0292 ^a	0.0000 $\pm$ 0.0001
	Tau	0.0228 $\pm$ 0.0606	0.0491 $\pm$ 0.1014	0.3110 $\pm$ 0.4838 ^c
	β -Amyloid	0.0681 $\pm$ 0.2398	0.1260 $\pm$ 0.1385 ^b	0.3891 $\pm$ 0.3491 ^b

Notes: Data are the mean ratio of area of staining to total region of interest $\pm$ standard deviation.

^ap < 0.05 compared with control and AD.

^bp < 0.05 compared with control.

^cp < 0.05 compared with control and DLB.

In DLB cases duration of disease was significantly negatively correlated with neuronal density in the SGI ($r_s = -0.726$, $N = 10$, $p = 0.018$). However, no significant correlations were found between duration of disease and neuronal density in the SGS and SGP. In AD duration of disease was not significantly correlated with neuronal density in any layer.

Neuropathology

Comparisons of neuropathology between groups are detailed in Table 2. In DLB cases there was a significant difference in the degree of α -synuclein pathology across the three layers (Friedman $\chi^2 = 12.600$, $df = 2$, $p = 0.002$; Table 2). The SGI (mean = 0.0268 [95% CI: 0.0180–0.0356]; Wilcoxon $z = 2.803$, $p = 0.005$, $d = 0.85$) and the SGP (mean = 0.0317 [95% CI: 0.0108–0.0525]; Wilcoxon $z = 2.497$, $p = 0.013$, $d = 1.09$) had a significantly greater burden of α -synuclein pathology than the SGS (mean = 0.0090 [95% CI: 0.0025–0.0155]). There was no significant difference in the burden of α -synuclein pathology between the SGI and SGP. There were no significant correlations between α -synuclein pathology and neuronal density across DLB, AD, and comparison cases.

Across DLB, AD, and comparison cases, there was a significant difference in the degree of tau pathology between the three layers (Friedman $\chi^2 = 26.375$, $p < 0.001$; Table 2). The SGI (mean = 0.0929 [95% CI: 0.0074–0.1783]; Wilcoxon $z = 4.330$, $p < 0.001$, $d = 0.32$) and SGP (mean = 0.1241 [95% CI: 0.0130–0.2352]; Wilcoxon $z = 4.031$, $p < 0.001$, $d = 0.46$) had a significantly greater burden of tau pathology than the SGS

(mean = 0.0207 [95% CI: 0.0038–0.0376]). There was no significant difference in the burden of tau pathology between the SGI and SGP. Tau pathology was significantly negatively correlated with neuronal density in the SGI in all cases ($r_s = -0.404$, $N = 33$, $p = 0.022$) and the SGP ($r_s = -0.451$, $N = 33$, $p = 0.009$). However, there was no significant relationship between tau pathology and neuronal density in the SGS.

Across DLB, AD, and comparison cases, there was no significant difference in β -amyloid expression across the three layers of the superior colliculus (Table 2). There were no significant correlations between β -amyloid pathology and neuronal density across DLB, AD, and comparison cases.

Clinicopathologic Correlation

There were no significant correlations between age at death, postmortem delay or fixation time in formalin, and neuronal density in any layer of the superior colliculus. In DLB cases NPI (hall) score was significantly positively correlated with neuronal density in the SGS ($r_s = 0.874$, $N = 8$, $p = 0.005$). No other stereologic or pathologic variables were correlated with NPI (hall) score in DLB cases.

DISCUSSION

This study demonstrates significant reductions in neuronal density in the SGI in DLB compared with significant neuronal loss in all laminae in AD. α -Synuclein

Pathology of Visual Hallucinations in DLB

pathology was not related to neuronal density in any layer in DLB, but tau was inversely related to neuronal density in the SGI and SGP. There was no relationship between α -synuclein pathology and hallucination severity and frequency, as assessed by NPI (hall). However, neuronal density in the SGS was positively correlated with the severity and frequency of visual hallucinations in DLB. Distinct topographic patterns of deposition were observed for α -synuclein and tau pathology, with the SGI and SGP affected more severely than the SGS.

The SGS receives retinal input³⁶ and projects to the medial temporal visual area (area V5/MT) through the inferior pulvinar^{37,38} in a pathway believed to be involved in the nonconscious perception of motion.³⁹ In contrast, the SGI and SGP receive inputs from wide-ranging cortical and subcortical regions, such as motor and association sensory cortices, the locus ceruleus, and dorsal raphe,⁴⁰ and have important roles in the selection of visual targets⁴¹ and in directing the resulting motor output.⁴²

Within the superior colliculus, neuropathologic lesions occurred in a stereotypical manner, with higher levels of α -synuclein and tau found in the SGI and SGP compared with the SGS. This may reflect the SGI and SGP receiving inputs from the dorsal raphe, locus ceruleus, and pedunculo pontine nucleus, which are vulnerable to the accumulation of Lewy body and tau pathology.^{43,44} In contrast, the SGS receives predominant innervation from the retina and primary visual cortex,¹⁷ which do not appear to accumulate significant Lewy body pathology in DLB and develop tau pathology only at the final stage of AD pathology.^{27,45–47} Because tau and α -synuclein may spread throughout the brain in a manner reminiscent of prion protein, these distinct patterns of deposition may reflect greater connectivity with areas that are vulnerable to early neuropathologic lesion formation.⁴⁸ However, a neuropathologic study of the superior colliculus has reported greater tau pathology in the SGI and SGP in individuals with chronic traumatic encephalopathy, indicating that these laminae may have an intrinsic vulnerability to tau pathology.⁴⁹

The present study has demonstrated that significant neuronal loss in the superior colliculus was only found in the SGI in DLB cases. This was in contrast to AD, where neuronal density reductions were found in all laminae of the superior colliculus. However, no significant difference in neuronal density was found

between DLB and AD in the SGS and SGP. DLB had significantly increased tau in the SGI, but not in other layers of the superior colliculus. Because tau was negatively correlated with neuronal density in the SGI and SGP across all cases, this may indicate that tau is driving neuronal density reductions in DLB. The greater levels of tau pathology in AD cases may similarly cause the reductions in neuronal density seen in all laminae of the superior colliculus.

Previous studies in the lateral geniculate nucleus and primary visual cortex^{34,46} have not demonstrated neuronal loss in DLB. In contrast, neuronal loss has been identified in the lateral pulvinar⁵⁰ and in the SGI of the superior colliculus in DLB. One model of visual hallucinations has suggested that DLB patients have difficulty engaging the dorsal attentional network to focus attention on ambiguous stimuli, instead relying on the self-referential default mode network.²⁰ Because the lateral pulvinar⁵¹ and SGI¹⁷ are implicated in visual attention and target selection, respectively, it is tempting to speculate that neuronal losses in these regions could impair directed attention toward visual targets. However, visual attentional measures were not conducted on the present cases, so it is not possible to confirm whether these patients were impaired in these faculties or whether this related to more severe visual hallucinations. Additionally, visual hallucinations in DLB are unlikely to result purely from neuronal loss in the SGI and lateral pulvinar, and these changes likely act in concert with degenerative changes elsewhere in the visual system to elicit visual hallucinations.

Visual dysfunction, including hallucinations, in Lewy body disease have been linked to dysfunctional pathways involved in blindsight, the phenomenon whereby a blind individual may react to moving or emotional visual stimuli.²¹ These pathways implicate the SGS, a region found in the present study to be preserved in individuals with more severe and frequent visual hallucinations. This may indicate that dysfunction of this region, or the pathway through it, is crucial for the occurrence of visual hallucinations. The present study was conducted on a relatively small sample, so it is difficult to draw strong conclusions about the relationship between neuronal number in the SGS and visual hallucinations.

This study was limited by the small sample size as a result of limitations on the availability of upper mid-brain tissue. Additionally, the superior colliculus was sampled on three adjacent slides rather than serially

through the entire z-axis. As a result, volume and total neuronal number could not be estimated. This was because of the superior colliculus being routinely dissected on two different planes (coronal and axial) and the inherent value of upper midbrain tissue. Few cases were available that had NPI data on visual hallucinations obtained during life, and this impacted the number of cases included in the study. However, the use of a comparator group that do not typically manifest visual hallucinations suggests that the reported findings may be related to the vulnerability of DLB patients to visual hallucinations.

In summary, the present study has demonstrated that the superior colliculus is subject to neurodegenerative changes in DLB. Specifically, neuronal loss was only found in the SGI in DLB. In contrast, AD cases had neuronal loss in all layers. Because the SGI is important in directing attention and aiding visual target selection, the present results may indicate dysfunction in these functions in DLB. Because dysfunctional visual attention and target selection have been related to visual hallucinations in DLB, the present results may provide evidence of neuropathologic changes that contribute to

the manifestation of visual hallucinations in DLB. However, these changes are unlikely to induce visual hallucinations on their own; thus, continued study of the visual system is necessary to further understand the pathologic changes that contribute to visual hallucinations in DLB.

Tissue for this study was provided by Newcastle Brain Tissue Resource, which is funded in part by a grant from the U.K. Medical Research Council (G0400074), by NIHR Newcastle Biomedical Research Centre and Unit awarded to the Newcastle upon Tyne NHS Foundation Trust and Newcastle University, and as part of the Brains for Dementia Research Programme jointly funded by Alzheimer's Research U.K. and Alzheimer's Society. This study was funded by the NHS National Institute of Health Research Biomedical Research Unit for Lewy body dementia at Newcastle upon Tyne Hospitals NHS Foundation Trust and Newcastle University.

Mr. Erskine was funded from the NHS National Institute for Health Research Biomedical Research Unit for Lewy body dementia from the Yvonne Emily Mairry bequest.

There are no conflicts of interest.

References

1. Vann Jones SA, O'Brien JT: The prevalence and incidence of dementia with Lewy bodies: a systematic review of population and clinical studies. *Psychol Med* 2014; 44:673–683
2. Heidebrink JL: Is dementia with Lewy bodies the second most common cause of dementia? *J Geriatr Psychiatry Neurol* 2002; 15:182–187
3. McKeith IG, Dickson DW, Lowe J, et al: Diagnosis and management of dementia with Lewy bodies: third report of the DLB Consortium. *Neurology* 2005; 65:1863–1872
4. Spillantini MG, Schmidt ML, Lee VM, et al: Alpha-synuclein in Lewy bodies. *Nature* 1997; 388:839–840
5. Barker RA, Williams-Gray CH: The spectrum of clinical features seen with alpha synuclein pathology. *Neuropathol Appl Neurobiol* 2016; 42:6–19
6. Perry RH, Irving D, Blessed G, et al: Senile dementia of Lewy body type. A clinically and neuropathologically distinct form of Lewy body dementia in the elderly. *J Neurol Sci* 1990; 95:119–139
7. Collerton D, Burn D, McKeith I, et al: Systematic review and meta-analysis show that dementia with Lewy bodies is a visual-perceptual and attentional-executive dementia. *Dement Geriatr Cogn Disord* 2003; 16:229–237
8. Perriol MP, Dujardin K, Derambure P, et al: Disturbance of sensory filtering in dementia with Lewy bodies: comparison with Parkinson's disease dementia and Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 2005; 76:106–108
9. Diederich NJ, Pieri V, Goetz CG: Coping strategies for visual hallucinations in Parkinson's disease. *Mov Disord* 2003; 18:831–832
10. Collerton D, Perry E, McKeith I: Why people see things that are not there: a novel perception and attention deficit model for recurrent complex visual hallucinations. *Behav Brain Sci* 2005; 28:737–757
11. McKeith IG: Consensus guidelines for the clinical and pathologic diagnosis of dementia with Lewy bodies (DLB): report of the Consortium on DLB International Workshop. *J Alzheimers Dis* 2006; 9(3 suppl):417–423
12. Burghaus L, Eggers C, Timmermann L, et al: Hallucinations in neurodegenerative diseases. *CNS Neurosci Ther* 2012; 18:149–159
13. Mosimann UP, Rowan EN, Partington CE, et al: Characteristics of visual hallucinations in Parkinson disease dementia and dementia with Lewy bodies. *Am J Geriatr Psychiatry* 2006; 14:153–160
14. Weintraub D, Moberg PJ, Duda JE, et al: Effect of psychiatric and other nonmotor symptoms on disability in Parkinson's disease. *J Am Geriatr Soc* 2004; 52:784–788
15. Bostrom F, Jonsson L, Minthon L, et al: Patients with dementia with Lewy bodies have more impaired quality of life than patients with Alzheimer disease. *Alzheim Dis Assoc Disord* 2007; 21:150–154
16. Ricci M, Guidoni SV, Sepe-Monti M, et al: Clinical findings, functional abilities and caregiver distress in the early stage of dementia with Lewy bodies (DLB) and Alzheimer's disease (AD). *Arch Gerontol Geriatr* 2009; 49:e101–e104
17. Krauzlis RJ, Lovejoy LP, Zenon A: Superior colliculus and visual spatial attention. *Annu Rev Neurosci* 2013; 36:165–182
18. Mosimann UP, Muri RM, Burn DJ, et al: Saccadic eye movement changes in Parkinson's disease dementia and dementia with Lewy bodies. *Brain* 2005; 128(Pt 6):1267–1276

Pathology of Visual Hallucinations in DLB

19. Aizawa H, Wurtz RH: Reversible inactivation of monkey superior colliculus. I. Curvature of saccadic trajectory. *J Neurophysiol* 1998; 79:2082–2096
20. Shine JM, O'Callaghan C, Halliday GM, et al: Tricks of the mind: visual hallucinations as disorders of attention. *Progr Neurobiol* 2014; 116:58–65
21. Diederich NJ, Stebbins G, Schiltz C, et al: Are patients with Parkinson's disease blind to blindsight? *Brain* 2014; 137(Pt 6):1838–1849
22. McKhann GM, Knopman DS, Chertkow H, et al: The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheim Dement* 2011; 7:263–269
23. Montine TJ, Phelps CH, Beach TG, et al: National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease: a practical approach. *Acta Neuropathol* 2012; 123:1–11
24. Thal DR, Rub U, Orantes M, et al: Phases of A beta-deposition in the human brain and its relevance for the development of AD. *Neurology* 2002; 58:1791–1800
25. Gearing M, Mirra SS, Hedreen JC, et al: The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part X. Neuropathology confirmation of the clinical diagnosis of Alzheimer's disease. *Neurology* 1995; 45(3 Pt 1):461–466
26. Braak H, Muller CM, Rub U, et al: Pathology associated with sporadic Parkinson's disease—where does it end? *J Neural Transm* 2006; 70(suppl):89–97
27. Braak H, Alafuzoff I, Arzberger T, et al: Staging of Alzheimer disease-associated neurofibrillary pathology using paraffin sections and immunocytochemistry. *Acta Neuropathol* 2006; 112:389–404
28. Cummings JL, Mega M, Gray K, et al: The Neuropsychiatric Inventory: comprehensive assessment of psychopathology in dementia. *Neurology* 1994; 44:2308–2314
29. Olszewski J, Baxter D: *Cytoarchitecture of the Human Brain Stem*. Basel: Karger, 1954
30. Wallack MN: Lattices of high histochemical activity occur in the human, monkey, and cat superior colliculus. *Neuroscience* 1988; 25:569–583
31. Khundakar A, Morris C, Oakley A, et al: Morphometric analysis of neuronal and glial cell pathology in the dorsolateral prefrontal cortex in late-life depression. *Br J Psychiatry* 2009; 195:163–169
32. Gundersen HJ, Jensen EB: The efficiency of systematic sampling in stereology and its prediction. *J Microsc* 1987; 147(Pt 3):229–263
33. Herculano-Houzel S, von Bartheld CS, Miller DJ, et al: How to count cells: the advantages and disadvantages of the isotropic fractionator compared with stereology. *Cell Tissue Res* 2015; 360:29–42
34. Erskine D, Taylor JP, Ffrench-McCormick MJ, et al: Changes to the lateral geniculate nucleus in Alzheimer's disease but not dementia with Lewy bodies. *Neuropathol Appl Neurobiol* 2016; 42:366–376
35. Burn DJ, Rowan EN, Minett T, et al: Extrapyrmidal features in Parkinson's disease with and without dementia and dementia with Lewy bodies: a cross-sectional comparative study. *Mov Disord* 2003; 18:884–889
36. Perry VH, Cowey A: Retinal ganglion cells that project to the superior colliculus and pretectum in the macaque monkey. *Neuroscience* 1984; 12:1125–1137
37. Stepniewska I, Qi HX, Kaas JH: Do superior colliculus projection zones in the inferior pulvinar project to MT in primates? *Eur J Neurosci* 1999; 11:469–480
38. Stepniewska I, Qi HX, Kaas JH: Projections of the superior colliculus to subdivisions of the inferior pulvinar in New World and Old World monkeys. *Vis Neurosci* 2000; 17:529–549
39. Lanyon LJ, Giaschi D, Young SA, et al: Combined functional MRI and diffusion tensor imaging analysis of visual motion pathways. *J Neuro-ophthalmol* 2009; 29:96–103
40. Sparks DL, Hartwich-Young R: The deep layers of the superior colliculus. *Rev Oculomot Res* 1989; 3:213–255
41. Sparks DL: Conceptual issues related to the role of the superior colliculus in the control of gaze. *Curr Opin Neurobiol* 1999; 9:698–707
42. Gandhi NJ, Katnani HA: Motor functions of the superior colliculus. *Annu Rev Neurosci* 2011; 34:205–231
43. Dugger BN, Murray ME, Boeve BF, et al: Neuropathological analysis of brainstem cholinergic and catecholaminergic nuclei in relation to rapid eye movement (REM) sleep behaviour disorder. *Neuropathol Appl Neurobiol* 2012; 38:142–152
44. Rub U, Del Tredici K, Schultz C, et al: The evolution of Alzheimer's disease-related cytoskeletal pathology in the human raphe nuclei. *Neuropathol Appl Neurobiol* 2000; 26:553–567
45. Ho CY, Troncoso JC, Knox D, et al: Beta-amyloid, phospho-tau and alpha-synuclein deposits similar to those in the brain are not identified in the eyes of Alzheimer's and Parkinson's disease patients. *Brain Pathol* 2014; 24:25–32
46. Khundakar AA, Hanson PS, Erskine D, et al: Analysis of primary visual cortex in dementia with Lewy bodies indicates GABAergic involvement associated with recurrent complex visual hallucinations. *Acta Neuropathol Commun* 2016; 4:66
47. Beach TG, Carew J, Serrano G, et al: Phosphorylated alpha-synuclein-immunoreactive retinal neuronal elements in Parkinson's disease subjects. *Neurosci Lett* 2014; 571:34–38
48. Goedert M: Neurodegeneration. Alzheimer's and Parkinson's diseases: the prion concept in relation to assembled Abeta, tau, and alpha-synuclein. *Science* 2015; 349:601–610
49. Armstrong RA, McKee AC, Cairns NJ: Pathology of the superior colliculus in chronic traumatic encephalopathy. *Optom Vis Sci* 2017; 94:33–42
50. Erskine D, Thomas AJ, Attems J, et al: Specific patterns of neuronal loss in the pulvinar nucleus in dementia with Lewy bodies. *Mov Disord* 2017; doi:10.1002/mds.26887. in press
51. Benarroch EE: Pulvinar: associative role in cortical function and clinical correlations. *Neurology* 2015; 84:738–747