Bellgrove A, McKenzie PF, Cameron H, Pocklington JB.

[Restoring rocky intertidal communities: Lessons from a benthic macroalgal ecosystem engineer.](#)

Marine Pollution Bulletin 2016, 117(1-2), 17-27.

Copyright:

© 2017. This manuscript version is made available under the [CC-BY-NC-ND 4.0 license](#)

DOI link to article:

<http://doi.org/10.1016/j.marpolbul.2017.02.012>

Date deposited:

12/05/2017

Embargo release date:

12 February 2018

This work is licensed under a

[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International licence](#)

Restoring rocky intertidal communities: lessons from a benthic macroalgal ecosystem engineer

Alecia Bellgrove^{a*}, Prudence F. McKenzie^a, Hayley Cameron^{a1}, Jacqueline B.
Pocklington^{b2}

^aDeakin University, Geelong, Australia. School of Life and Environmental
Sciences, Centre for Integrative Ecology, Warrnambool Campus, P.O. Box 423,
Warrnambool, Victoria 3280, Australia

^bDepartment of Zoology, University of Melbourne, Parkville, Victoria 3010,
Australia and Marine Invertebrates, Museum of Victoria, GPO Box 666,
Melbourne, Victoria 3001, Australia

¹Present address: School of Biological Sciences, Monash University, Clayton,
Victoria 3800, Australia,

²Present address: School of Marine Science and Technology, Newcastle
University, Dove Marine Laboratory, Cullercoats, Tyne & Wear NE30 4PZ,
United Kingdom

*Corresponding author. E-mail: alecia.bellgrove@deakin.edu.au, Fax: +61-3-5563-
3462; Address: School of Life and Environmental Sciences, Deakin University, P.O.
Box 423, Warrnambool, Victoria 3280, Australia

Co-author E-mails: PF McKenzie: prue.francis@deakin.edu.au, H Cameron:
hayley.cameron@monash.edu, JB Pocklington: jacqueline.pocklington@ncl.ac.uk

Abstract

As coastal population growth increases globally, effective waste management practices are required to protect biodiversity. Water authorities are under increasing pressure to reduce the impact of sewage effluent discharged into the coastal environment and restore disturbed ecosystems. We review the role of benthic macroalgae as ecosystem engineers and focus particularly on the temperate Australasian fucoid *Hormosira banksii* as a case study for rocky intertidal restoration efforts. Research focussing on the roles of ecosystem engineers is lagging behind restoration research of ecosystem engineers. As such, management decisions are being made without a sound understanding of the ecology of ecosystem engineers. For successful restoration of rocky intertidal shores it is important that we assess the thresholds of engineering traits (discussed herein) and the environmental conditions under which they are important.

Keywords: facilitation; foundation species; fucoid algae; *Hormosira banksii*; positive interactions; sewage

1 Introduction

Coastlines around the world are facing increasing pressure from anthropogenic developments associated with population growth. Disposal of domestic sewage effluent into the marine environment is increasingly becoming a contentious issue as protection of biodiversity comes into the public eye (de-la-Ossa-Carretero *et al.*, 2016; Walker and Kendrick, 1998). Management authorities are actively seeking ways to reduce the

1 impact of the wastewater disposal on the nearshore coastal environment (Adams et al.,
2
3 2008; Molloy et al., 2004; Molloy et al., 2007) and restore biodiversity to disturbed
4
5 regions. Ecosystem engineers – species that create habitat and modify the environment
6
7 and resources for other species (Jones et al., 1994, 1997) – may be appropriate species
8
9 on which to focus restoration efforts because of their direct and indirect interactions
10
11 with a diverse suite of species (Byers et al., 2006; Crain and Bertness, 2006).
12
13
14
15

16 In this review, we examine the role of intertidal benthic macroalgae as ecosystem
17
18 engineers and focus particularly on the temperate Australasian fucoid *Hormosira*
19
20 *banksii* as a case study for rocky intertidal restoration efforts. *H. banksii* provides an
21
22 excellent system for studying ecosystem engineering and restoration. It is an important
23
24 habitat-forming brown macroalga that dominates much of the southern Australasian
25
26 coastline (Osborn, 1948; Schiel, 2004); it contributes tangibly to the biodiversity of
27
28 rocky intertidal communities, and where this species is lost (or reduced), the
29
30 biodiversity is also changed (Brown et al., 1990; Lilley and Schiel, 2006). Furoid algae,
31
32 including *H. banksii*, have been shown to be very sensitive to anthropogenic
33
34 disturbances such as human trampling (Araujo et al., 2009; Keough and Quinn, 1998;
35
36 Schiel and Taylor, 1999), coastal sedimentation (Chapman and Fletcher, 2002; Schiel et
37
38 al., 2006) and sewage effluent discharge (Bellgrove et al., 1997; Brown et al., 1990;
39
40 Fairweather, 1990). We focus on the impact of sewage effluent discharge on benthic
41
42 macroalgae, using *H. banksii* as a case study for intertidal restoration. Based on past
43
44 restoration efforts of macroalgae (from anthropogenic disturbances), we suggest
45
46 avenues for study that will lead to an improved understanding of the roles of fucoid
47
48 algae in community and ecosystem function, with the aim of improving restoration
49
50 efforts of rocky intertidal shores affected by sewage effluent pollution.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2 Intertidal benthic macroalgae as ecosystem engineers

Autogenic ecosystem engineers are species that create habitats, changing the environment by their physical presence (*e.g.* increased structural complexity) and by directly or indirectly modifying the abiotic conditions and/or biotic interactions between species, and consequently the availability of resources (Jones *et al.*, 1994, 1997). We note that other authors have used different terminology for the same concept stressing the importance of positive interactions, including *foundation species* (Bruno and Bertness, 2001; Dayton, 1972) and *facilitators* (Altieri and van de Koppel, 2014; Bruno and Bertness, 2001), but we prefer the simplicity and operational definitions (Jones *et al.*, 1994) of *ecosystem engineers*. We contrast ecosystem engineers to *keystone species*, whose activities and abundance determine the integrity and persistence of a community based on trophic and/or competitive interactions (Paine, 1966, 1969a, b). We emphasise that understanding the variation in the strength of the interactions between species (particularly relative to abundance) is integral to assessing their role in community/ecosystem structure and for conservation and management of these systems (Bruno and Bertness, 2001; Crain and Bertness, 2006; Jones *et al.*, 1997; Menge *et al.*, 1994).

2.1 Interactions with habitat-forming intertidal benthic macroalgae

Macroalgae are often dominant space-holders in intertidal and shallow subtidal regions, particularly on temperate coasts. This section focuses on the current literature highlighting intertidal furoid macroalgae as important ecosystem engineers. Understanding the direct and indirect effects macroalgae can have on the community will provide the necessary insight for restoration ecology practices.

1 Intertidal rocky shores are physically stressful environments, with gradients in wave
2
3 action, heat and desiccation stress. We may expect that in such environments,
4
5 ecosystem engineers that can ameliorate these physical stresses might be particularly
6
7 important, especially at lower latitudes (Bertness and Leonard, 1997; Bertness et al.,
8
9 1999; Crain and Bertness, 2006). Macroalgae can have both direct (providers of habitat
10
11 and shelter) and indirect (*e.g.* habitat facilitation, altered species interactions) effects on
12
13 intertidal community structure, and either may be equally important (Menge, 1995).
14
15 Moreover, changes in the intertidal abiotic conditions by ecosystem engineers may have
16
17 both positive and negative effects on associated species.
18
19
20
21
22
23

24 While many species of foliose algae are often abundant in the rocky intertidal, creating
25
26 and modifying habitats and resource availability for associated species (Gosselin and
27
28 Chia, 1995; Jernakoff, 1986; Kelaher et al., 2001; Sanchez-Moyano et al., 2001; Worm
29
30 and Chapman, 1998), furoid algae are often the dominant habitat-forming algae on
31
32 intertidal rocky shores (Schiel, 2004) and there is much evidence of their importance as
33
34 ecosystem engineers (Fig. 1). The complex three dimensional structure of furoid
35
36 canopies provides more microhabitats than less complex abiotic habitats, and have been
37
38 shown to be important for biodiversity (Bertness et al., 1999; Bishop et al., 2012;
39
40 Bishop et al., 2013; Fredriksen et al., 2005; Hily and Jean, 1997; Jenkins et al., 2004;
41
42 Schiel and Lilley, 2011). Additionally, furoid canopies can ameliorate the physical
43
44 stresses associated with temperature, desiccation and wave action. For example, a
45
46 canopy of *Fucus gardneri* reduced rock surface temperatures and desiccation rates, and
47
48 increased survival of conspecific germlings (van Tamelen et al., 1997). Similarly a
49
50 canopy of *Silvetia compressa* prevented dehydration of agarose beads after 4.1 h
51
52 emersion in summer, and this correlated with 100 % survival of zygotes of *S. compressa*
53
54
55
56
57
58
59
60
61
62
63
64
65

(c.f. 0 % survival on bare rock and in canopy removal plots; Brawley and Johnson, 1991, 1993). For *Ascophyllum nodosum* and *Fucus distichus* on Rhode Island, survivorship of conspecifics high on the shore was greater under natural canopies than under experimentally thinned canopies due to reduced heat and desiccation stress (Bertness and Leonard, 1997). A canopy of *F. distichus* also increased barnacle recruitment on a northwestern Pacific shore where desiccation was high, suggesting that the canopy moderated the desiccation stress (Dayton, 1971). Similarly, on a sheltered shore in the Gulf of Maine, rock temperatures and evaporative water loss were greatly reduced below a canopy of *A. nodosum* compared to canopy removal plots, positively affecting recruitment, growth and survival of understorey species in the high intertidal (Bertness et al., 1999). As well as reducing heat stress, fucoid canopies can provide insulation and reduce thermal stress to understorey species caused by freezing temperatures (McCook and Chapman, 1991; Thompson et al., 1996).

Hydrodynamic forces created by wave action can place severe mechanical stress on organisms living in wave-swept environments and can constrain the sizes of intertidal plants and invertebrates (Blanchette, 1997 and references therein). Although constructed from brittle material, macroalgae are highly extensible and therefore remarkably resilient to the mechanical forces of waves (Denny et al., 1989). Canopies of *Fucus vesiculosus* and *A. nodosum* can greatly reduce wave action, which can reduce mechanical stress for understorey species and create a depositional environment for the entrapment of larvae (Leonard, 1999; McCook and Chapman, 1991; Thompson et al., 1996), and presumably algal propagules. Certainly some studies have found initial propagule settlement to be highest under an adult canopy (Johnson and Brawley, 1998).

While the reductions in physical stresses of the intertidal environment afforded by furoid canopies can have direct positive effects, they may also have indirect positive or negative effects on associated species (Fig. 1). Brawley and Johnson (1991) and Johnson and Brawley (1998) noted that despite high settlement and initial survival of zygotes under a canopy, *S. compressa* recruits were rarely found beneath a canopy. They suggested negative indirect effects of increased herbivory under the canopy or whiplash by adult plants might be possible causes. Whiplash was also found to affect germling survival of *F. gardneri*, but the positive canopy effects of reduced heat and desiccation stress were stronger (van Tamelan et al., 1997). Barnacles have been shown to be negatively affected by whiplash from an algal canopy in some studies (Beermann et al., 2013; Grant, 1977; Leonard, 1999) despite positive effects through amelioration of physical stress (Beermann et al., 2013; Leonard, 1999), while other studies have shown no effect of whiplash (Dayton, 1971; Underwood, 1999).

Dense intertidal algal canopies reduce the amount of light reaching the understory and this can also have direct and indirect, positive and negative effects on understory species. When a canopy of *Fucus serratus* was removed, encrusting coralline algae bleached and died within one week (Hawkins and Harkin, 1985). Similarly, red algal understory turfs (primarily *Chondrus crispus*, *Gelidium* spp. and *Corallina officinalis*) became bleached when an *A. nodosum* canopy was removed, even in winter (Jenkins et al., 1999). In contrast, removals of *Cystoseira* canopies in the Mediterranean resulted in increases in red algal turfs, including articulated corallines, but encrusting corallines were unaffected (Benedetti-Cecchi and Cinelli, 1992; Benedetti-Cecchi et al., 2001; Bulleri et al., 2002); recruitment of *F. vesiculosus* and ephemeral algae were enhanced when *F. vesiculosus* was removed (McCook and Chapman, 1991); and in *A. nodosum*

canopy-removal plots, *F. vesiculosus*, *F. serratus* and *A. nodosum* recruitment were enhanced (Jenkins *et al.*, 1999), suggesting that the shade of the canopies suppressed these species' growth.

Additionally, several studies have recorded changes in the species composition of understory algae and invertebrates in response to canopy removals, which incorporate complex direct and indirect interactions (Fig. 1; Benedetti-Cecchi and Cinelli, 1992; Benedetti-Cecchi *et al.*, 2001; Bertocci *et al.*, 2014; Bulleri *et al.*, 2002; Hawkins and Harkin, 1985; Jenkins *et al.*, 1999; Jenkins *et al.*, 2004; Marzinelli *et al.*, 2012; McCook and Chapman, 1991; Migné *et al.*, 2015; Schiel and Lilley, 2011). Furoid canopies can indirectly affect the abundances of other species by causing changes in consumer pressure (Bertness and Leonard, 1997). For example, the shade provided by a *F. distichus* canopy, while increasing survival of conspecific germlings, also positively affected the densities of the grazing snail *Littorina littorea*, which had an indirect negative effect on growth and survival of *F. distichus* germlings through increased grazing pressure (Bertness and Leonard, 1997). *Fucus evanescens* provides habitat for a diverse taxa of epiphytic algae that facilitate many free-living epifauna (Wikstrom and Kautsky, 2004), which might affect algal recruitment (Buschmann and Bravo, 1990; Buschmann and Vergara, 1993). Green crab (*Carcinus maenus*) predation was higher beneath an *A. nodosum* canopy and probably limited densities of mussels (*Mytilus edulis*) in the understory (Bertness *et al.*, 1999). A canopy of *A. nodosum* indirectly affected the grazing range of the limpet *Patella vulgata* by facilitating red algal turf that limited suitable substrata for limpets (Jenkins *et al.*, 1999). Moreover, the presence of *Fucus vesiculosus* canopies affected the intraspecific competition of small and large *P.*

1 *vulgata* leading to differences in population structure and thus grazing pressure across
2
3 the shore (Marzinelli *et al.*, 2012).
4
5

6
7 Recent research focussing on metabolic processes of canopy-forming fucoids suggest
8
9 they are the most productive systems of intertidal habitats (Bordeyne *et al.*, 2015;
10
11 Crowe *et al.*, 2011a; Gollty *et al.*, 2008; Mign *et al.*, 2015; Tait *et al.*, 2014; Tait and
12
13 Schiel, 2011) and where canopies are lost (removed), productivity is also lost. For
14
15 example, photosynthetic rate decreased by 50% after the removal of the fucoid, *H.*
16
17 *banksii* (Tait and Schiel, 2011). Furthermore, field experiments have shown canopy
18
19 removal of *F. serratus* (and other canopy-forming algae) dramatically reduced the
20
21 community's primary productivity, respiration and biomass levels (Crowe *et al.*, 2011a;
22
23 Mign *et al.*, 2015). It has also been shown that the fucoids *A. nodosum*, *F. vesiculosus*
24
25 and *F. serratus* contribute to most of the carbon fluxes in their zones (Bordeyne *et al.*,
26
27 2015; Gollty *et al.*, 2008). Therefore canopy structure appears to influence metabolic
28
29 processes of algal assemblages, and consequently the removal of fucoid species may
30
31 also negatively impact ecosystem functions such as coastal primary productivity and
32
33 nutrient cycling.
34
35
36
37
38
39
40
41

42 **3 Effects of treated sewage effluent on benthic macroalgae**

43
44

45 Discharge of treated sewage effluent (wastewater) into the nearshore marine
46
47 environment has been shown to have detrimental effects on rocky intertidal and shallow
48
49 subtidal communities around the world (Brown *et al.*, 1990; Bustamante *et al.*, 2012;
50
51 Coleman *et al.*, 2008; Cormaci and Furnari, 1999; Dez *et al.*, 2013; Fairweather, 1990;
52
53 Hirose, 1978; Littler and Murray, 1975; May, 1985). Our understanding of these effects
54
55 in the intertidal zone is summarised in Fig. 2. The dominant brown algae
56
57
58
59
60
61
62
63
64
65

(predominately fucoids or laminarians) that are often important ecosystem engineers, generally decline in abundance or are absent altogether from polluted sites, and are replaced by an increase in the abundance of opportunistic and turf-forming species such as *Ulva* (including *Enteromorpha* Borowitzka, 1972; Brown et al., 1990; Fairweather, 1990; Golubic, 1970; Hirose, 1978; Klavestad, 1978; Littler and Murray, 1975; May, 1985; Tewari and Joshi, 1988), *Ceramium* (Brown et al., 1990; Klavestad, 1978), *Gelidium* (Littler and Murray, 1975, 1978) and *Capreolia implexa* (Brown et al., 1990; as *Gelidium pusillum* May, 1985; Womersley, 1994). These ephemeral and turf-forming species may benefit from high nutrient loads because of their high surface-area:volume ratios and rapid nutrient uptake (Rosenberg and Ramus, 1984), facilitating the high levels of recruitment of ephemeral species observed at polluted sites (Bellgrove et al., 1997).

In some studies the articulated coralline alga, *Corallina officinalis*, has been abundant in the immediate vicinity of a sewage outfall (Littler and Murray, 1975; May, 1985), but in others this species was absent from outfall sites, but formed extensive turfs at intermediately polluted sites (Bellgrove et al., 1997; Bellgrove et al., 2010; Borowitzka, 1972; Brown et al., 1990). Bellgrove *et al.* (1997) suggested that high phosphate concentrations near the outfall may inhibit calcification of the cell walls (following from Brown et al., 1977). Clearly mat-forming algae are more tolerant or actively benefit from anthropogenic stressors such as sewage effluent discharge (Strain et al., 2014).

Sites that are affected by sewage effluent discharge typically show a decrease in species richness (Borowitzka, 1972; Brown et al., 1990; Fairweather, 1990; Littler and Murray, 1975). The reduction in diversity is possibly a consequence of both direct effects of the

sewage effluent on species and indirect effects of the loss of a brown algal canopy. The loss of the important habitat-forming species such as the low-shore laminarian *Eisenia arborea*, resulted in decreased habitat stratification and complexity and probably contributed to the loss of diversity of understory flora and fauna (Littler and Murray, 1975). In fringing areas where the canopy of *E. arborea* was maintained, the changes to the community were minimised. Changes in species richness at effluent-affected sites may also be driven by high nutrient loads favouring certain species that will then out-compete a variety of space-holders, lowering diversity. Recruitment of some fucoid algae in nutrient-enriched conditions has been reduced by competition for space by ulvoid and filamentous opportunistic algae (Korpinen and Jormalainen, 2008; Steen, 2004). However, other studies (Bokn et al., 2003) have shown that opportunistic algae are unable to displace established dominant fucoid algae under enriched conditions unless space is made available.

Although many studies have recorded similar patterns of community change in response to sewage effluent discharge (Fig. 2), we still have relatively little understanding of the mechanisms that are causing these changes. For example, Bellgrove *et al.* (1997) did not find any negative effect of sewage effluent on the recruitment of macroalgae. However, no perennial species, including the pre-discharge dominant *H. banksii* (Manning, 1979), recruited during their study (even at unpolluted shores), which may have been an artefact of the type of artificial substrata used in their experiments (Bellgrove et al., 2004). While field evidence confirming that effluent exposure can limit fucoid recruitment is scarce, laboratory-based toxicity assays have demonstrated that the early developmental stages (fertilization, germination, cell wall formation and cell division) of fucoids can be negatively affected by sewage effluent. In particular,

ammonium concentrations (EC_{50} values of 0.08 mg – 3.19 NL⁻¹ for *H. banksii*) and salinity appear to be the most harmful effluent constituents to early life-stages, but effects are variable amongst developmental stages and amongst studies (Kevekordes, 2001), Myers et al., 2006; (Stauber et al., 2003)).

Discrepancies in reported EC_{50} values for the fucoid *H. banksii* may be due at least in part to different concentrations of ammonium and ammonia (and associated differences in toxicity of the two compounds), with tests run at different temperatures and pH amongst studies (Kevekordes, 2001; Myers et al., 2006; Stauber et al., 2003). The concentrations of ammonia and ammonium are in dynamic equilibrium in solution and are affected moderately by temperature, but dramatically by changes in pH. Thus, to ensure comparable EC_{50} values between different toxicity tests for individual species, it is essential that the protocols used for bioassays are identical and that the pH and temperatures of test solutions are the same. However, if laboratory toxicity tests are to be meaningful for the environmental conditions surrounding the sewage outfalls, then it would be sensible to repeat these bioassays under a range of temperatures and pH conditions that reflect seasonal fluctuations of the coastal receiving waters. Moreover, effluent volumes, total ammonia concentrations and salinity can vary over fine and coarse temporal scales (Batley and Simpson, 2009; Molloy et al., 2004). Thus to be able to set realistic water quality targets for restoration of coastal ecosystems, it is imperative that we also understand the effects of pulses of toxic constituents on the fucoid algae, requiring both carefully designed laboratory and field experiments (e.g. Mayer-Pinto and Ignacio, 2015; Mayer-Pinto et al., 2016) comparing pulses vs press disturbances of effluents under natural conditions.

Reducing the toxicity of sewage effluent discharged into the marine environment alone may not be sufficient to restore furoid populations. There are multiple factors that might limit the recovery of furoids and other lost seaweed species into remediated areas (Bellgrove *et al.*, 2010; Díez *et al.*, 2013), highlighting the need to compliment laboratory studies with manipulative field experiments to appropriately predict outcomes of management decisions aimed at restoring communities affected by sewage effluent discharge. Furthermore, the recruitment potential of early-life stages and long-term survival of adults on remediated shores should be investigated in the field to evaluate the effectiveness of wastewater upgrades in achieving a water quality that permits re-establishment and self-perpetuation of furoid populations.

4 Ecosystem engineers and restoration

Ecosystem engineers may have positive effects on some species and negative effects on others (Bertness *et al.*, 1999; Bruno and Bertness, 2001; Jones *et al.*, 1997). This may increase or decrease species richness and abundances on local scales (Bellgrove *et al.*, 2004; Thompson *et al.*, 1996), but at larger spatial scales that incorporate both engineered and non-engineered environments the net effect on biodiversity is likely to be positive (Bertness and Leonard, 1997; Halpern *et al.*, 2007; Jones *et al.*, 1997); with a mosaic of habitats and associated species and potentially a transfer of energy between them (Bustamante *et al.*, 1995; Estes and Palmisano, 1974; Griffiths *et al.*, 1983; Halpern *et al.*, 2007; Robertson and Lenanton, 1984; Thomsen *et al.*, 2013).

The importance or role of a species as an ecosystem engineer depends on 1) patch size, 2) density/cover, 3) architectural complexity, 4) spatial arrangement/connectivity, and 5) the stability or residence of traits 1 – 4 (Bruno and Bertness, 2001). Furthermore,

there are likely to be thresholds between these five traits and environmental stress (Bruno and Bertness, 2001). In reviewing the current literature, the role, and sometimes the importance, of ecosystem engineers varies along environmental gradients (Crain and Bertness, 2006; but see Viejo *et al.*, 2008) and with environmental context (Crowe *et al.*, 2011b). The example given about the interaction between canopy cover of *H. banksii*, whelk predation and barnacle densities (Underwood, 1999; see *H. banksii* section of review) illustrates the potential for such thresholds.

In addition to the traits of ecosystem engineers defined by Bruno and Bertness (2001), Jones *et al.* (1994, 1997) suggested that 6) the amount of engineered habitat created by an individual in its lifetime, 7) the type and durability of the constructed habitat and impacts once the original engineer has died (or left), and 8) the number of resource flows that are modulated by the engineer and the number of species dependent on these flows will also be important for scaling the impact of an ecosystem engineer. To maximise the effectiveness of restoration efforts, it would thus be ideal to understand the thresholds for each of these traits and the environmental conditions under which they operate. Indeed, there are yet few notable attempts to examine this for intertidal fucoid macroalgae (e.g. Pocklington *et al.*, 2017; Schiel and Lilley, 2007)

5 Lessons from the recovery of intertidal benthic macroalgae

The recovery rates of canopy-forming fucoid algae have been shown to vary both between (e.g. Hawkins and Harkin, 1985; Jenkins *et al.*, 2004) and within (Keough and Quinn, 1998; Lilley and Schiel, 2006; Schiel, 2006) species. Following experimental removals of canopies of *Fucus serratus* from the intertidal, recruitment and recovery of all canopy species was rapid (6 mth to 1 y) (Hawkins and Harkin, 1985). However,

1 while the habitat had been restored, the associated algal assemblages remained different
2
3 from controls after 1 y (Hawkins and Harkin, 1985). In contrast however, removal of a
4
5 canopy of *Durvillaea antarctica* from the lower intertidal zone of an exposed shore
6
7 resulted in an altered assemblage, where turfing articulated coralline algae (*Corallina*
8
9 *officinalis*, *Jania* spp., *Haliptilon roseum*) had increased cover with little recovery of *D.*
10
11 *antarctica* (Taylor and Schiel, 2005).
12
13
14
15

16 Experimental removal of *Ascophyllum nodosum* on the Isle of Man, UK, resulted in a
17
18 mixed canopy of *Fucus vesiculosus* and *F. serratus* that provided similar shade to
19
20 turfing species as the original canopy (Jenkins et al., 1999). However, despite
21
22 successful recruitment (Jenkins et al., 1999), *A. nodosum* had not regained the canopy
23
24 even 12 y post initial removals, apparently due to very slow growth (Jenkins et al.,
25
26 2004). Changes to the community from the initial loss of the *A. nodosum* canopy were
27
28 still present after 12 y (Jenkins et al., 1999; Jenkins et al., 2004). In the Gulf of Maine
29
30 (Bertness et al., 1999), recruitment of *A. nodosum* was comparably high (Jenkins et al.,
31
32 1999), but was restricted to shaded and canopy plots from which grazers were excluded
33
34 high on the shore. Bertness *et al.* (1999) suggested that the amelioration of abiotic
35
36 conditions by the canopy could facilitate recruitment of conspecifics providing positive
37
38 feedbacks (Jones et al., 1997) once a canopy was established (Fig. 1).
39
40
41
42
43
44
45
46

47 In the Mediterranean, removals of canopies of *Cystoseira* spp. have resulted in rapid
48
49 changes to assemblages dominated by coralline turfs (Benedetti-Cecchi and Cinelli,
50
51 1992; Benedetti-Cecchi et al., 2001; Bulleri et al., 2002). Once established the turfs
52
53 appear to prevent the recruitment of the canopy algae such that the changes to the
54
55 assemblages are long-lasting (Benedetti-Cecchi and Cinelli, 1992; Bulleri et al., 2002).
56
57
58
59
60
61
62
63
64
65

The above examples illustrate that the effects of removing ecosystem engineers can be long-lasting, even in cases where there is an ample supply of propagules and good recruitment. For many of these species there is no functional replacement after canopy loss (Schiel, 2006). For anthropogenic disturbances, the negative effects on ecosystem engineers are usually over much larger spatial scales than those adopted for many canopy-removal experiments (Benedetti-Cecchi *et al.*, 2001; Brown *et al.*, 1990; Chapman *et al.*, 1995). For example, recovery of *Fucus gardneri* following broad-scale decimation of populations due to the Exxon Valdez oil spill and subsequent clean up, was initially slow because of low densities of reproductive plants and poor dispersal (van Tamelen *et al.*, 1997). Additionally, habitat characteristics such as substratum rugosity and provision of a canopy (and associated reduction in heat and desiccation), improved recruitment for restoration efforts (van Tamelen *et al.*, 1997).

Recovery of furoid assemblages following wastewater remediation appears to be limited in many parts of the world; with efforts focusing on seeding (Bellgrove *et al.*, 2010; Hernandez-Carmona *et al.*, 2000; Stekoll and Deysher, 1996; Yoon *et al.*, 2014; Yu *et al.*, 2012) and/or transplantation of juvenile or adult thalli (Campbell *et al.*, 2014; Hernandez-Carmona *et al.*, 2000; Whitaker *et al.*, 2010) to sites where the populations were lost, or by artificially increasing the substratum complexity (Deysher *et al.*, 2002; Terawaki *et al.*, 2001; Yu *et al.*, 2012). For example, Soltan *et al.* (2001) noted that even 8 years following upgrades to industrial and domestic effluent treatment in Marseille, France, populations of the furoid *Cystoseria amantacea* were yet to recolonize; even at sites most distant from the point of discharge. Similarly, improvements to the effluent discharged into the Nervión River in northern Spain has allowed some recovery of macroalgal communities along the Basque coast, with these

communities progressing towards a state similar to a reference site (Diez et al., 2009; acknowledging the limitations of sampling only one reference site). However, complete restoration has not been achieved within the low intertidal for any of these sites, which is largely driven by the continued absence of the fucoids *Bifurcaria bifurcaria* and *Cystoseria tamariscifolia*. In Sydney, Australia the subtidal fucoid *Phyllospora comosa* was lost along a 70 km stretch of coastline following discharge of primary-treated sewage effluent and has not recovered naturally since sewage treatment was improved in the 1990's (Coleman et al., 2008). Preliminary transplants of fertile adult plants back into this region is looking promising for restoration even after 18 months post-transplantation (Marzinelli et al., 2014; Marzinelli et al., 2016). Results indicate limited dispersal may have prevented re-establishment following remediation (Campbell et al., 2014).

Most studies investigating recovery of algal assemblages following wastewater remediation (such as the examples above) focus on changes in community composition and abundances of adult plants. While it is informative to track changes in algal communities in response to wastewater remediation, such studies cannot distinguish mechanisms contributing to recovery failure that may need to be regulated for successful restoration to be achieved. Consequently, there is an urgent need for future manipulative studies to untangle potential limitations to recolonization in order for implementation of appropriate restoration initiatives. It is also clear that we must have a thorough understanding of the factors that lead to the initial demise of ecosystem engineers, to ensure that initial drivers of loss are reversed.

An understanding of connectivity between populations is essential for understanding the patterns of gene flow between populations and how they may recover from disturbances (Schiel, 2004). Understanding gene flow is also essential if restoration involves re-establishing ecosystem engineers by augmentation of populations from remnant sites, in order to maintain genetic diversity and avoid genetic pollution (McKay *et al.*, 2005). While populations of marine organisms with a pelagic dispersal stage have traditionally been considered to be ‘open’, with local reproduction and recruitment de-coupled (Underwood and Fairweather, 1989) there is a lot of evidence from both oceanographic modelling (Cowen *et al.*, 2000), field and laboratory experimentation (Santelices, 1990) and genetic studies (Coleman *et al.*, 2011; Kinlan and Gaines, 2003; Williams and Di Fiori, 1996) that suggest for most macroalgae and many invertebrates, dispersal may be limited. In particular, for fucoid algae, with large non-motile and negatively-buoyant eggs, dispersal is often thought to be limited to within metres or 10s m from the parent (Bellgrove *et al.*, 2004; reviewed in Santelices, 1990; Schiel, 2004). However, while the potential for long-distance dispersal of macroalgae by floating fertile fragments has been discussed (Santelices, 1990; van den Hoek, 1987) the importance of this mode of dispersal has had little investigation (Gagnon *et al.*, 2015; Hernández-Carmona *et al.*, 2006; Macaya *et al.*, 2005; McKenzie and Bellgrove, 2008, 2009; Rothausler *et al.*, 2015). For coastal areas affected by anthropogenic disturbances where the habitat-forming fucoid algae often disappear from large stretches of coast (e.g. Benedetti-Cecchi *et al.*, 2001; Brown *et al.*, 1990), an understanding of the dispersal capabilities of the ecosystem engineers is an essential precursor to understanding whether populations can re-establish from distant undisturbed populations, if inhibitory conditions are removed.

6 *Hormosira banksii*: a case study

6.1 *Hormosira banksii* as an ecosystem engineer

In southern Australasia, the only fucoid species that forms extensive canopies in the intertidal zone is the endemic *Hormosira banksii* (Osborn, 1948; Schiel and Taylor, 1999; Womersley, 1987). However, despite its abundance on both wave-exposed and sheltered shores throughout southern Australasia (Bellgrove *et al.*, 2004; Keough and Quinn, 1998; Schiel and Taylor, 1999; Underwood, 1998; Womersley, 1987), we know relatively little about the role *H. banksii* plays in intertidal community structure, with only a handful of studies, but all suggest an important autogenic engineering role (Fig. 1).

Like other perennial fucoids, *H. banksii* is long-lived (>5 y; Kain (Jones), 2015; Keough and Quinn, 1998), and can form extensive mats often with 70 – 100 % cover (Bellgrove *et al.*, 2004; Keough and Quinn, 1998; Schiel and Taylor, 1999; Underwood, 1998).

Recruitment of *H. banksii* was higher under the adult canopy than in an adjacent low-relief mixed algal assemblage (Bellgrove *et al.*, 2004), suggesting either very limited dispersal or amelioration of physical stresses beneath the canopy. For example, reduced wave action might improve ‘stickability’ of zygotes (Taylor and Schiel, 2003) and the moisture maintained beneath the canopy on emersion (Davenport *et al.*, 1999, A.

Bellgrove personal observations; Keough and Quinn, 1998) may facilitate post-settlement survival. The absence of an adult canopy and increased susceptibility to heat and desiccation stress of settlers, may explain slow recovery of *H. banksii* in canopy removal experiments (Schiel, 2006). Thus, there may be positive feedbacks to *H. banksii* through the provision of an extensive canopy (Jones *et al.*, 1997). Vegetative regeneration of senesced fronds from the basal holdfast (Schiel and Taylor, 1999) may

also ensure the relative stability of the canopy in undisturbed conditions (Keough and Quinn, 1998; Schiel and Taylor, 1999).

The thallus of *H. banksii* consists of chains of hollow vesicles that contain internal reserves of water for replenishing dehydrated tissues on emersion (Osborn, 1948). The thallus is thus resistant to desiccation at low tide and also appears to moderate the water loss and heat stress in the understory (Davenport et al., 1999; Keough and Quinn, 1998; but see Pocklington, 2003). Temperature increases of 4 – 9.5 °C have been recorded in canopy removal plots compared to controls (Lilley and Schiel, 2006). The cell walls of *H. banksii* contain phenolic compounds (Schoenwaelder and Clayton, 1999) that have been shown to protect the inner photosynthetic tissues from UV radiation (Schoenwaelder, 2002). The canopy of *H. banksii* is thus also able to provide shade and UV protection for understory species during summer low tides. However, some canopy loss of *H. banksii* occurs in most summers (Keough and Quinn, 1998) due to ‘thallus sunburn’ (Schoenwaelder, 2002) associated with daytime summer low tides exacerbated by barometric influences; and a single, very hot (> 28 °C) spring day caused ~ 30 % reduction in percentage cover of the canopy (Keough and Quinn, 1998).

H. banksii supports a relatively diverse, and in some cases unique, assemblage of understory algae (Bellgrove, 1998; Bellgrove et al., 2004; Lilley and Schiel, 2006; Schiel, 2006), epifauna (Davenport et al., 1999; Schiel, 2006; Schreider et al., 2003) and macroinvertebrates (Bishop et al., 2009; Keough and Quinn, 1998; King, 1992; Schiel, 2006) and can also provide important shelter for the fish *Enneapterygius rufopileus* (Silberschneider and Booth, 2001). Similarly, as for some northern hemisphere fucoids, several studies have shown that articulated and encrusting corallines are rapidly

bleached and fragmented when a canopy of *H. banksii* is removed (Keough and Quinn, 1998; Schiel and Taylor, 1999; Underwood, 1999). However, experimental canopy removals of *H. banksii* in southern New Zealand resulted in initial bleaching of articulated coralline algae, followed by recovery to become the primary benthic cover. Further, removals had positive effects on the abundance of ephemeral algae but an overall negative effect on floral and faunal diversity with a 37 % decline relative to controls, primarily because of strong non-trophic interactions between *H. banksii* and understorey species (Lilley and Schiel, 2006). Another long-term canopy removal experiment in New Zealand, showed that removal of *H. banksii* only resulted in death and removal of articulated coralline turf in the higher mid-intertidal region (0.8 m) at one site (Schiel, 2006). In this case, there was up to 42 % decline in taxon richness in canopy removals compared to controls across sites (Schiel, 2006).

Unlike abiotic habitats (*e.g.* cracks and crevices in rocks), biotic habitats are dynamic and can vary in space and time (Underwood, 1999), where such fluctuations could modify the roles of autogenic engineers (Jones et al., 1997). There is evidence that the amount of cover provided by the canopy of *H. banksii* directly or indirectly affects associated species. Abundances of the grazing snail *Austrocochlea constricta* and percentage cover of articulated and encrusting corallines increased with canopy cover (Keough and Quinn, 1998). *H. banksii* provided shelter for whelks, which increased their effectiveness as predators of barnacles, indirectly affecting the abundance of barnacles (Underwood, 1999). However, this interaction was modified by the size and extent of the *H. banksii* canopy, which is in turn modified by unpredictable processes, such as storm events (Underwood, 1999), extreme thallus sunburn (Keough and Quinn, 1998) or fluctuations in recruitment (Bellgrove et al., 2004). Additionally, no

significant differences in desiccation or temperature in mixed species (including *H. banksii*) canopy and canopy removal plots in Lady Bay (Adelaide, Australia) were found (Pocklington, 2003). However, further investigation revealed that microtopography was highly rugose at this site, and the canopy was patchy with short fronds (Pocklington, 2003). This suggests that both the quality of the canopy-forming alga and the alternative microhabitats available at a site can influence its role as an autogenic engineer.

Varying densities of ecosystem engineers such as fucoid macroalgae, can also be an important mechanism influencing the associated community (Schiel and Lilley, 2007; Speidel et al., 2001). For example, the density of *H. banksii* was manipulated over varying levels on a rocky shore in New Zealand, where results showed that taxon richness declined when the density of *H. banksii* was reduced (Schiel and Lilley, 2007). Additionally, densities of engineers may play an important role in relation to rehabilitation programs, as disturbances (e.g. anthropogenic or restoration programs) may move populations over important density thresholds. When densities of the fucoid, *Fucus gardneri* were manipulated, results showed robustness towards various disturbance intensities as long as a few individuals were present to maintain recruitment (Speidel et al., 2001). Therefore, when *F. gardneri* is faced with a disturbance (e.g. oil spill), and the population is completely removed, we know based on the above findings that recovery will be slow, however if a few individuals remain unaffected recovery will be much quicker (Speidel et al., 2001).

There is considerable morphological plasticity in the thallus form of *H. banksii* (Clarke and Womersley, 1981), with broad differences between estuarine and marine habitats,

and also differences between microhabitats within these habitats (Bishop *et al.*, 2009; Ralph *et al.*, 1998). However, even within morphometric forms associated with microhabitats, particularly on eulittoral rock platforms, there is quite a lot of variability in vesicle attributes, chain length and branching pattern (Bishop *et al.*, 2009; McKenzie and Bellgrove, 2008; Ralph *et al.*, 1998). This means that the three-dimensional structure created by *H. banksii* might vary at different sites and in different microhabitats within sites. Therefore, the role of *H. banksii* as an ecosystem engineer may change depending on its morphology (Bruno and Bertness, 2001; Jones *et al.*, 1997). For instance, Bishop *et al.* (2009) conducted transplant experiments of *H. banksii* thalli from both rocky shore and estuarine habitats into a mangrove forest along the New South Wales coast in eastern Australia. They found that while both algal types enhanced molluscan abundance and species richness, estuarine *H. banksii*, which is characterized by larger vesicles and longer thalli, supported higher diversity and abundances of mollusc assemblages than did rocky shore plants. Subsequent experiments demonstrated that thallus length was most important, but also that the morphology of an intermediary facilitator (mangrove pneumatophores that trap floating *H. banksii* thalli) may affect the epifaunal assemblages (Bishop *et al.*, 2013).

6.2 Recovery rates of *Hormosira banksii*

Although Australasian studies have shown that removal of a canopy of *H. banksii* can in some cases result in bleaching and death of understorey coralline turfs (Keough and Quinn, 1998; Schiel, 2006; Schiel and Taylor, 1999) there is also evidence that where *H. banksii* is weakened by intermediate levels of pollution, articulated coralline turfs can take over dominance (Bellgrove *et al.*, 2010; Brown *et al.*, 1990). It is possible that articulated coralline turfs and fucoid canopies are alternate community states (Petratis

and Latham, 1999) where the switch is intermediate-level pollution that disrupts dominance by *H. banksii*. Recruitment of *H. banksii* into coralline turfs could only occur when turfs had been reduced to either < 2 mm (by human trampling, Schiel and Taylor, 1999) or up to 10 mm (Alestra et al., 2014). Similarly, recovery of *H. banksii* in canopy removal plots may have been impeded by thick (30 mm) coralline turfs (Schiel, 2006) and associated bound sediments (Alestra et al., 2014; Schiel, 2006). It is therefore possible that extensive turfs (up to 40 mm thick, Bellgrove et al., 2010) of *Corallina officinalis* at sites at an intermediate distance to ocean outfalls (Bellgrove et al., 2010; Brown et al., 1990) might inhibit re-establishment of *H. banksii*, even if water quality was conducive to early development and growth. While it has been suggested that recruitment of furoid algae is enhanced in red algal turfs because reduced wave motion facilitates strong zygote attachment and/or the turf structure traps dislodged embryos (Brawley and Johnson, 1991), it is also possible that the reduction in nutrient delivery and shading effects (Hay, 1981) inhibit photosynthesis in developing embryos. In addition, (Bellgrove et al., 2010) found coralline turfs inhibited *H. banksii* recruitment in field and laboratory studies. They suggested surface tension holding propagules above the substratum (and thus preventing attachment) as the most likely mechanism through which turfs impaired *H. banksii* recruitment, but also acknowledged the potential roles of sediment accumulation (Airoldi, 2003; Alestra et al., 2014; Schiel et al., 2006) alleopathy (Suzuki et al., 1998) and sloughing of epithallus layers (Littler and Littler, 1999).

A number of studies have demonstrated variable recovery of *H. banksii* in Australia and New Zealand and at different spatial scales, suggesting different mechanistic processes. Cover of *H. banksii* was reduced from ~ 60 % to < 5 % on a Sydney shore by a severe

1 storm in 1974, but had recovered to ~ 60 % cover by 1979 (Underwood, 1999).
2
3 However, recovery was highly variable even on small scales and was dependent on the
4
5 amount of damage done to holdfasts and the activities of grazers: recovery was much
6
7 slower when holdfasts were removed than when they were undamaged; grazing
8
9 molluscs and seastars slowed the rate of recovery but did not prevent it altogether
10
11 (Underwood, 1998).
12
13
14
15

16 In an experimental test of the effects of trampling on *H. banksii* at two sites separated
17
18 by 320 km (Kaikoura and Moeraki) in southern New Zealand, despite almost 100 %
19
20 reduction in cover in some treatments, *H. banksii* recovered to control levels at both
21
22 sites in all treatments except one at Moeraki (where final cover was ~ 25 % less than
23
24 controls) within 21 months of the original pulse disturbance (Schiel and Taylor, 1999).
25
26 However, larger scale (3 × 3 m plots) experimental canopy removals of *H. banksii* at the
27
28 same sites resulted in very different patterns of recovery between sites (Lilley and
29
30 Schiel, 2006). Plots had recovered to control levels (almost 100 % cover) at Kaikoura
31
32 after 2 y, but at Moeraki cover was only ~ 20 % (c.f. ~ 100 % in controls) after 2 y.
33
34 Furthermore, although *H. banksii* had recovered to form almost a closed canopy similar
35
36 in cover to control plots at both sites 8 y after initial canopy removals, density and
37
38 biomass of *H. banksii* fronds was significantly greater in 0.25 m² quadrats at Kaikoura
39
40 than Moeraki, and control plots always had higher frond densities and biomass than
41
42 pulse and press plots (Schiel and Lilley, 2011). It is worth noting that where community
43
44 data were recorded for the New Zealand example, the communities in removal plots
45
46 were still very different from those in controls (as in Hawkins and Harkin, 1985; Lilley
47
48 and Schiel, 2006; Schiel and Lilley, 2011) even when canopy recovery was complete.
49
50
51 This suggests that the residence time (Bruno and Bertness, 2001; Jones et al., 1994,
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1997; Konar, 2000) might be particularly important for benthic communities that rely on recruitment (and associated variability) of component species.

In Victoria, Australia, canopy recovery in response to different levels of trampling was also variable, where high trampling plots only returned to 50 % of the initial cover after 420 days (Povey and Keough, 1991). Sites subject to heavy trampling may never recover fully because of annual pulses of trampling pressure (Povey and Keough, 1991).

When these shores were examined over longer temporal scales, large site-to-site differences in recovery were seen for trampling experiments at sites separated by only 1 km (Keough and Quinn, 1998). Two sites on the same large rock platform showed a series of pulse responses to repeated pulse disturbances, but a third site 1 km away showed a press response to repeated pulse disturbances over a 5 y period. At the end of the 5 y experiment the two adjacent sites had comparable canopy cover in all treatments, but the third site cover was < 10 % cover in high trampling treatments and 60 – 70 % in intermediate trampling treatments (c.f. > 90 % in controls) (Keough and Quinn, 1998).

We do not currently fully understand the causes for these large differences in recovery between sites (Keough and Quinn, 1998; Lilley and Schiel, 2006), although there is evidence that differences (of 0.4 m) in height on the shore can affect recovery rates (Schiel, 2006). However, it is clear that monitoring programs focusing on *H. banksii* (and potentially other furoid algae), including those designed to monitor restoration success, must account for the potential variability between sites in recovery of this species, by using well-designed monitoring programs with multiple control/impact sites and replication at meaningful temporal and spatial scales (Chapman et al., 1995; Keough and Quinn, 1998).

7 Discrepancies in knowledge from experimental field ecology and application by restoration ecologists

Despite the important role that habitat-forming species play in structuring communities and ecosystems, ecological research has focused more on negative interactions (*e.g.* competition and predation) than positive interactions between species (Bertness and Leonard, 1997; Bertness *et al.*, 1999; Bruno and Bertness, 2001; Jones *et al.*, 1994). This is also true for intertidal rocky shores that have been the focus of a plethora of experimental studies and development of current ecological theory. Bruno and Bertness (2001) surveyed publications in the prominent marine ecological journal, *Marine Ecology Progress Series*, from 1994 – 1996 and found a 5:1 ratio of papers examining predation compared to ecosystem engineers (facilitators) (Table 1). They made a plea for more research to help us understand the strength and importance of interactions between ecosystem engineers and associated species. However, 20 years on and the focus on the roles of ecosystem engineers has declined. A survey of the papers in *Marine Ecology Progress Series* from 2003 – 2005 showed only 1.5% of papers focussing on the roles of ecosystem engineers and then ten years later from 2013 – 2015, the percentage of papers dropped to 0.5%. The predation: ecosystem engineer ratio in the later years has changed to an approximately 11:1 (Table 1). The picture is not much different if we look at broader ecological journals, such as *Ecology*, *Oikos* and *Oecologia*, where the percentages of papers focusing on ecosystem engineering from 2003 – 2005 were 4.1 %, 2.5 % and 2.5 %, respectively. However, a decade later (2013 – 2015) the percentages have slightly increased (5.5 %, 5.9 % and 4.7 %, respectively). In contrast, 22 % of papers published in *Restoration Ecology* from 2003 – 2005 focused on restoring ecosystem engineers to facilitate restoration of associated assemblages, but

overtime this trend has dramatically decreased to 1.8 % from 2013 – 2015, with a shift in emphasis to studies focusing on sampling techniques for restoration (Table 2). Additionally, Byers *et al.* (2006) proposed that incorporating ecosystem engineers into restoration efforts may result in cheaper, easier, faster and more sustainable restoration. This suggests that managers and restoration ecologists have to make decisions about best practice for biogenic habitat restoration, without necessarily a sound understanding of the interactions between species and the conditions in which these interactions may be strongest.

It is also important to note that restoration of ecosystem engineers, regardless of how much is known of the positive and negative interactions between species, is unlikely to be successful unless the physico-chemical environment is conducive to sustaining the restored populations (Nienhuis *et al.*, 2002). Less than 10 % of studies published in *Restoration Ecology* since 2003 have focussed on restoring the physico-chemical environment, but this exceeds those focusing on the restoration of biotic habitats in recent years (Table 2).

8 Restoration and future directions

For sound management of our coastal ecosystems we must understand the role of individual species and their interactions within communities (Marzinelli *et al.*, 2014; Schiel, 2006; Wahl *et al.*, 2015). Management decisions to focus on the re-establishment of important ecosystem engineers such as canopy-forming fucoid algae, should result in effective restoration of rocky intertidal ecosystems (Benedetti-Cecchi *et al.*, 2001; Bruno and Bertness, 2001; Byers *et al.*, 2006; Crain and Bertness, 2006; Jones *et al.*, 1994, 1997; Nienhuis *et al.*, 2002; Wahl *et al.*, 2015; Wright and Jones, 2006). In

temperate Australasia, *H. banksii* has many characteristics that suggest it is an important ecosystem engineer: it forms extensive, long-lived and stable canopies; it integrates and may modulate many abiotic resources (slows flow, decreases heat and desiccation, decreases UV radiation, provides microhabitats) in the physically and physiologically stressful environment of the rocky intertidal zone; has direct and indirect, positive and negative interactions with a wide range of species and undoubtedly increases biodiversity on seascape scales (Jones et al., 1994, 1997). However, for a full understanding of the ecosystem function of *H. banksii* and other furoid algae, it will be important to conduct carefully planned manipulative experiments to assess the types and strength of interactions with associated species (Díez et al., 2013; Jones et al., 1997; Menge et al., 1994), the influence of the quality of canopy cover and the environmental conditions under which these interactions vary (Bertness et al., 1999; Bruno and Bertness, 2001; Crain and Bertness, 2006; Jones et al., 1997; Wright and Jones, 2006).

Additionally, to facilitate effective restoration of canopy-forming furoid algae and their associated communities affected by anthropogenic disturbances such as sewage effluent discharge, we must first have a thorough understanding of the factors that lead to the initial demise of these populations at polluted sites (as suggested in Wahl et al., 2015), including effects on early life-history stages as well as established plants (e.g. Sales et al., 2011), the spatial and temporal scales at which the impacts can be demonstrated and the processes that operate at these scales (Benedetti-Cecchi et al., 2001). Only then can we set realistic water quality targets for restoration efforts.

Finally, for restoration of both furoid engineers and associated species we must understand the connectivity and the patterns of gene flow between populations

(Coleman et al., 2011; McKenzie and Bellgrove, 2006, 2008, 2009; Schiel, 2004), and the variability in, and factors that may limit, the associated processes. This will allow us to make informed predictions about the probability of natural re-establishment and the conditions under which it might be maximised. Finally, in order to assess the success of intertidal restoration projects, we need to carefully design monitoring programmes that are replicated over appropriate spatial and temporal scales to detect positive or negative effects of restoration over and above the high degree of variability that naturally exists in these systems (Chapman et al., 1995; Keough and Quinn, 1998; Schiel, 2006).

Acknowledgements

This review was funded by the Environment Protection Authority Victoria (EPA Victoria) and the School of Life & Environmental Sciences, Deakin University. B. Light (of EPA Victoria) is thanked for early discussions in framing the review, but the EPA Victoria had no role in the data collection, analysis or writing of this review. J. Sherwood and J. Mondon are thanked for discussions relating to toxicity tests and B. Robson and S. Lilley for reviewing an early draft. The manuscript was improved by comments from anonymous referees. Authors are listed in order of contribution to this review.

References

Adams, M.S., Stauber, J.L., Binet, M.T., Molloy, R., Gregory, D., 2008. Toxicity of a secondary-treated sewage effluent to marine biota in Bass Strait, Australia: Development of action trigger values for a toxicity monitoring program. *Mar. Pollut. Bull.* 57, 587-598.

1 Airoidi, L., 2003. The effects of sedimentation on rocky coast assemblages. *Oceanogr*
2
3
4 *Mar Biol* 41, 161-236.

5
6
7 Alestra, T., Tait, L.W., Schiel, D.R., 2014. Effects of algal turfs and sediment
8
9 accumulation on replenishment and primary productivity of furoid assemblages. *Mar.*
10
11 *Ecol. Prog. Ser.* 511, 59-70, doi: 10.3354/Meps10932.

12
13
14
15 Altieri, A.H., van de Koppel, J., 2014. Foundation species in marine ecosystems, in:
16
17 Bertness, M.D., Bruno, J.F., Silliman, B.R., Stachowicz, J.J. (Eds.), *Marine community*
18
19 *ecology and conservation*. Sinauer Associates, Inc, Sunderland, Massachusetts, USA, pp.
20
21
22 37-56.

23
24
25
26 Araujo, R., Vaselli, S., Almeida, M., Serrao, E., Sousa-Pinto, I., 2009. Effects of
27
28 disturbance on marginal populations: human trampling on *Ascophyllum nodosum*
29
30 assemblages at its southern distribution limit. *Mar Ecol Prog Ser* 378, 81-92, doi:
31
32 10.3354/meps07814.

33
34
35
36
37 Batley, G.E., Simpson, S.L., 2009. Development of guidelines for ammonia in estuarine
38
39 and marine water systems. *Mar. Pollut. Bull.* 58, 1472-1476, doi:
40
41 10.1016/j.marpolbul.2009.06.005.

42
43
44
45 Beermann, A.J., Ellrich, J.A., Molis, M., Scrosati, R.A., 2013. Effects of seaweed
46
47 canopies and adult barnacles on barnacle recruitment: The interplay of positive and
48
49 negative influences. *J. Exp. Mar. Biol. Ecol.* 448, 162-170, doi:
50
51 10.1016/j.jembe.2013.07.001.

52
53
54
55
56 Bellgrove, A., 1998. Recruitment of intertidal macroalgae on a wave-exposed rocky
57
58 coast. Monash University, Clayton, Australia, p. 130.

1 Bellgrove, A., Clayton, M.N., Quinn, G.P., 1997. Effects of secondarily treated sewage
2
3 effluent on intertidal macroalgal recruitment processes. *Mar Freshw Res* 48, 137-146.
4
5

6
7 Bellgrove, A., Clayton, M.N., Quinn, G.P., 2004. An integrated study of the temporal
8
9 and spatial variation in the supply of propagules, recruitment and assemblages of
10
11 intertidal macroalgae on a wave-exposed rocky coast, Victoria, Australia. *J. Exp. Mar.*
12
13 *Biol. Ecol.* 310, 207-225.
14
15

16
17 Bellgrove, A., McKenzie, P.F., McKenzie, J.L., Sfilgoj, B., 2010. Restoration of the
18
19 habitat-forming furoid alga *Hormosira banksii* at effluent-affected sites: competitive
20
21 exclusion by coralline turfs. *Mar. Ecol. Prog. Ser.* 419, 47-56.
22
23
24

25
26 Benedetti-Cecchi, L., Cinelli, F., 1992. Canopy removal experiments in *Cystoseira*-
27
28 dominated rockpools from the Western coast of the Mediterranean (Ligurian Sea). *J.*
29
30 *Exp. Mar. Biol. Ecol.* 155, 69-83.
31
32

33
34 Benedetti-Cecchi, L., Pannacciulli, F., Bulleri, F., Moschella, P.S., Airolidi, L., Relini,
35
36 G., Cinelli, F., 2001. Predicting the consequences of anthropogenic disturbance: large-
37
38 scale effects of loss of canopy algae on rocky shores. *Mar. Ecol. Prog. Ser.* 214, 137-
39
40 150.
41
42
43

44
45 Bertness, M.D., Leonard, G.H., 1997. The role of positive interactions in communities:
46
47 lessons from intertidal habitats. *Ecology* 78, 1976-1989.
48
49

50
51 Bertness, M.D., Leonard, G.H., Levine, J.M., Schmidt, P.R., Ingraham, A.O., 1999.
52
53 Testing the relative contribution of positive and negative interactions in rocky intertidal
54
55 communities. *Ecology* 80, 2711-2726.
56
57
58
59
60
61
62
63
64
65

Bertocci, I., Seabra, M.I., Dominguez, R., Jacinto, D., Ramirez, R., Coca, J., Tuya, F.,
2014. Effects of loss of algal canopies along temperature and irradiation gradients in
continental Portugal and the Canary Islands. *Mar. Ecol. Prog. Ser.* 506, 47-U355, doi:
10.3354/meps10785.

Bishop, M.J., Byers, J.E., Marcek, B.J., Gribben, P.E., 2012. Density-dependent
facilitation cascades determine epifaunal community structure in temperate Australian
mangroves. *Ecology* 93, 1388-1401.

Bishop, M.J., Fraser, J., Gribben, P.E., 2013. Morphological traits and density of
foundation species modulate a facilitation cascade in Australian mangroves. *Ecology* 94,
1927-1936, doi: 10.1890/12-1847.1.

Bishop, M.J., Morgan, T., Coleman, M.A., Kelaher, B.P., Hardstaff, L.K., Evenden,
R.W., 2009. Facilitation of molluscan assemblages in mangroves by the fuclean alga
Hormosira banksii. *Marine Ecology Progress Series* 392, 111-122.

Blanchette, C.A., 1997. Size and survival of intertidal plants in response to wave action:
A case study with *Fucus gardneri*. *Ecology* 78, 1563-1578.

Bokn, T.L., Duarte, C.M., Pedersen, M.F., Marba, N., Moy, F.E., Barron, C., Bjerkeng,
B., Borum, J., Christie, H., Engelbert, S., Fotel, F.L., Hoell, E.E., Karez, R., Kersting,
K., Kraufvelin, P., Lindblad, C., Olsen, M., Sanderud, K.A., Sommer, U., Sorensen, K.,
2003. The response of experimental rocky shore communities to nutrient additions.
Ecosystems 6, 577-594.

1 Bordeyne, F., Migne, A., Davoult, D., 2015. Metabolic activity of intertidal *Fucus* spp.
2 communities: evidence for high aerial carbon fluxes displaying seasonal variability. *Mar.*
3 *Biol.* 162, 2119-2129, doi: 10.1007/s00227-015-2741-6.
4
5

6
7
8
9 Borowitzka, M.A., 1972. Intertidal algal species diversity and the effect of pollution.
10 *Aust J Mar Freshwater Res* 23, 73-84.
11
12

13
14
15 Brawley, S.H., Johnson, L.E., 1991. Survival of furoid embryos in the intertidal zone
16 depend upon developmental stage and microhabitat. *J. Phycol.* 27, 179-186.
17
18

19
20
21 Brawley, S.H., Johnson, L.E., 1993. Predicting desiccation stress in microscopic
22 organisms: the use of agarose beads to determine evaporation within and between
23 intertidal microhabitats. *J. Phycol.* 29, 528-535.
24
25

26
27
28
29 Brown, V., Ducker, S.C., Rowan, K.S., 1977. The effect of orthophosphate
30 concentration on the growth of articulated coralline algae (Rhodophyta). *Phycologia* 16,
31 125-131.
32
33

34
35
36
37
38 Brown, V.B., Davies, S.A., Synnot, R.N., 1990. Long-term monitoring of the effects of
39 treated sewage effluent on the intertidal macroalgae community near Cape Schanck,
40 Victoria, Australia. *Bot. Mar.* 33, 85-98.
41
42

43
44
45
46 Bruno, J., Bertness, M.D., 2001. Habitat modification and facilitation in benthic marine
47 communities, in: Bertness, M.D., Gaines, S.D., Hay, M.E. (Eds.), *Marine Community*
48 *Ecology*. Sinauer Associates, Inc., Sunderland, Massachusetts, pp. 201-218.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 Bulleri, F., Benedetti-Cecchi, L., Acunto, S., Cinelli, F., Hawkins, S.J., 2002. The
2 influence of canopy algae on vertical patterns of distribution of low-shore assemblages
3 on rocky coasts in the northwest Mediterranean. *J. Exp. Mar. Biol. Ecol.* 267, 89-106.
4
5
6
7
8
9 Buschmann, A.H., Bravo, A., 1990. Intertidal amphipods as potential dispersal agents of
10 carpospores of *Irideae laminarioides* (Gigartinales, Rhodophyta). *J. Phycol.* 26, 417-
11 420.
12
13
14
15
16
17 Buschmann, A.H., Vergara, P.A., 1993. Effect of rocky intertidal amphipods on algal
18 recruitment: a field study. *J. Phycol.* 29, 154-159.
19
20
21
22
23 Bustamante, M., Bevilacqua, S., Tajadura, J., Terlizzi, A., Saiz-Salinas, J.I., 2012.
24 Detecting human mitigation intervention: Effects of sewage treatment upgrade on rocky
25 macrofaunal assemblages. *Mar. Environ. Res.* 80, 27-37, doi:
26 10.1016/j.marenvres.2012.06.007.
27
28
29
30
31
32
33
34 Bustamante, R.H., Branch, G.M., Eekhout, S., 1995. Maintenance of an exceptional
35 intertidal grazer biomass in South Africa: subsidy by subtidal kelps. *Ecology* 76, 2314-
36 2329.
37
38
39
40
41
42 Byers, J.E., Cuddington, K., Jones, C.G., Talley, T.S., Hastings, A., Lambrinos, J.G.,
43 Crooks, J.A., Wilson, W.G., 2006. Using ecosystem engineers to restore ecological
44 systems. *Trends Ecol. Evol.* 21, 493-500.
45
46
47
48
49
50
51 Campbell, A.H., Marzinelli, E.M., Verges, A., Coleman, M.A., Steinberg, P.D., 2014.
52 Towards Restoration of Missing Underwater Forests. *PLoS ONE* 9, e84106, doi:
53 10.1371/journal.pone.0084106.
54
55
56
57
58
59
60
61
62
63
64
65

1 Chapman, A.S., Fletcher, R.L., 2002. Differential effects of sediments on survival and
2 growth of *Fucus serratus* embryos (Fucales, Phaeophyceae). J. Phycol. 38, 894-903.
3
4

5
6
7 Chapman, M.G., Underwood, A.J., Skilleter, G.A., 1995. Variability at different spatial
8 scales between a subtidal assemblage exposed to the discharge of sewage and two
9 control assemblages. J. Exp. Mar. Biol. Ecol. 189, 103-122.
10
11
12

13
14
15 Clarke, S.M., Womersley, H.B.S., 1981. Cross-fertilization and Hybrid Development of
16 forms of the Brown Alga *Hormosira banksii* (Turner) Decaisne. Aust. J. Bot. 29, 497-
17 505.
18
19
20
21

22
23 Coleman, M.A., Chambers, J., Knott, N.A., Malcolm, H.A., Harasti, D., Jordan, A.,
24 Kelaher, B.P., 2011. Connectivity within and among a network of temperate marine
25 reserves. PLoS ONE 6, 1-9.
26
27
28
29
30

31
32 Coleman, M.A., Kelaher, B.P., Steinberg, P.D., Millar, A.J.K., 2008. Absence of a large
33 brown macroalga on urbanized rocky reefs around Sydney, Australia, and evidence for
34 historical decline. J. Phycol. 44, 897-901, 10.1111/j.1529-8817.2008.00541.x.
35
36
37
38
39

40 Cormaci, M., Furnari, G., 1999. Changes of the benthic algal flora of the Tremiti Islands
41 (southern Adriatic) Italy. Hydrobiologia 398/399, 75-79.
42
43
44
45

46 Cowen, R.K., Lwiza, K.M.M., Sponaugle, S., Paris, C.B., Olson, D.B., 2000.
47 Connectivity of marine populations: open or closed? Science 287, 857-859.
48
49
50

51
52 Crain, C.M., Bertness, M.D., 2006. Ecosystem engineering across environmental
53 gradients: implications for conservation and management. Bioscience 56, 211-218.
54
55
56
57
58
59
60
61
62
63
64
65

1 Crowe, T.P., Frost, N.J., Hawkins, S.J., 2011a. Interactive effects of losing key grazers
2 and ecosystem engineers vary with environmental context. *Mar. Ecol. Prog. Ser.* 430,
3
4 223-234, doi: 10.3354/meps09023.
5
6

7
8
9 Crowe, T.P., Frost, N.J., Hawkins, S.J., 2011b. Interactive effects of losing key grazers
10 and ecosystem engineers vary with environmental context. *Mar Ecol Prog Ser* 430, 223-
11
12 234, doi: 10.3354/meps09023.
13
14
15

16
17 Davenport, J., Butler, A., Cheshire, A., 1999. Epifaunal composition and fractal
18 dimensions of marine plants in relation to emersion. *J. Mar. Biol. Assoc. U.K.* 79, 351-
19
20 355.
21
22
23

24
25 Dayton, P.K., 1971. Competition, Disturbance, and Community Organization: The
26 Provision and Subsequent Utilization of Space in a Rocky Intertidal Community. *Ecol.*
27
28 *Monogr.* 41, 351-389.
29
30
31

32
33 Dayton, P.K., 1972. Toward an understanding of community resilience and the potential
34 effects of enrichment to the benthos as McMurdo Sound, Antarctica, *Proceedings of the*
35
36 *Colloquium on Conservation problems in Antarctica*, pp. 81-96.
37
38
39

40
41 de-la-Ossa-Carretero, J.A., Del-Pilar-Ruso, Y., Giménez-Casalduero, F., Sánchez-
42
43 Lizaso, J.L., 2016. Monitoring the effects of wastewater treatment strategies. *Environ.*
44
45 *Monit. Assess.* 188, 110, doi: 10.1007/s10661-016-5092-y.
46
47
48

49
50 Denny, M., Brown, V., Carrington, E., Kraemer, G., Miller, A., 1989. Fracture
51 mechanics and the survival of wave-swept macroalgae. *J. Exp. Mar. Biol. Ecol.* 127,
52
53 211-228.
54
55
56
57
58
59
60
61
62
63
64
65

1 Deysher, L.E., Dean, T.A., Grove, R.S., Jahn, A., 2002. Design considerations for an
2 artificial reef to grow giant kelp (*Macrocystis pyrifera*) in Southern California. *ICES J.*
3
4
5
6 *Mar. Sci.* 59, S201-S207.

7
8
9 Díez, I., Santolaria, A., Muguerza, N., Gorostiaga, J.M., 2013. Measuring restoration in
10 intertidal macrophyte assemblages following sewage treatment upgrade. *Mar. Environ.*
11
12
13 *Res.* 84, 31-42, doi: 10.1016/j.marenvres.2012.11.006.

14
15
16
17 Díez, I., Santolaria, A., Secilla, A., Maria Gorostiaga, J., 2009. Recovery stages over
18 long-term monitoring of the intertidal vegetation in the 'Abra de Bilbao' area and on the
19
20
21 adjacent coast (N. Spain). *Eur J Phycol* 44, 1-14, Pii 908501138, doi:
22
23
24 10.1080/09670260802158642.

25
26
27
28 Estes, J.A., Palmisano, J., 1974. Sea otters: their role in structuring nearshore
29
30
31 communities. *Science* 185, 1058-1060.

32
33
34 Fairweather, P.G., 1990. Sewage and the biota on seashores: assessment of impact in
35
36
37 relation to natural variability. *Environ. Monit. Assess.* 14, 197-210.

38
39
40 Fredriksen, S., Christie, H., Saethre, B.A., 2005. Species richness in macroalgae and
41
42
43 macrofauna assemblages on *Fucus serratus* L. (Phaeophyceae) and *Zostera marina* L.
44
45 (Angiospermae) in Skagerrak, Norway. *Mar. Biol. Res.* 1, 2-19.

46
47
48 Gagnon, K., McKindsey, C.W., Johnson, L.E., 2015. Roles of dispersal mode, recipient
49
50
51 environment and disturbance in the secondary spread of the invasive seaweed *Codium*
52
53
54 *fragile*. *Biol. Invasions* 17, 1123-1136, doi: 10.1007/s10530-014-0782-2.

- Gollty, C., Migne, A., Davoult, D., 2008. Benthic metabolism on a sheltered rocky shore: Role of the canopy in the carbon budget. *J. Phycol.* 44, 1146-1153, doi: 10.1111/j.1529-8817.2008.00569.x.
- Golubic, S., 1970. Effect of organic pollution on benthic communities. *Mar. Pollut. Bull.* 1, 56-57.
- Gosselin, L.A., Chia, F., 1995. Distribution and dispersal of early juvenile snails: effectiveness of intertidal microhabitats as refuges and food sources. *Mar. Ecol. Prog. Ser.* 128, 213-223.
- Grant, W.S., 1977. High intertidal community organization on a rocky headland in Maine, USA. *Mar. Biol.* 44, 15-25.
- Griffiths, C.L., Stanton-Dozey, J., Koop, K., 1983. Kelp wrack and flow of energy through a sandy beach ecosystem., in: McLachlan, A., Erasmus, T. (Eds.), *Sandy Beaches as Ecosystems*. Dr W Junk, The Hague, pp. 547-556.
- Halpern, B.S., Silliman, B.R., Olden, J.D., Bruno, J.P., Bertness, M.D., 2007. Incorporating positive interactions in aquatic restoration and conservation. *Front. Ecol. Environ.* 5, 153-160, doi: 10.1890/1540-9295(2007)5[153:Ipiaar]2.0.Co;2.
- Hawkins, S.J., Harkin, E., 1985. Preliminary canopy removal experiments in algal dominated communities low on the shore and in the shallow subtidal on the Isle of Man. *Bot. Mar.* 28, 223-230.
- Hay, M.E., 1981. The functional morphology of turf-forming seaweeds: persistence in stressful marine habitats. *Ecology* 62, 739-750.

Hernandez-Carmona, G., Garcia, O., Robledo, D., Foster, M., 2000. Restoration techniques for *Macrocystis pyrifera* (Phaeophyceae) populations at the southern limit of their distribution in Mexico. *Bot. Mar.* 43, 273-284, doi: 10.1515/Bot.2000.029.

Hernández-Carmona, G., Hughes, B., Graham, M.H., 2006. Reproductive longevity of drifting kelp *Macrocystis pyrifera* (Phaeophyceae) in Monterey Bay, USA. *J Phycol* 42, 1199-1207.

Hily, C., Jean, F., 1997. Macrobenthic biodiversity in intertidal habitats of the Iroise biosphere reserve (Brittany, France). *J. Mar. Biol. Assoc. U.K.* 77, 311-323.

Hirose, H., 1978. Composition of benthic marine algae in relation to pollution in the Seto Inland Sea, Japan. *Proceedings of the International Seaweed Symposium* 9, 173-179.

Jenkins, S.R., Hawkins, S.J., Norton, T.A., 1999. Direct and indirect effects of a macroalgal canopy and limpet grazing in structuring a sheltered inter-tidal community. *Mar. Ecol. Prog. Ser.* 188, 81-92.

Jenkins, S.R., Norton, T.A., Hawkins, S.J., 2004. Long term effects of *Ascophyllum nodosum* canopy removal on mid shore community structure. *J. Mar. Biol. Assoc. U.K.* 84, 327-329.

Jernakoff, P., 1986. Experimental investigation of interactions between the perennial red alga *Gelidium pusillum* and barnacles on a New South Wales rocky shore. *Mar. Ecol. Prog. Ser.* 28, 259-263.

1 Johnson, L.E., Brawley, S.H., 1998. Dispersal and recruitment of a canopy-forming
2
3 intertidal alga: the relative roles of propagule availability and post-settlement processes.
4
5
6 *Oecologia* 117, 517-526.

7
8
9 Jones, C.G., Lawton, J.H., Shachak, M., 1994. Organisms as ecosystem engineers.
10
11
12 *Oikos* 69, 373-386.

13
14
15 Jones, C.G., Lawton, J.H., Shachak, M., 1997. Positive and negative effects of
16
17 organisms as physical ecosystem engineers. *Ecology* 78, 1946-1957.

18
19
20
21 Kain (Jones), J.M., 2015. *Hormosira banksii* (Phaeophyceae): a tough
22
23
24 survivor in the harsh conditions of high intertidal pools in southeast Australia. *Eur. J.*
25
26
27 *Phycol.* 50, 408-421, doi: 10.1080/09670262.2015.1075594.

28
29
30 Kelaher, B.P., Chapman, M.G., Underwood, A.J., 2001. Spatial patterns of diverse
31
32 macrofaunal assemblages in coralline turf and their associations with environmental
33
34 variables. *J. Mar. Biol. Assoc. U.K.* 81, 917-930.

35
36
37
38 Keough, M.J., Quinn, G.P., 1998. Effects of periodic disturbances from trampling on
39
40
41 rocky intertidal algal beds. *Ecol. Appl.* 8, 141-161.

42
43
44 Kevekordes, K., 2001. Toxicity tests using developmental stages of *Hormosira banksii*
45
46 (Phaeophyta) identify ammonium as a damaging component of secondary treated
47
48 sewage effluent discharged into Bass Strait, Victoria, Australia. *Mar. Ecol. Prog. Ser.*
49
50
51 219, 139-148.

52
53
54
55 King, A., 1992. Human activity and effects on marine intertidal plant and animal
56
57
58 populations. University of Melbourne, Melbourne.

59
60
61
62
63
64
65

- Kinlan, B.P., Gaines, S.D., 2003. Propagule dispersal in marine and terrestrial environments: a community perspective. *Ecology* 84, 2007-2020.
- Klavestad, N., 1978. The marine algae of the polluted inner part of the Oslofjord. *Bot. Mar.* 21, 71-97.
- Konar, B., 2000. Limited effects of a keystone species: trends of sea otters and kelp forests at the Semichi Islands, Alaska. *Mar. Ecol. Prog. Ser.* 199, 271-280.
- Korpinen, S., Jormalainen, V., 2008. Grazing and nutrients reduce recruitment success of *Fucus vesiculosus* L. (Fucales : Phaeophyceae). *Estuarine Coastal Shelf Sci* 78, 437-444.
- Leonard, G.H., 1999. Positive and negative effects of intertidal algal canopies on recruitment and survival of barnacles. *Mar. Ecol. Prog. Ser.* 178, 241-249.
- Lilley, S.A., Schiel, D.R., 2006. Community effects following the deletion of a habitat-forming alga from rocky marine shores. *Oecologia* 148, 672-681.
- Littler, M.M., Littler, D.S., 1999. Epithallus sloughing: a self-cleaning mechanism for coralline algae. *Coral Reefs* 18, 204.
- Littler, M.M., Murray, S.N., 1975. Impact of sewage on the distribution, abundance and community structure of rocky intertidal macro-organisms. *Mar. Biol.* 30, 277-291.
- Littler, M.M., Murray, S.N., 1978. Influence of domestic wastes on energetic pathways in rocky intertidal communities. *J. Appl. Ecol.* 15, 583-595.

Macaya, E.C., Boltaña, S., Hinojosa, I.A., Macchiavello, J.E., Valdivia, N.A., Vásquez, N.R., Buschmann, A.H., Vásquez, J.A., Alonso Vega, J.M., Thiel, M., 2005. Presence of sporophylls in floating kelp rafts of *Macrocystis* spp. (Phaeophyceae) along the Chilean Pacific Coast. *J Phycol* 41, 913-922.

Manning, P.F., 1979. The biological effects of the discharge of secondarily treated sewage effluent at Boags Rocks, Gunnamatta Beach, Victoria. La Trobe University, Bundoora, Australia.

Marzinelli, E.M., Burrows, M.T., Jackson, A.C., Mayer-Pinto, M., 2012. Positive and Negative Effects of Habitat-Forming Algae on Survival, Growth and Intra-Specific Competition of Limpets. *PLoS ONE* 7.

Marzinelli, E.M., Campbell, A.H., Verges, A., Coleman, M.A., Kelaher, B.P., Steinberg, P.D., 2014. Restoring seaweeds: does the declining furoid *Phyllospora comosa* support different biodiversity than other habitats? *J. Appl. Phycol.* 26, 1089-1096, doi: 10.1007/s10811-013-0158-5.

Marzinelli, E.M., Leong, M.R., Campbell, A.H., Steinberg, P.D., Verges, A., 2016. Does restoration of a habitat-forming seaweed restore associated faunal diversity? *Restor. Ecol.* 24, 81-90, doi: 10.1111/rec.12292.

May, V., 1985. Observations on algal floras close to two sewerage outlets. *Cunninghamia* 1, 385-394.

Mayer-Pinto, M., Ignacio, B.L., 2015. Effects of chemical disturbances on intertidal benthic assemblages. *Sci. Total Environ.* 506, 10-17.

1 Mayer-Pinto, M., Matias, M.G., Coleman, R.A., 2016. The interplay between habitat
2 structure and chemical contaminants on biotic responses of benthic organisms. *Peerj* 4.
3
4

5
6
7 McCook, L.J., Chapman, A.R.O., 1991. Community succession following massive ice-
8 scour on an exposed rocky shore: effects of *Fucus* canopy algae and mussels during late
9 succession. *J. Exp. Mar. Biol. Ecol.* 154, 131-169.
10
11
12

13
14
15 McKay, J.K., Christian, C.E., Harrison, S., Rice, K.J., 2005. "How local is local?" - A
16 review of practical and conceptual issues in the genetics restoration. *Restor. Ecol.* 13,
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

432-440.
McKenzie, P.F., Bellgrove, A., 2006. No outbreeding depression at a regional scale for
a habitat-forming intertidal alga with limited dispersal. *Mar Freshw Res* 57, 655-663.

McKenzie, P.F., Bellgrove, A., 2008. Dispersal of *Hormosira banksii* (Phaeophyceae)
via detached fragments: reproductive viability and longevity. *J. Phycol.* 44, 1108-1115.

McKenzie, P.F., Bellgrove, A., 2009. Dislodgment and attachment strength of the
intertidal macroalga, *Hormosira banksii* (Fucales, Phaeophyceae). *Phycologia* 48, 335-
343.

Menge, B.A., 1995. Indirect effects in marine rocky intertidal interaction webs: patterns
and importance. *Ecol. Monogr.* 65, 21-74.

Menge, B.A., Berlow, E.L., Blanchette, C.A., Navarrete, S.A., Yamada, S.B., 1994. The
keystone species concept: variation in interaction strength in a rocky intertidal habitat.
Ecol. Monogr. 64, 249-286.

1 Migné, A., Gollety, C., Davoult, D., 2015. Effect of canopy removal on a rocky shore
2 community metabolism and structure. *Mar. Biol.* 162, 449-457, doi: 10.1007/s00227-
3 014-2592-6.
4
5
6

7
8
9 Molloy, R., Fox, D., Stauber, J., Beardall, J., 2004. Eastern Treatment Plant: Monitoring
10 and Receiving Environment. Commonwealth Scientific and Industrial Research
11 Organisation, Melbourne, p. 64.
12
13
14
15

16
17 Molloy, R., Robinson, L., Stauber, J., Fox, D., Beardall, J., 2007. Eastern Treatment
18 Plant: Monitoring the Receiving Environment. 2004-2007 Monitoring Report.
19
20 Commonwealth Scientific and Industrial Research Organization, Melbourne.
21
22
23
24

25
26 Myers, J.H., Duda, S., Gunthorpe, L., Allinson, G., 2006. Assessing the performance of
27 *Hormosira banksii* (Turner) Desicaine germination and growth assay using four
28 reference toxicants. *Ecotoxicol. Environ. Saf.* 64, 304-311, doi:
29 10.1016/j.ecoenv.2005.04.013.
30
31
32
33
34
35

36
37 Nienhuis, P.H., Bakker, J.P., Grootjans, A.P., Gulati, R.D., de Jonge, V.N., 2002. The
38 state of the art of aquatic and semi-aquatic ecological restoration projects in the
39 Netherlands. *Hydrobiologia* 478, 219-233.
40
41
42
43

44
45 Osborn, J.E.M., 1948. The structure and life history of *Hormosira banksii* (Turner)
46 Decaisne. *Trans R Soc NZ* 77, 47-71.
47
48
49

50
51 Paine, R.T., 1966. Food web complexity and species diversity. *Am. Nat.* 100, 65-75.
52
53

54
55 Paine, R.T., 1969a. A note on trophic complexity and community stability. *Am. Nat.*
56 103, 91-93.
57
58
59
60
61
62
63
64
65

- Paine, R.T., 1969b. The *Pisaster-Tegula* interaction: prey patches, predator food preference, and intertidal community structure. *Ecology* 50, 950-961.
- Petraitis, P.S., Latham, R.E., 1999. The importance of scale in testing the origins of alternative community states. *Ecology* 80, 429-442.
- Pocklington, J.B., 2003. Positive interactions on rocky seashores: are they important, or can they be ignored? Flinders University, Bedford Park, Australia.
- Pocklington, J.B., Jenkins, S.R., Bellgrove, A., Keough, M.J., O'Hara, T.D., Masterson-Algar, P.E., Hawkins, S.J., 2017. Disturbance alters ecosystem engineering by a canopy-forming alga. *J. Mar. Biol. Assoc. U.K.*, doi: 10.1017/S0025315416002009.
- Povey, A., Keough, M.J., 1991. Effects of trampling on plant and animal populations on rocky shores. *Oikos* 61, 355-368.
- Ralph, P.J., Morrison, D.A., Addison, A., 1998. A quantitative study of the patterns of morphological variation within *Hormosira banksii* (Turner) Decaisne (Fucales : Phaeophyta) in south-eastern Australia. *J. Exp. Mar. Biol. Ecol.* 225, 285-300.
- Robertson, A., Lenanton, R., 1984. Fish community structure and food chain dynamics in the surf zone of sandy beaches: the role of detached macrophyte detritus. *J. Exp. Mar. Biol. Ecol.* 84, 265-283.
- Rosenberg, G., Ramus, J., 1984. Uptake of inorganic nitrogen and seaweed surface area : volume ratios. *Aquat. Bot.* 19, 65-72.

1 Rothausler, E., Corell, H., Jormalainen, V., 2015. Abundance and dispersal trajectories
2 of floating *Fucus vesiculosus* in the Northern Baltic Sea. *Limnol. Oceanogr.* 60, 2173-
3 2184, doi: 10.1002/lno.10195.
4
5

6
7
8
9 Sales, M., Cebrian, E., Tomas, F., Ballesteros, E., 2011. Pollution impacts and recovery
10 potential in three species of the genus *Cystoseira* (Fucales, Heterokontophyta).
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Estuarine Coastal Shelf Sci 92, 347-357, doi: 10.1016/j.ecss.2011.01.008.

Sanchez-Moyano, J.E., Garcia-Adiego, E.M., Estacio, F.J., Garcia-Gomez, J.C., 2001.
Influence of the density of *Caulerpa prolifera* (Chlorophyta) on the composition of the
macrofauna in a meadow in Algeciras Bay (southern Spain). *Cienc. Mar.* 27, 47-71.

Santelices, B., 1990. Patterns of reproduction, dispersal and recruitment in seaweeds.
Oceanogr. Mar. Biol. Annu. Rev. 28, 177-276.

Schiel, D.R., 2004. The structure and replenishment of rocky shore intertidal
communities and biogeographic comparisons. *J. Exp. Mar. Biol. Ecol.* 300, 309-342.

Schiel, D.R., 2006. Rivets or bolts? When single species count in the function of
temperate rocky reef communities. *J. Exp. Mar. Biol. Ecol.* 338, 233-252.

Schiel, D.R., Lilley, S.A., 2007. Gradients of disturbance to an algal canopy and the
modification of an intertidal community. *Mar. Ecol. Prog. Ser.* 339, 1-11, doi:
10.3354/meps339001.

Schiel, D.R., Lilley, S.A., 2011. Impacts and negative feedbacks in community recovery
over eight years following removal of habitat-forming macroalgae. *Journal of*

Experimental Marine Biology and Ecology 407, 108-115, doi:

10.1016/j.jembe.2011.07.004.

Schiel, D.R., Taylor, D.I., 1999. Effects of trampling on a rocky intertidal algal assemblage in southern New Zealand. J. Exp. Mar. Biol. Ecol. 235, 213-235.

Schiel, D.R., Wood, S.A., Dunmore, R.A., Taylor, D.I., 2006. Sediment on rocky intertidal reefs: Effects on early post-settlement stages of habitat-forming seaweeds. J. Exp. Mar. Biol. Ecol. 331, 158-172, doi: 10.1016/j.jembe.2005.10.015.

Schoenwaelder, M.E.A., 2002. Physode distribution and the effect of 'thallus sunburn' in *Hormosira banksii* (Fucales, Phaeophyceae). Bot. Mar. 45, 262-266.

Schoenwaelder, M.E.A., Clayton, M.N., 1999. The presence of phenolic compounds in isolated cell walls of brown algae. Phycologia 38, 161-166.

Schreider, M.J., Glasby, T.M., Underwood, A.J., 2003. Effects of height on the shore and complexity of habitat on abundances of amphipods on rocky shores in New South Wales, Australia. J. Exp. Mar. Biol. Ecol. 293, 57-71.

Silberschneider, V., Booth, D.J., 2001. Resource use by *Enneapterygius rufopileus* and other rockpool fishes. Environ. Biol. Fishes 61, 195-2004.

Soltan, D., Verlaque, M., Boudouresque, C.F., Francour, P., 2001. Changes in macroalgal communities in the vicinity of a Mediterranean sewage outfall after the setting up of a treatment plant. Mar. Pollut. Bull. 42, 59-70, doi: 10.1016/S0025-326x(00)00116-8.

1 Speidel, M., Harley, C., Wonham, M., 2001. Recovery of the brown alga *Fucus*
2
3 *gardneri* following a range of removal intensities. *Aquat Bot* 71, 273-280.
4
5

6
7 Stauber, J.L., Binet, M., Duda, S., Krassoi, R., 2003. Toxicity assessment of secondary
8
9 treated sewage effluent from Melbourne Water's ETP - Final Report. CSIRO Energy
10
11 Technology Centre for Advanced Analytical Chemistry.
12
13

14
15 Steen, H., 2004. Interspecific competition between *Enteromorpha* (Ulvaes :
16
17 Chlorophyceae) and *Fucus* (Fucales : Phaeophyceae) germlings: effects of nutrient
18
19 concentration, temperature, and settlement density. *Mar. Ecol. Prog. Ser.* 278, 89-101.
20
21
22

23
24 Stekoll, M.S., Deysher, L., 1996. Recolonization and restoration of upper intertidal
25
26 *Fucus gardneri* (Fucales, Phaeophyta) following the Exxon Valdez oil spill.
27
28 *Hydrobiologia* 327, 311-316.
29
30

31
32 Strain, E.M.A., Thomson, R.J., Micheli, F., Mancuso, F.P., Airoidi, L., 2014.
33
34 Identifying the interacting roles of stressors in driving the global loss of canopy-forming
35
36 to mat-forming algae in marine ecosystems. *Global Change Biol.* 20, 3300-3312, doi:
37
38 10.1111/gcb.12619.
39
40
41

42
43 Suzuki, Y., Takabayashi, T., Kawaguchi, T., Matsunaga, K., 1998. Isolation of an
44
45 allelopathic substance from the crustose coralline algae, *Lithophyllum* spp., and its
46
47 effect on the brown alga, *Laminaria religiosa* Miyabe (Phaeophyta). *J Exp Mar Biol*
48
49 *Ecol* 225, 69-77, doi: 10.1016/s0022-0981(97)00208-6.
50
51
52

53
54 Tait, L.W., Hawes, I., Schiel, D.R., 2014. Shining Light on Benthic Macroalgae:
55
56 Mechanisms of Complementarity in Layered Macroalgal Assemblages. *PLoS ONE* 9,
57
58 e114146, doi: 10.1371/journal.pone.0114146.
59
60
61
62
63
64
65

1 Tait, L.W., Schiel, D.R., 2011. Legacy Effects of Canopy Disturbance on Ecosystem
2
3 Functioning in Macroalgal Assemblages. PLoS ONE 6, e26986, doi:
4
5 10.1371/journal.pone.0026986.
6
7

8
9 Taylor, D.I., Schiel, D.R., 2003. Wave-related mortality in zygotes of habitat-forming
10
11 algae from different exposures in southern New Zealand: the importance of 'stickability'.
12
13 J. Exp. Mar. Biol. Ecol. 290, 229-245.
14
15

16
17 Taylor, D.I., Schiel, D.R., 2005. Self-replacement and community modification by the
18
19 southern bull kelp *Durvillaea antarctica*. Mar. Ecol. Prog. Ser. 288, 87-102.
20
21

22
23 Terawaki, T., Hasegawa, H., Arai, S., Ohno, M., 2001. Management-free techniques for
24
25 restoration of *Eisenia* and *Ecklonia* beds along the central Pacific coast of Japan. J. Appl.
26
27 Phycol. 13, 13-17, doi: 10.1023/A:1008135515037.
28
29

30
31 Tewari, A., Joshi, H.V., 1988. Effect of domestic sewage and industrial effluents on
32
33 biomass and species diversity of seaweeds. Bot. Mar. 31, 389-397.
34
35

36
37 Thompson, R.C., Wilson, B.J., Tobin, M.L., Hill, A.S., Hawkins, S.J., 1996.
38
39 Biologically generated habitat provision and diversity of rocky shore organisms at a
40
41 hierarchy of spatial scales. J. Exp. Mar. Biol. Ecol. 202, 73-84.
42
43

44
45 Thomsen, M.S., Staehr, P.A., Nejrup, L., Schiel, D.R., 2013. Effects of the invasive
46
47 macroalgae *Gracilaria vermiculophylla* on two co-occurring foundation species and
48
49 associated invertebrates. Aquat Invasions 8, 133-145, doi: 10.3391/ai.2013.8.2.02.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Underwood, A.J., 1998. Grazing and disturbance: an experimental analysis of patchiness in recovery from a severe storm by the intertidal algal *Hormosira banksii* on rocky shores in New South Wales. J. Exp. Mar. Biol. Ecol. 231, 291-306.

Underwood, A.J., 1999. Physical disturbances and their direct effect on an indirect effect: responses of an intertidal assemblage to a severe storm. J. Exp. Mar. Biol. Ecol. 232, 125-140.

Underwood, A.J., Fairweather, P.G., 1989. Supply-side ecology and benthic marine assemblages. Trends Ecol. Evol. 4, 16-19.

van den Hoek, C., 1987. The possible significance of long-range dispersal for the biogeography of seaweeds. Helgolander Meeresuntersuchungen 41, 261-272.

van Tamelan, P.G., Stekoll, M.S., Deysher, L., 1997. Recovery process of the brown alga *Fucus gardneri* following the 'Exxon Valdez' oil spill: settlement and recruitment. Marine Ecology Progress Series 160, 265-277.

van Tamelen, P.G., Stekoll, M.S., Deysher, L., 1997. Recovery process of the brown alga *Fucus gardneri* following the 'Exxon Valdez' oil spill: settlement and recruitment. Mar. Ecol. Prog. Ser. 160, 265-277.

Viejo, R.M., Arenas, F., Fernandez, C., Gomez, M., 2008. Mechanisms of succession along the emersion gradient in intertidal rocky shore assemblages. Oikos 117, 376-389, doi: 10.1111/j.2007.0030-1299.16206.x.

Wahl, M., Molis, M., Hobday, A.J., Dudgeon, S., Neumann, R., Steinberg, P., Campbell, A.H., Marzinelli, E., Connell, S., 2015. The responses of brown macroalgae to

1 environmental change from local to global scales: direct versus ecologically mediated
2
3 effects. *Perspectives in Phycology* 2, 11-29.
4

5
6 Walker, D.I., Kendrick, G.A., 1998. Threats to macroalgal diversity: marine habitat
7
8 destruction and fragmentation, pollution and introduced species. *Botanica Marina* 41,
9
10 105-112.
11
12

13
14 Whitaker, S.G., Smith, J.R., Murray, S.N., 2010. Reestablishment of the Southern
15
16 California Rocky Intertidal Brown Alga, *Silvetia compressa*: An Experimental
17
18 Investigation of Techniques and Abiotic and Biotic Factors That Affect Restoration
19
20 Success. *Restor. Ecol.* 18, 18-26, doi: 10.1111/j.1526-100X.2010.00717.x.
21
22
23

24
25 Wikstrom, S.A., Kautsky, L., 2004. Invasion of a habitat-forming seaweed: effects on
26
27 associated biota. *Biol. Invasions* 6, 141-150.
28
29

30
31 Williams, S.L., Di Fiori, R.E., 1996. Genetic diversity and structure in *Pelvetia*
32
33 *fastigiata* (Phaeophyta: Fucales): does a small effective neighborhood size explain fine-
34
35 scale genetic structure? *Mar. Biol.* 126, 371-382.
36
37
38

39
40 Womersley, H.B.S., 1987. The marine benthic flora of southern Australia, Part II. South
41
42 Australian Government Printing Division, Adelaide.
43
44

45
46 Womersley, H.B.S., 1994. The marine benthic flora of southern Australia, Part IIIA.
47
48 Australian Biological Resources Study, Canberra.
49
50

51
52 Worm, B., Chapman, A.R.O., 1998. Relative effects of elevated grazing pressure and
53
54 competition from a red algal turf on two post-settlement stages of *Fucus evanescens* C.
55
56
57 Ag. *J. Exp. Mar. Biol. Ecol.* 220, 247-268.
58
59
60
61
62
63
64
65

1 Wright, J.P., Jones, C.G., 2006. The concept of organisms as ecosystem engineers ten
2 years on: progress, limitations, and challenges. *Bioscience* 56, 203-209.
3
4

5
6
7 Yoon, J.T., Sun, S.M., Chung, G., 2014. Sargassum bed restoration by transplantation
8 of germlings grown under protective mesh cage. *J. Appl. Phycol.* 26, 505-509, doi:
9 10.1007/s10811-013-0058-8.
10
11
12

13
14
15 Yu, Y.Q., Zhang, Q.S., Tang, Y.Z., Zhang, S.B., Lu, Z.C., Chu, S.H., Tang, X.X., 2012.
16 Establishment of intertidal seaweed beds of *Sargassum thunbergii* through habitat
17 creation and germling seeding. *Ecol. Eng.* 44, 10-17, doi:
18 10.1016/j.ecoleng.2012.03.016.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Number of papers and percentage of total in each study type published in *Marine Ecology Progress Series* from 1994 – 1996, 2003 – 2005 and 2013 – 2015. Data for 1994 -1996 are from Bruno and Bertness (2001).

nr, not recorded.

	1994 – 1996		2003 – 2005		2013 – 2015	
Study Type	Number	%	Number	%	Number	%
Population Ecology	nr	nr	412	18.0	226	15.1
Predation/Herbivory/Parasitism/ Scavenger	133	10.6	139	6.1	79	5.3
Physiology/Behaviour	144	11.5	284	12.4	205	13.7
Diet/Feeding Ecology	nr	nr	122	5.3	114	7.6
Consumers/Food Webs	nr	nr	37	1.6	101	6.7
Supply side ecology	76	6.1	114	5.0	87	5.8
Morphology, Biomechanics, Structure, Composition	14	1.1	62	2.7	110	7.3
Biogeochemistry	nr	nr	296	12.9	92	6.1
Management, Conservation, Monitoring, Marine Reserves	35	2.8	80	3.5	80	5.3
Migration/Movement Patterns	nr	nr	151	6.6	70	4.7
Disturbance	53	4.2	87	3.8	49	3.3
Ecosystem Engineers (Facilitation)	27	2.2	36	1.5	7	0.5
Natural/Life History	231	18.5	146	6.4	60	4

Genetics, phylogeny	32	2.6	92	4.0	56	3.7
Global Warming/Climate Change	nr	nr	9	0.4	46	3.1
Introduced/Invasive spp.	nr	nr	33	1.4	27	1.8
Pollution, Bioassay	45	3.6	40	1.7	13	0.9
Mortality Patterns/Diseases/Virus	nr	nr	37	1.6	23	1.5
Sediment	nr	nr	16	0.7	16	1.1
Transport/Erosion/Sedimentation						
Metal concentration/trace	nr	nr	25	1.1	3	0.2
metals/metal uptake						
Intracellular Symbiosis	6	0.5	7	0.3	7	0.5
Competition	15	1.2	32	1.4	12	0.8
Impact of antifouling	nr	nr	6	0.3	nr	nr
Other	441	35.2	31	1.3	14	0.9
Total	1252	100	2294	100	1497	100

Table 2. Number of papers and percentage of total in each study type published in *Restoration Ecology* from 2003 – 2005 and 2013 – 2015.

Study Type	2003 – 2005		2013 – 2015	
	Number	%	Number	%
Restoration of ecosystem engineers/facilitators	69	21.6	6	1.8
Disturbance (Natural/Human) +ve and -ve effects on restoration.	50	15.7	52	15.4
Population Ecology	40	12.5	35	10.4
Ecological theory to restoration ecology/Management/Monitoring/	35	11.0	22	6.5
Society & Environment/ Development/Conservation			35	10.4
Restoration of the physio-chemical environment	31	9.7	24	7.1
Introduced/Invasive/Exotic Species	25	7.8	16	4.7
Predators/Grazers/Competition/	23	7.2	29	8.6
Dispersal/Recruitment				
Plant Life Histories/Germination	11	3.5	25	7.4
Addition of single charismatic or rare species (including Endangered/Threatened species)	10	3.1	8	2.4
Restoration Success (Theory)	9	2.8	11	3.3
Sampling techniques for restoration treatments/methods for monitoring restoration success	7	2.2	54	16.0
Genetics of restoration/Gene flow	3	1.0	4	1.2

Food Web Management	3	1.0	6	1.8
Land Uses	2	0.6	5	1.5
Other	1	0.3	5	1.5
Total	319	100	337	100

Figure captions

Fig. 1. Schematic diagram expressing the current understanding of the autogenic ecosystem engineering roles of intertidal furoid algae (see text for further explanation and citations). Grey arrows indicate an increase (solid) or decrease (dashed) in physical factors due to the presence of a furoid canopy. Consequent positive (solid) and negative (coarse dashed) direct and indirect (fine dashed) interactions with other organisms and ecological processes are indicated by black arrows. Other indirect interactions are implied by the series of arrows.

Fig. 2. Schematic diagram expressing the current understanding of the effects of treated sewage effluent disposed into the marine environment on rocky intertidal communities (see text for further explanation and citations). Solid arrows indicate direct positive interactions, dashed arrows indicate direct negative interactions and perforated dashed arrows indicate indirect negative interactions. Cross indicates no interaction; question mark indicates that the possibility of an interaction has not yet been thoroughly explored and is thus unknown.

Figure(s)