Saitta ET, Rogers C, Brooker RA, Abbott GD, Kumar S, O'Reilly SS, Donohoe P,
Dutta S, Summons RE, Vinther J.

[Low fossilization potential of keratin protein revealed by experimental
taphonomy.](#)

Palaeontology (2017)

DOI: <https://doi.org/10.1111/pala.12299>

Copyright:

This is the peer reviewed version of the following article, which has been published in final form at <https://doi.org/10.1111/pala.12299>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

Date deposited:

06/06/2017

Embargo release date:

08 May 2018

1 Low fossilisation potential of keratin protein revealed by experimental taphonomy

2
3 Evan T. Saitta¹, Chris Rogers², Richard A. Brooker¹, Geoffrey D. Abbott³, Sumit Kumar⁴, Shane
4 S. O'Reilly^{5,6}, Paul Donohoe³, Suryendu Dutta⁴, Roger E. Summons⁶, & Jakob Vinther¹

5
6 ¹School of Earth Sciences, University of Bristol, Bristol BS8 1RJ, United Kingdom; e-mails:
7 evansaitta@gmail.com (ORCiD ID: orcid.org/0000-0002-9306-9060),
8 richard.brooker@bristol.ac.uk (ORCiD ID: [orchid.org/000-0003-4931-9912](https://orcid.org/000-0003-4931-9912)),
9 jakob.vinther@bristol.ac.uk (ORCiD ID: orcid.org/0000-0002-3584-9616)

10 ²School of Biological, Earth and Environmental Sciences, University College Cork, Cork T23
11 XA50, Ireland; e-mail: christopher.rogers@ucc.ie

12 ³School of Civil Engineering and Geosciences, Drummond Building, Newcastle University,
13 Newcastle NE1 7RU, UK; e-mails: geoff.abbott@newcastle.ac.uk (ORCiD ID: [orcid.org/0000-](https://orcid.org/0000-0001-9803-8215)
14 [0001-9803-8215](https://orcid.org/0001-9803-8215)), paul.donohoe@newcastle.ac.uk

15 ⁴Department of Earth Sciences, Indian Institute of Technology Bombay, Mumbai, Maharashtra
16 400 076, India; e-mails: sumit.kumar@iitb.ac.in, s.dutta@iitb.ac.in

17 ⁵ School of Earth Sciences, University College Dublin, Dublin 4, Ireland; e-mail:
18 shane.oreilly@ucd.ie (ORCiD ID: orcid.org/0000-0001-6303-4329)

19 ⁶Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of
20 Technology, Cambridge, MA 02139, USA; e-mail: rsummons@mit.edu (ORCiD ID:
21 [orchid.org/0000-0002-7144-8537](https://orcid.org/0000-0002-7144-8537))

ABSTRACT: Recent studies have suggested the presence of keratin in Cenozoic- to Mesozoic-aged fossils. However, ultrastructural studies revealing exposed melanosomes in many fossil keratinous tissues suggest that keratin should rarely, if ever, be preserved. In this study, keratin's stability through diagenesis was tested using microbial decay and maturation experiments on various keratinous structures. The residues were analyzed by pyrolysis-gas chromatography-mass spectrometry and compared to unpublished feather and hair fossils and published fresh and fossil melanin from squid ink. Results show that highly matured feathers (200–250 °C/250 bars/24 hours) become a volatile-rich, thick fluid with semi-distinct pyrolysis compounds from those observed in less degraded keratins (i.e., fresh, decayed, moderately matured, and decayed and moderately matured), suggesting hydrolysis of peptide bonds and potential degradation of free amino acids. Neither melanization nor keratin (secondary) structure (e.g., α - vs. β -keratin) produced different pyrograms – melanin pyrolysates are largely a subset of those from proteins and proteins have characteristic pyrolysates. Analyses of fossil fur and feather lacked amides, succinimide, and piperazines (present even in highly matured keratin) and showed pyrolysis compounds more similar to fossil and fresh melanin than to non-matured or matured keratin. Although the highly matured fluid was not water soluble at room temperature, it readily dissolved at elevated temperatures easily attained during diagenesis, meaning it can leach away from the fossil. Future interpretations of fossils must consider that calcium phosphate and pigments are the only components of keratinous structures known to survive fossilisation in mature sediments.

KEY WORDS: keratin, protein, experimental taphonomy, Py-GC-MS, fossilisation

BIOMOLECULES vary in preservation potential. From least to most robust, these are roughly: nucleic acids, proteins, carbohydrates, aromatics, and lipids (Briggs and Summons 2014). The fossilization potential of these biomolecules can also be influenced by the interaction with inorganic components (Curry *et al.* 1991). Keratin is a diverse family of fibrous structural proteins and a common component of vertebrate tissues. Some keratinous structures are hardened through deposition of calcium phosphate salts (Pautard 1964) such as baleen, claws and feather rachises (Blakey *et al.* 1963). These salts are known from the fossil record (Mayr *et al.* 2016; Vinther *et al.* 2016) as thin, usually white, phosphate sheets (Benton *et al.* 2008; Bergmann *et al.* 2010). Tissues bearing melanin—a widely distributed group of pigments, also preserve well as original organic remains (Vinther *et al.* 2008; Vinther 2015). However, the survival of keratin protein itself remains controversial given that keratinous materials such as feathers or textiles are considered to have poor preservation in the archaeological record (Hargrave 1960; Messinger 1965; Brom 1986; Reinhard & Bryant 1992; Rogers *et al.* 2002; Dove *et al.* 2005). Studies have claimed to have found evidence of intact keratin based on immunohistochemistry experiments (Schweitzer *et al.* 1999a, 1999b; Moyer *et al.* 2016; Pan *et al.* 2016). These experiments indicate the preservation of tertiary protein structure, but use a method prone to false positives or statistical artefacts (Buckley *et al.* 2007; True 2008; Bern *et al.* 2009). Other studies on fossil keratinous structures have relied on infrared spectroscopy, recovering amide bands (Manning *et al.* 2009; Edwards *et al.* 2011), although pyrolysis-gas chromatography-mass spectrometry (Py-GC-MS) data did not yield any unambiguous markers for amino acids. Here we determine through taphonomic experiments what signatures degraded and intact keratin might leave in the fossil record and clarify if keratin is likely to persist in exceptionally preserved fossil material over million-year time scales.

MATERIAL AND METHODS

Various integumentary structures, such as feathers, hair, and scales, were subjected to microbial decay and maturation treatments (Table 1; extended methods in Saitta *et al.* 2017). Samples were either (a) fresh, (b) microbially decayed (with a treatment using cultured, naturally occurring feather microbes in a salt broth at ~37 °C/50 days), (c) matured and not decay treated, or (d) matured after decay treatment. These were chemically analysed using Py-GC-MS (analytical details in Saitta *et al.* 2017). Samples were sealed in noble metal (Au⁹⁰ Pd¹⁰) capsules and maturation was performed in a cold-seal, water pressurized autoclave ranging from 100–250 °C (measured with a calibrated K-type thermocouple) and 250 bars for 24 hours. The product of the highest condition maturation experiments (250 °C/250 bars/24 hours) on feather samples was subjected to solubility tests in water at room temperature for a year and then in a steam autoclave at 121 °C for 45 minutes.

Analyses were compared to previously obtained and unreported Py-GC-MS data on fossil hair from the Eocene *Palaeochiropteryx* (Messel Shale, SMF-ME 11406a) and feather material from a Lower Eocene bird head (Fur Formation, Danekræ 200, MGUH 28.929). Associated sediment was also analysed as a control and these were compared to published fossil and modern squid ink data as a reference for melanin (Glass *et al.* 2012).

Institutional abbreviations

Fossil specimens for which novel data is presented are deposited at the following institutions:
Geological Museum of Copenhagen (MGUH), Copenhagen, Denmark and Naturmuseum
Senckenberg (SMF), Frankfurt, Germany.

RESULTS

Very similar pyrolysis compounds were observed in the fresh, decayed, moderately matured, and decayed and moderately matured samples, regardless of keratin type or melanisation (Fig. 1; extended results in Saitta *et al.* 2017) including acetic acid, nitriles, pyridines, benzenes, toluene, amides, pyrrole-related compounds, indoles, phenols, styrene, and 2,5-diketopiperazine. Highly matured feathers (250 °C/250 bars/24 hours) lost all structural integrity and became a viscous liquid with a strong odour. Pyrolysis compounds for these highly matured samples did not differ between the white and dark feathers. Highly matured feathers shared many similar categories of compounds with the less degraded samples (i.e., nitriles, benzenes, toluene, amides, pyrrole-related compounds, phenols, styrene, and 2,5-diketopiperazine) but specific compounds showed distinct variations and their pyrograms produced noticeably different ‘fingerprints’ (Fig. 1). Highly matured feathers had more diverse nitriles and amides and less diverse benzenes, pyrrole-related compounds, and phenols compared to the less degraded samples. The highly matured feather fluid product showed no sign of dissolution when left in water at room temperature for over one year. However, placing this same vial in a steam autoclave (121 °C/45 minutes) resulted in rapid dissolution. It should be noted that earlier maturation runs on white feathers matured at 200 °C and 250 °C (250 bars/24 hours) in an argon gas autoclave resulted in a similar

fluid that leaked out of the unsealed sample tubes and could not be analysed, which prompted the protocol ultimately used here.

Both the fossil hair and feathers and their associated sediments yielded pyrolysates containing nitrogen or sulphur (Table 2). Slight differences were apparent between the two localities, and the sediments yielded many compounds with similar affinities to the fossils, which may be due to the presence of aromatics, such as algal porphyrins. Although some categories of compounds are shared between the fossils and experimental keratin (Table 2), fossils lacked acetic acid, pyridines, toluene, amides, indoles, styrene, succinimide, and piperazines found in the experimental keratin. Furthermore, the fossils shared many categories of compounds with the melanin of squid ink: nitriles, benzenes, pyrrole-related compounds, thiophenes, and phenols (Table 2). The fossils, compared to the fresh and treated experimental keratin and melanin, showed many unique ketones, pyrrole-related compounds, and thiophenes. Caution must be taken when comparing Py-GC-MS datasets obtained separately (Fossil data analysed at MIT, keratin data analysed at the University of Newcastle); therefore, it is important to compare more general categories of compounds rather than precise chemical species between the experiment keratin and the fossils/sediment.

DISCUSSION

The pyrolysis results from the experimental keratin are consistent with previous studies using Py-GC-MS on keratin (Brebou and Spiridon 2011) and protein in general (Bland *et al.* 1998; Reeves and Francis 1998). Py-GC-MS can detect protein or degraded protein but cannot distinguish between different keratin types nor can it distinguish between melanized and non-

136 melanized keratin. This is not entirely unexpected given that proteins appear to produce fairly
137 characteristic pyrolysates. Different keratin types are not molecularly diverse enough to result in
138 distinct pyrolysates. Also, pyrolysis compounds from melanin (Glass *et al.* 2012; Dzierżęga-
139 Łęcznar *et al.* 2012) are largely a subset of those from protein; melanin is a polymer formed from
140 oxidation products of the amino acid tyrosine. Py-GC-MS can distinguish between highly
141 matured and less degraded keratin. Succinimide was found in the highly matured feathers. It is
142 known to be linked to non-enzymatic protein degradation, acting as an intermediate during
143 deamidation, racemization, and isomerization (Geiger and Clarke 1987; Stephenson and Clarke
144 1989). However, these studies observed low temperature reactions, and the succinimide detected
145 here would be a product of pyrolysis. Combined with the observed liquefaction, the results
146 suggest a breakdown of the protein structure and peptide bonds, is likely coupled with further
147 breakdown of the free amino acids in the highly matured feathers since they experienced
148 temperatures within the range of many amino acid decomposition temperatures (Dunn and
149 Brophy 1932; Lien and Nawar 1974). Diketopiperazines were found in all of the experimental
150 keratin. Although they can be byproducts of peptide terminal cleavage (Martins and Carvalho
151 2007), those identified here likely formed during pyrolysis of either peptides or free amino acids.
152 It is possible that side chains from at least one amino acid are preserved in 2,5-diketopiperazine
153 homologs. Although beyond the scope of this study, such information could provide insight into
154 the original protein amino acid composition (e.g., keratins are highly conserved with high Cys,
155 Gly, Pro, and Ser and low His, Lys, Trp, and Glu). Similar insight might be gathered from
156 detailed examination of other characteristic protein pyrolysates such as amides and succinimide.
157 Thus, if a fossil were to contain these protein pyrolysis markers, there is potential to further
158 investigate their structure and elucidate the composition of the proteins they derived from.

159 However, such markers were not found in the fossil samples analysed here, making this line of
160 investigation unnecessary, and the proposed method should first be verified to determine just
161 how much variation in pyrolysates would be expected from different proteins and whether such
162 variation could be diagnostically useful. Similarly, methodological advances allowing for
163 analysis of Py-GC-MS data in a statistical framework would further improve comparisons of
164 chemical composition.

165   No unambiguous protein markers were found in the geologic samples, either in the
166 sediment or the fossil material. Unambiguous protein pyrolysis markers are identified here as
167 those categories of compounds that are present in fresh or degraded experimental keratin and
168 absent from melanins. The similarities in pyrolysates between the sediment and the fossil from
169 both localities, including a suite of aromatic compounds, could suggest protein or amino acid
170 presence. However, these pyrolysates have been found in melanins and could also be products of
171 other aromatic compounds, such as humic acids and porphyrins (Meuzelaar 1977). Aromatic
172 compounds are also formed during pyrolysis of biomacromolecules. Regardless, the lack of
173 amides, succinimide, and piperazines in the fossils demonstrate that proteinaceous material is
174 lacking. Therefore, the compounds in the fossils appear overall consistent with melanin than with
175 keratin protein. TOF SIMS analysis of the same fossil mammal hair and other feathers from the
176 Fur Formation confirm melanin preservation in these structures (Colleary *et al.* 2015, Gren *et al.*
177 2016). Furthermore, both melanin and the fossils contained thiophenes, absent from the
178 experimental keratin – although thiophenes were found in what was likely melanized keratin by
179 Brebu and Spiridon (2011), in addition to thiazoles, which are known phaeomelanin markers
180 (Dzierżęga-Lęcznar *et al.* 2012). Thiophenes in fossils also occur through diagenetic
181 incorporation of sulfur in iron-poor depositional environments (Glass *et al.* 2012). Manning *et al.*

(2009) and Edwards *et al.* (2011) presented Py-GC-MS data on fossil reptile and hadrosaur skin, claiming it supported the presence of keratin. Although some of the compounds reported included nitriles, thiophenes, and benzene derivatives which are known from proteins, these are also known from fossil melanin. Similar to the fur and feather fossils examined here, the compounds are not a good match to those expected from keratin, which should present itself as a broader suite of pyrolysates. Their reports of aliphatic compounds also strongly resemble those reported through lipid *in situ* geopolymerization (Briggs 1999; Gupta *et al.* 2007). Fourier transform infrared spectroscopy has also been used to identify amides in the fossil reptile and hadrosaur skin (Manning *et al.* 2009; Edwards *et al.* 2011). However, the diverse amides present in the highly matured feather fluid prove, not surprisingly, that the presence of amides does not necessitate the presence of intact proteins, as amides can be present in simpler biomolecules. Amides were not found in pyrolysis studies of the fossil material, which raises questions as to the validity of the FTIR results and suggests the need to confirm the presence of amides detected with FTIR using other methods such as Py-GC-MS. In general, Manning *et al.* (2009) and Edwards *et al.* (2011) fail to prove the presence of intact keratin or distinguish between potential protein breakdown products from melanin or other contaminants.

Artificial maturation is known to mimic the chemical changes that occur during diagenesis. Temperatures and pressures are kept high (e.g., 200–250 °C, which corresponds to the last stages of the oil window) for short durations (e.g., days–weeks) in order to speed up chemical reactions that would occur at relatively more moderate conditions but over longer durations (e.g., millions of years). With regards to organic preservation it is important to regard both the age of the fossil and its burial history. The Messel Oil Shale, for example, is considered “thermally immature” (Ocampo *et al.* 1985) so our failure to find key protein pyrolysates in a

Messel fossil bones poorly for keratin preservation potential in fossils of equal or greater age than the Lower Eocene and possibly even younger fossils.

Our inability to reproduce the results of McNamara *et al.* (2013, p.2) is explained by a *lapsus calami* in their methods which should state “each [maturation] experiment lasted for 1 h” not “24 h” (McNamara personal communication 2016). However, the absence of keratin proteins in the fossil record suggests the timeframe of our maturation experiments, in which original keratin protein was absent, may be a better approximation of the conditions that occur during diagenesis. Maturation experiments must be of a duration long enough to ensure accurate modeling of diagenetic chemical reactions, and the one hour experiments of McNamara *et al.* (2013) are not sufficient to draw conclusions as to the preservation potential of keratin in the fossil record.

Although the highly matured feather fluid was not water soluble at room temperature, which is expected given the relatively high abundance of hydrophobic and neutral amino acids compared to hydrophilic amino acids (at pH 7) in β -keratin (Dalla Valle *et al.* 2009), it was easily dissolved in our experiments at elevated temperatures that might be expected during diagenesis. This provides a model for how hydrolyzed and degraded proteins might aqueously dissolve and leach away from fossils.

Feathers and textiles produced from keratin are rare from archaeological sites. Toxic metals can slow keratin degradation and allow the peptides to persist in some sites, but the presence of iron only allows for the preservation of a corrosion cast. Although keratin appears robust compared to many other proteins, it is not resistant to millennia of diagenetic forces (Hargrave 1960; Messinger 1965; Brom 1986; Reinhard & Bryant 1992; Rogers *et al.* 2002; Dove *et al.* 2005). Studies of well-preserved fossil feathers have found that organic residues

consist of melanosomes, usually in alignment along barb and barbule axes (Vinther *et al.* 2008; Vinther 2015). Furthermore, colour patterns are frequently observed (Vinther *et al.* 2008; Field *et al.* 2013; Vinther *et al.* 2016), which are observed through the macroscopic presence or absence of organic material composed of melanosomes. Areas lacking melanosomes preserve nothing but feather impressions in the rock matrix (Vinther *et al.* 2008). No substance conclusively attributable to keratin has been found in any well-preserved fossil feathers, even though amorphous organic residues in more weathered specimens from localities such as the Jehol or Yanliao biota and the Green River Formation have been suggested to be keratin by some (Schweitzer *et al.* 1999a; McNamara *et al.* 2013). The nature of this amorphous organic residue remains to be fully described. However, Lindgren *et al.* (2015) found signatures consistent with melanin using TOF SIMS and FTIR on feathers preserved in a Late Jurassic specimen of *Anchiornis huxleyi* preserved as solid organic material along with traces of melanosomes as impressions and merged blocks of melanosomes. The only other constituent preserved in the feathers were calcium phosphate, either derived from the hardened keratin as observed elsewhere (Mayr *et al.* 2016; Vinther *et al.* 2016) or through secondary mineralisation.

CONCLUSION

The production of a viscous fluid from highly matured feathers with a characteristic Py-GC-MS signature compared to less degraded keratin suggests that keratin degrades during diagenesis and catagenesis. Py-GC-MS of fossil keratinous structures supports our experimental data, showing that non-lipid, nitrogen and sulfur containing compounds more closely match recalcitrant fossil and fresh melanin than keratin protein. Keratin can decay through microbial action and

diagenetic hydrolysis of peptide bonds. An analogous process to collagen gelatinization (Pfretzschner 2006) might occur in keratin to produce a viscous fluid. Unlike proteins encapsulated within a mineralized matrix, such as bone or shell, epidermal keratin is not stabilised nor shielded from the environment, making it more prone to microbial decay, aqueous hydrolysis, and ultimately dissolution, leaching from the fossil at higher temperatures during later diagenesis. Keratin degradation means that calcium phosphate and melanin are likely the only remaining components of keratinous structures detectable in fossils. Reports of keratin proteins in fossils, especially of Mesozoic age (Schweitzer *et al.* 1999*a*, 1999*b*; Moyer *et al.* 2016; Pan *et al.* 2016), should be reexamined. Furthermore, thermal experiments that report preserved morphological structures of feathers with weak and diffuse antibody signal persisting after a decade-long exposure at 350 °C (Moyer *et al.* 2016) not only contradict the results here, they are likely impossible given reported thermal decomposition points of amino acids (Dunn & Brophy 1932; Lien & Nawar 1974). Immunohistological keratin detection in such experimental and fossil cases are likely false positives from humic acids formed through degradation of biomolecules (Collins *et al.* 1992). This is best exemplified by the claims for immunodetection of keratin in feathers which had been converted to ash (covered in foil and placed in a 350 °C dry oven for a decade) (Moyer *et al.* 2016). Py-GC-MS, in contrast, presents itself as a useful tool for identifying protein-derived organics (i.e., those containing amino acids) in fossil samples.

Acknowledgments. Many thanks to E. Thornberry (University of Bristol) and I. Bull (University of Bristol) for feedback regarding Py-GC-MS, A. Soler (La Ferme aux Crocodiles) for dissecting and shipping the crocodile skin. SOR acknowledges funding from the Marie Skłodowska-Curie Actions Research Fellowship Programme and the Irish Research Council (ELEVATE Career-

Development Fellowship ELEVATEPD/2014/47). Work at MIT was also supported by an award (NNA13AA90A) from the NASA Astrobiology Institute. Thanks to the helpful comments of all those who reviewed the work including M. Collins (University of York) and anonymous reviewers. We thank M. McNamara for clarification on her experimental parameters. ETS writes in memory of Robert S. Kearney.

DATA ARCHIVING STATEMENT

Data for this study are available in the Dryad Digital Repository:

<http://datadryad.org/review?doi=doi:10.5061/dryad.h02q0>

REFERENCES

- BENTON, M. J., ZHOU, Z., ORR, P. J., ZHANG, F. and KEARNS, S. L. 2008. The remarkable fossils from the Early Cretaceous Jehol Biota of China and how they have changed our knowledge of Mesozoic life. *Proceedings of the Geologists' Association*, **119**, 209–228.
- BERGMANN, U., MORTON, R. W, MANNING, P. L., SELLERS, W. I., FARRAR, S., HUNTLEY, K. G., WOGELIUS, R. A. and LARSON, P. 2010. *Archaeopteryx* feathers and bone chemistry fully revealed via synchrotron imaging. *Proceedings of the National Academy of Sciences*, **107**, 9060–9065.
- BERN, M., PHINNEY, B. S. and GOLDBERG, D. 2009. Reanalysis of *Tyrannosaurus rex* mass spectra. *Journal of Proteome Research*, **8**, 4328–4332.

296 BLAKEY, P. R., EARLAND, C. and STELL, J. G. P. 1963. Calcification of keratin. *Nature*,
 297 **198**, 481.

298 BLAND, H. A., VAN BERGEN, P. F., CARTER, J. F. and EVERSHED, R. P. 1998. Early
 299 diagenetic transformations of proteins and polysaccharides in archaeological plant
 300 remains. In STANKIEWICZ, B. A. and VAN BERGEN, P. F. (eds). *Nitrogen-*
 301 *Containing Macromolecules in the Bio- and Geosphere*. American Chemical Society,
 302 Washington, DC, **113-31**, 370 pp.

303 BREBU, M. and SPIRIDON, I. 2011. Thermal degradation of keratin waste. *Journal of*
 304 *Analytical and Applied Pyrolysis*, **91**, 288–295.

305 BRIGGS, D. E. 1999. Molecular taphonomy of animal and plant cuticles: selective preservation
 306 and diagenesis. *Philosophical Transactions of the Royal Society of London B: Biological*
 307 *Sciences*, **354**, 7–17.

308 — and SUMMONS, R. E. 2014. Ancient biomolecules: their origins, fossilization, and role in
 309 revealing the history of life. *BioEssays*, **36**, 482–490.

310 BROM, T. G. 1986. Microscopic identification of feathers and feather. *Bijdragen tot de*
 311 *dierkunde*, **56**, 181–204.

312 BUCKLEY, M., WALKER, A., HO, S. Y., YANG, Y., SMITH, C., ASHTON, P., OATES, J.
 313 T., CAPPELLINI, E., KOON, H., PENKMAN, K., ELSWORTH, B., ASHFORD, D.,
 314 SOLAZZO, C., ANDREWS, P., STRAHLER, J., SHAPIRO, B., OSTROM, P.,
 315 GANDHI, H., MILLER, W., RANEY, B., ZYLBER, M. I., GILBERT, M. T.,
 316 PRIGODICH, R. V., RYAN, M., RIJSDIJK, K. F., JANOO, A. and COLLINS, M. J.
 317 2007. Protein sequences from mastodon and *Tyrannosaurus rex* revealed by mass
 318 spectrometry. Comment. *Science*, **318**, 33c.

319 COLLEARY, C., DOLOCAN, A., GARDNER, J., SINGH, S., WUTTKE, M., RABENSTEIN,
320 R., HABERSETZER, J., SCHAAL, S., FESEHA, M., CLEMENS, M. and JACOBS, B.
321 F. 2015 Chemical, experimental, and morphological evidence for diagenetically altered
322 melanin in exceptionally preserved fossils. *Proceedings of the National Academy of*
323 *Sciences*, **112**, 12592–12597.

324 COLLINS, M. J., WESTBROEK, P., MUYZER, G. and DE LEEUW, J. W. 1992. Experimental
325 evidence for condensation reactions between sugars and proteins in carbonate skeletons.
326 *Geochimica et Cosmochimica Acta*, **56**, 1539–1544.

327 CURRY, G. B., CUSACK, M., WALTON, D., ENDO, K., CLEGG, H., ABBOTT, G.,
328 ARMSTRONG, H., BADA, J. L. AND ENGEL, M. H. 1991. Biogeochemistry of
329 brachiopod intracrystalline molecules [and discussion]. *Philosophical Transactions of the*
330 *Royal Society of London B: Biological Sciences*, **333**, 359–366.

331 DALLA VALLE, L., NARDI, A., TONI, M., EMERA, D. and ALIBARDI, L. 2009.
332 Beta-keratins of turtle shell are glycine-proline-tyrosine rich proteins similar to those of
333 crocodilians and birds. *Journal of Anatomy*, **214**, 284–300.

334 DOVE, C. J., HARE, P. G. and HEACKER, M. 2005. Identification of ancient feather fragments
335 found in melting alpine ice patches in southern Yukon. *Arctic*, **58**, 38–43.

336 DUNN, M. S. and BROPHY, T. W. 1932. Decomposition points of the amino acids. *Journal of*
337 *Biological Chemistry*, **99**, 221–229.

338 DZIERŻĘGA-LĘCZNAR, A., KURKIEWICZ, S. and STĘPIEŃ, K. 2012. Detection and
339 quantitation of a pheomelanin component in melanin pigments using pyrolysis–gas
340 chromatography/tandem mass spectrometry system with multiple reaction monitoring
341 mode. *Journal of Mass Spectrometry*, **47**, 242–245.

342 EDWARDS, N. P., BARDEN, H. E., VAN DONGEN, B. E., MANNING, P. L., LARSON, P.
 343 L., BERGMANN, U., SELLERS, W. I. and WOGELIUS, R. A. 2011. Infrared mapping
 344 resolves soft tissue preservation in 50 million year-old reptile skin. *Proceedings of the*
 345 *Royal Society of London B: Biological Sciences*, **278**, 3209–3218.

346 FIELD, D. J., D’ALBA, L., VINTHER, J., WEBB, S. M., GEARTY, W. and SHAWKEY, M. D.
 347 2013. Melanin concentration gradients in modern and fossil feathers. *PloS one*, **8**,
 348 e59451.

349 GEIGER, T. and CLARKE, S. 1987. Deamidation, isomerization, and racemization at
 350 asparaginy and aspartyl residues in peptides. Succinimide-linked reactions that
 351 contribute to protein degradation. *Journal of Biological Chemistry*, **262**, 785–794.

352 GLASS, K., ITO, S., WILBY, P.R., SOTA, T., NAKAMURA, A., BOWERS, C.R., VINTHER,
 353 J., DUTTA, S., SUMMONS, R., BRIGGS, D.E. and WAKAMATSU, K. 2012. Direct
 354 chemical evidence for eumelanin pigment from the Jurassic period. *Proceedings of the*
 355 *National Academy of Sciences*, **109**, 10218–10223.

356 GREN, J. A., SJÖVALL, P., ERIKSSON, M. E., SYLVESTERSEN, R. L., MARONE, F.,
 357 SIGFRIDSSON CLAUSS, K. G., TAYLOR, G. J., CARLSON, S., UVDAL, P. and
 358 LINDGREN, J. 2016. Molecular and microstructural inventory of an isolated fossil bird
 359 feather from the Eocene Fur Formation of Denmark. *Palaeontology*, **60**, 73–90.

360 GUPTA, N. S., BRIGGS, D. E. G., COLLINSON, M. E., EVERSHED, R. P., MICHELS, R.,
 361 JACK, K. S. and PANCOST, R. D. 2007. Evidence for the *in situ* polymerisation of
 362 labile aliphatic organic compounds during the preservation of fossil leaves: implications
 363 for organic matter preservation. *Organic Geochemistry*, **38**, 499–522.

364 HARGRAVE, L. L. 1960. Identification of archeological feathers from Glen Canyon, Utah.

365 *University of Utah Anthropological Papers*, **44**, 239–241.
 366 LIEN, Y. C. and NAWAR, W. W. 1974. Thermal decomposition of some amino acids. Valine,
 367 leucine and Isoleucine. *Journal of Food Science*, **39**, 911–913.
 368 MANNING, P.L., MORRIS, P.M., MCMAHON, A., JONES, E., GIZE, A., MACQUAKER,
 369 J.H., WOLFF, G., THOMPSON, A., MARSHALL, J., TAYLOR, K.G. and LYSON, T.
 370 2009. Mineralized soft-tissue structure and chemistry in a mummified hadrosaur from the
 371 Hell Creek Formation, North Dakota (USA). *Proceedings of the Royal Society of London*
 372 *B: Biological Sciences*, **276**, 3429–3437.
 373 MARTINS, M. B. and CARVALHO, I. 2007. Diketopiperazines: biological activity and
 374 synthesis. *Tetrahedron*, **63**, 9923–9932.
 375 MAYR, G., PITTMAN, M., SAITTA, E., KAYE, T. G. and VINTHER, J. 2016. Structure and
 376 homology of *Psittacosaurus* tail bristles. *Palaeontology*, **59**, 793–802.
 377 MCNAMARA, M.E., BRIGGS, D. E. G., ORR, P. J., FIELD, D. J. and WANG, Z. 2013.
 378 Experimental maturation of feathers: implications for reconstructions of fossil feather
 379 colour. *Biology Letters*, **9**, 20130184.
 380 MESSINGER, N. G. 1965. Methods used for identification of feather remains from Wetherill
 381 Mesa. *Memoirs of the Society for American Archaeology*, **19**, 206–215.
 382 MEUZELAAR, H. L. C., HAIDER, K., NAGAR, B. R. and MARTIN, J. P. 1977. Comparative
 383 studies of pyrolysis-mass spectra of melanins, model phenolic polymers, and humic
 384 acids. *Geoderma*, **17**, 239–252.
 385 MOYER, A. E., ZHENG, W. and SCHWEITZER, M. H. 2016. Keratin durability has
 386 implications for the fossil record: results from a 10 year feather degradation experiment.
 387 *PloS one*, **11**, e0157699.

388 OCAMPO, R., CALLOT, H. J. and ALBRECHT, P. 1985. Identification of polar porphyrins in
 389 oil shale. *Journal of the Chemical Society, Chemical Communications*, **4**, 198–200.

390 PAN, Y., ZHENG, W., MOYER, A.E., O’CONNOR, J.K., WANG, M., ZHENG, X., WANG,
 391 X., SCHROETER, E.R., ZHOU, Z. And SCHWEITZER, M.H. 2016. Molecular
 392 evidence of keratin and melanosomes in feathers of the Early Cretaceous bird
 393 *Eoconfuciusornis*. *Proceedings of the National Academy of Sciences*, **113**, E7900–E7907.

394 PAUTARD, F. G. E. 1964. Calcification of keratin. In ROOK, A. and CHAMPION, R. H. (eds).
 395 *Progress in the biological sciences in relation to dermatology – 2*. Cambridge University
 396 Press, London, **227-40**, 499 pp.

397 PFRETZSCHNER, H. U. 2006. Collagen gelatinization: the key to understand early bone-
 398 diagenesis. *Palaeontographica Abteilung A*, **278**, 135–148.

399 REEVES, J. B. III and FRANCIS, B. A. 1998. Pyrolysis-gas chromatography for the analysis of
 400 proteins: with emphasis on forages. In STANKIEWICZ, B. A. and VAN BERGEN, P. F.
 401 (eds). *Nitrogen-Containing Macromolecules in the Bio- and Geosphere*. American
 402 Chemical Society, Washington, DC, **47-62**, 370 pp.

403 REINHARD, K. J. and BRYANT, V. M. 1992. Coprolite analysis: a biological perspective on
 404 archaeology. *Archaeological Method and Theory*, **4**, 245–288.

405 ROGERS, J. D., DOVE, C. J., HEACKER, M. and GRAVES, G. R. 2002. Identification of
 406 feathers in textiles from the Craig Mound at Spiro, Oklahoma. *Southeastern Archaeology*,
 407 **21**, 245–251.

408 SAITTA, E. T., ROGERS, C., BROOKER, R. A., ABBOTT, G. D., KUMAR, S., O'REILLY, S.
 409 S., DONOHOE, P., DUTTA, S., SUMMONS, R. E., and VINTHER, J. 2017. Data from:

- Low fossilisation potential of keratin protein revealed by experimental taphonomy. Dryad Digital Repository. <http://dx.doi.org/10.5061/dryad.h02q0>
- SCHWEITZER, M. H., WATT, J. A., AVCI, R., FORSTER, C. A., KRAUSE, D. A., KNAPP, L., ROGERS, R. R., BEECH, I. and MARSHALL, M. 1999a. Keratin immunoreactivity in the Late Cretaceous bird *Rahonavis ostromi*. *Journal of Vertebrate Paleontology*, **19**, 712–722.
- WATT, J. A., AVCI, R., KNAPP, L., CHIAPPE, L., NORELL, M. and MARSHALL, M., 1999b. Beta-keratin specific immunological reactivity in feather-like structures of the cretaceous alvarezsaurid, *Shuvuuia deserti*. *Journal of Experimental Zoology*, **285**, 146–157.
- STEPHENSON, R. C. and CLARKE, S. 1989. Succinimide formation from aspartyl and asparaginyl peptides as a model for the spontaneous degradation of proteins. *Journal of Biological Chemistry*, **264**, 6164–6170.
- TRUE, L. D. 2008. Quality control in molecular immunohistochemistry. *Histochemistry and Cell Biology*, **130**, 473–480.
- VINTHER, J., BRIGGS, D. E., PRUM, R. O. and SARANATHAN, V. 2008. The colour of fossil feathers. *Biology Letters*, **4**, 522–525.
- 2015. A guide to the field of palaeo colour. *BioEssays*, **37**, 643–656.
- NICHOLLS, R., LAUTENSCHLAGER, S., PITTMAN, M., KAYE, T. G., RAYFIELD, E., MAYR, G. and CUTHILL, I. C. 2016. 3D Camouflage in an ornithischian dinosaur. *Current Biology*, **26**, 2456–2462.

FIGURES

433

434 **Fig. 1.** Total ion pyrograms of feathers. Subpanel shows highly matured viscous fluid from white
 435 feathers (scale bar represents 500 μm). ?, tentative compound identification. Letters, compounds
 436 present in less degraded samples. a, acetic acid. b, propanenitrile. c, 2-methyl propanenitrile. d,
 437 2-methyl butanenitrile. e, pyridine. f, pyrrole. g, toluene. h, methyl pyridine. i, methyl pyrrole. j,
 438 ethyl benzene. k, styrene. l, ethyl pyrrole. m, phenol. n, 4-methyl phenol. o, ethyl cyanobenzene.
 439 p, propyl cyanobenzene. q, indole. r, methyl indole. s, 2,5-diketopiperazine. t, trimethyl
 440 methoxyphenol. u, octadecanamide. Numbers, compounds exclusive to highly matured samples.
 441 1, benzene. 2, acetamide. 3, methyl pentanenitrile. 4, propanamide. 5, C1-styrene. 6, methyl
 442 butanamide. 7, C1-phenol. 8, hexanamide. 9, methyl pentanamide. 10, piperidinone. 11,
 443 hexadecyl nitrile. 12, hexadecanamide. 13, docosenenyl nitrile. 14, eicosanamide. 15,
 444 docosenamide. 16, docosanamide. 17, pyrrolidine, 1-acetyl-. 18, N-[2-hydroxyethyl]succinimide.
 445 Slight retention time misalignment for the same peak between samples is a result of running
 446 Py/GC/MS at different times.

447

448 TABLES

449

450 **Table 1.** List of maturation runs.

451

452 **Table 2.** Comparison of pyrolysates.

TABLE 1. LIST OF MATURATION RUNS

Sample	Notes
<u>Highly matured - 250°C, 250 bars, 24 hours</u>	
White feathers*	Chicken (<i>Gallus gallus</i>); Leg region
Dark feathers*	Turkey (<i>Meleagris gallopavo</i>); Leg region; Colour gradient (from proximal to distal) of white, grey, black, and iridescent
<u>Moderately matured - 100°C, 250 bars, 24 hours</u>	
Feathers – iridescent*	Turkey; Wing covert
Feathers – white*	Chicken; Wing covert
Feathers – black*	Chicken; Back/neck region; White fringe on vane
Avian scutate scales [†]	Turkey
Avian reticulate scales [§]	Turkey; Multiple scales with associated epidermis
Turkey beard – adult* [†]	Dried samples; Small portion of epidermis still attached
Turkey beard – juvenile* [†]	Portion of epidermis still attached
Crocodilian scale – black [§]	Nile crocodile (<i>Crocodylus niloticus</i>); Flank region; Predominantly black colouration
Crocodilian scale – white [§]	Nile crocodile; Flank region; Predominantly white/light colouration
Mammalian hair [#]	Horse (<i>Equus ferus</i>) mane; Black colour; ‘Outgroup’

Decayed feathers - iridescent*	Turkey; Back region
Decayed feathers - white*	Chicken; Back/neck region
Decayed feathers - black*	Chicken; Back/neck (white fringe on vane) and tail regions

---

Note: Samples that are not listed as decayed are fresh samples.

*Feather ϕ -keratin.

[†]Avian scale-type ϕ -keratin.

[§]Non-featherlike β -keratin.

[#] α -keratin.

TABLE 2. COMPARISON OF PYROLYSATES

Compound	White feather		Messel Shale		Fur Formation		Squid ink*	
	Fresh	Highly matured	Bat fur fossil	Sediment	Feather fossil	Sediment	Fossil	Modern
Acetic acid	X							
Toluene	X	X	?	?	?	?	X	X
2,3-epoxycarane					?			
N-[2-hydroxyethyl]succinimide		X						
2,5-diketopiperazine	X	X						
3,4-dimethyl benzaldehyde			X	X		?		
Dimethyl cyclopentene			?		X	?		
Dimethyl naphthalene			X	X	X	X		
<u>Various ketones</u>								
4-methyl-1-penten-3-one			?		?	?		
2-methyl cyclooctanone					X			
2-methyl cycloheptanone					X			
Acetophenone			?	X	?	X		
Piperidinone		X						
1-(2-vinylphenyl) ethanone					?	X		
2H-Inden-2-one, 1,3-dihydro-1-(1-oxopropoxy)-			?	?		?		
Benzophenone			X	?	X	X		
<u>Nitriles</u>								
Propanenitrile	X	X						
Methyl pentanenitrile		X						
Hexadecyl nitrile		X						
Docosenenyl nitrile		X						
2-methyl propanenitrile	X							
2-methyl butanenitrile	X							
Dimethyl benzonitrile			X	X				
2-phenyl acetone nitrile							X	X
3-phenyl propanenitrile							X	X
<u>Pyridines</u>								
Pyridine	X		?	?	?	?	X	X
Methyl pyridine	X						X	X
<u>Benzenes</u>								
Benzene		X	?	?		?		
Propyl cyanobenzene	X	X					X	
Ethyl benzene	X		X	X	?	X	X	
Ethyl cyanobenzene	X							
Propyl benzene			?	?	X	X		
<u>Amides</u>								
Acetamide		X						
Propanamide		X						
Methyl butanamide		X						
Hexanamide		X						
Methyl pentanamide		X						
Hexadecanamide		X						
Octadecanamide	X	X						
Eicosanamide		X						
Docosenamide		X						
Docosanamide		X						
<u>Pyrrole-related compounds</u>								
Ethyl pyrrole	X	X						
Pyrrolidine, 1-acetyl-		X						
Pyrrole	X						X	X
Methyl pyrrole	X		?	X		?	X	X
Dimethyl pyrrole			X	X				
Trimethyl pyrrole			?	?				
Ethyl-methyl-1H-pyrrole			X					
Ethyl-dimethyl-1H-pyrrole				?				
Ethylmethyl-1H-pyrrole-dione			X	?				

Thiophenes

Thiophene							X	
Methyl thiophene							X	
Ethyl thiophene							X	
Propyl thiophene			?	?	?	?	X	
Dimethyl thiophene			X	X				
Trimethyl thiophene			?			?		
Butyl thiophene			?	?	?	?		

Indoles

Indole	X		?	?		?	X	X
Methyl indole	X		?	?		?	X	X

Phenols

Phenol	X	X	X		X	X	X	X
Methyl phenol		X						
4-methyl phenol	X		?	?		?	X	X
5-methyl phenol							X	X
Ethyl (or dimethyl) phenol			?	?		?	X	X
Trimethyl methoxyphenol	X							
2-methyl phenol			?	?	?	?		
Di-ter-butyl phenol			?					

Styrene

Styrene	X	X	?	?		?		
Methyl styrene		X						

---

*from Glass et al. (2012).