Shakeel M, Jabeen F, Iqbal R, Chaudhry AS, Zafar S, Ali M, Khan MS, Khalid A, Shabbir S, Asghar MS.

[Assessment of Titanium Dioxide Nanoparticles \(TiO₂-NPs\) Induced Hepatotoxicity and Ameliorative Effects of *Cinnamomum cassia* in Sprague-Dawley Rats.](#)

Biological Trace Element Research 2017,
<https://doi.org/10.1007/s12011-017-1074-3>.

Copyright:

The final publication is available at Springer via <https://doi.org/10.1007/s12011-017-1074-3>.

Date deposited:

22/08/2017

Embargo release date:

19 June 2018

This work is licensed under a [Creative Commons Attribution-NonCommercial 3.0 Unported License](https://creativecommons.org/licenses/by-nc/3.0/)

Assessment of titanium dioxide nanoparticles (TiO₂-NPs) induced hepatotoxicity and ameliorative effects of *Cinnamomum cassia* in Sprague-Dawley rats

Muhammad Shakeel¹, Farhat Jabeen^{1*}, Rehana Iqbal², Abdul Shakoor Chaudhry³, Sadia Zafar⁴, Muhammad Ali¹, Muhammad Saleem Khan¹, Adeel Khalid¹, Samina Shabbir¹, Muhammad Saleem Asghar¹

1. Department of Zoology, Government College University, Faisalabad, Pakistan
2. Institute of Pure and Applied Biology, Bahauddin Zakariya University Multan
3. School of Agriculture, Food and Rural Development, Newcastle University UK
4. Department of Botany, Government College University, Faisalabad, Pakistan

ABSTRACT

This study assessed the protective effects of *Cinnamomum cassia* (cinnamon) bark extract in rats exposed to titanium dioxide nanoparticles or titanium dioxide bulk salt. For *in vivo* evaluation of the ameliorative role of the cinnamon extract, the experimental groups were orally administered with the cinnamon extract at different dose levels (50 or 100 or 150 mg/kg bodyweight) along with the subcutaneous injections of 150 mg/kg bodyweight titanium dioxide nanoparticles or titanium dioxide bulk salt. The extract showed significant ameliorative role on the antioxidant system in response to elevated levels of titanium dioxide nanoparticles or titanium dioxide bulk salt induced oxidative stress. It aided in the recovery of the antioxidant system as well as protective role in histological damages and some haematological parameters in the rat liver treated with titanium dioxide nanoparticles or titanium dioxide bulk salt.

Keywords: - TiO₂-NPs, TiO₂ Bulk Salt, Histopathology, hematology, Sprague Dawley rats

INTRODUCTION

Titanium dioxide (TiO₂) is one of the most commonly used materials and is among the top five nanoparticles being synthesized [1]. It is being used in nano-form in various industrial fields such as paints, pigments, dyes, alloys, rubber, ceramics, chemical fibres, sunscreens and other cosmetics, tooth pastes, electronics, pharmaceutical preparations, industrial photo-catalytic processes, personal care products, food additives, catalyst and metallurgy and in the environmental decontamination of air, soil, and water [2-9]. Due to the extensive application of TiO₂ nanoparticles (NPs) in the industrial field and on-going

commercialization of nanotechnology products, the exposure of the human body intentionally or unintentionally to nanoparticles via several possible routes, including dermal penetration, inhalation, oral ingestion or intravenous injection is possible and may continue to increase [3, 10]. Therefore, the analysis of its potential toxicity and distribution in the body, as well as the remedial activity of natural and easily available antioxidants is essential. Toxic effects of titanium dioxide nanoparticles and titanium dioxide bulk salt in the liver and blood of male Sprague-Dawley rats have been assessed by different assays in our previous study [11] and a number of studies regarding the toxicity due to TiO₂ nanoparticles in other body parts in different organisms have been reviewed by Shakeel et al. [12].

Drugs from natural sources are used for treating various diseases since the ancient times [13]. Plants have always been an ideal source of drugs. Many of the drugs that are currently available have been derived from the plants directly or indirectly. Medicinal plants are an integral or essential part of human society for the treatment of diseases from the beginning of civilization [14-16] and provide continued important therapeutic agents, in both the modern as well as traditional medicine [15, 17]. In recent times the utilization of medicinal plants belonging to different families is increasing because they are safe, effective, widely and readily available and inexpensive as compared to modern medicine [18-20]. At least 89% of Africans depend on herbal medicine for their healthcare [15, 21]. Because of the linked side effects of the modern medicine, the traditional medicines are gaining importance and are now being evaluated to discover the scientific basis of their healing activities [15, 22] in terms of their antioxidant, antimutagenic, and anticarcinogenic effects [23-25] and against a number of other diseases, such as anti-bacterial effect, since the plant products show fewer side effects and also may provide a way to the development of better new oral drugs [26, 27]. Nowadays, about 88% of the global populations turn to plant derived medicines as their first line of defence for their health maintenance and combating diseases [16, 19, 20, 26]. Plants possess a diverse variety of secondary metabolites, e.g., terpenoids, flavonoids, alkaloids, tannins, etc., that have been investigated *in vitro* to exhibit medicinal properties. About 120 secondary plant metabolites are used as drugs globally, 15% of all angiosperms have been evaluated chemically and of that 74% of pharmacologically active herbal components have been discovered [28]. Pharmacological studies have recognised the significance of medicinal plants as potential source of bioactive compounds [16, 19, 20, 26]. Extracts of some plants have been assessed in accordance with modern system of medicine and found to alter favourably the activities of several regulatory

enzymes [29]. Medicinal plants are known to exhibit their antioxidative effects by scavenging the reactive oxygen species (ROS) and modulating the antioxidant defence mechanisms [30, 31]. The world market for alternative medicine is being increased to approximately USD 10.895 billion according to a recent report by the World Health Organization (WHO) [32]. The global market for plant derived medicine alone has reached \$5 trillion and is increasing at a rate of about 11% per annum. This rapid growth in the market of the herbal/alternative medicines indicates its acceptance and popularity world-wide [32].

Cinnamon (*Cinnamomum cassia*, Family *Lauraceae*) bark is commonly used in Asian countries as a spice for most foods. In Eastern and Western folk medicine, it is used for treating abdominal and chest pains, chronic diarrhoea, hypertension, kidney disorders and rheumatism. It has strong antioxidant, analgesic, antiulcer, hypocholesterolaemic properties as well as antibacterial and anti-candidal, activities and regulates the lipogenesis [33-36].

To our knowledge no previous data are available on the Cinnamon (*C. cassia*) extract to combat metal induced toxicity in different organisms. Therefore, this study was designed to investigate the amelioration of the oxidative stress and hepatotoxicity induced by TiO₂ or its nanoparticles by Cinnamon extract in male Sprague-Dawley rats.

MATERIALS AND METHODS

The study was carried out at the Research Laboratory of the Department of Zoology, Government College University Faisalabad. The details of materials and methods are discussed below.

Animals

Thirty-five healthy post-weaning male Sprague-Dawley rats were procured from the animal house of Government College University Faisalabad, housed in groups in ventilated cages under standard lighting conditions and natural day/night cycle after approval from the local ethical committee of the Government College University Faisalabad. They were given a free access to water and food. Humidity and temperature (25°C ± 2°C) was controlled.

After a period of acclimatization for 7 days, the animals of similar mean initial body weights were randomly divided into groups each having five animals (Table 1). The treatment groups receiving the higher tested sub-lethal dose (150 mg/kg bodyweight) of

TiO₂ or TiO₂ nanoparticles were orally administered with the extract of Cinnamon to analyse its healing activities.

Table 1 Division of animals into different treatment groups.

Groups	Treatments
Group 1	Control (No treatment)
Group 1(a)	Placebo (Normal saline subcutaneously)
Group 2	TiO ₂ NPs 150 mg/kg BW subcutaneously + CE 50 mg/kg BW orally
Group 3	TiO ₂ NPs 150 mg/kg BW subcutaneously + CE 100 mg/kg BW orally
Group 4	TiO ₂ NPs 150 mg/kg BW subcutaneously + CE 150 mg/kg BW orally
Group 5	TiO ₂ BS 150 mg/kg BW subcutaneously + CE 50 mg/kg BW orally
Group 6	TiO ₂ BS 150 mg/kg BW subcutaneously + CE 100 mg/kg BW orally
Group 7	TiO ₂ BS 150 mg/kg BW subcutaneously + CE 150 mg/kg BW orally

A control group was fed by usual water and food while the other group was treated with normal saline subcutaneously for the equivalency of the shock due to the injection. The animals receiving 150mg/kg bodyweight TiO₂ NPs along with cinnamon extract at the dose of 50 or 100 or 150 mg/kg body weight orally were designated as Group 2, Group 3 and Group 4 respectively, whereas the animals receiving TiO₂ bulk salt at the dose of 150 mg/kg body weight along with cinnamon extract at the dose of 50 or 100 or 150 mg/kg body weight orally were designated as Group 5, Group 6 and Group 7, respectively. The treatment continued on alternate days for a period of 28 days. The dose of TiO₂ NPs and their bulk counterpart was selected on the basis of our previously determined sub-lethal dose at the rate of 150 mg/kg bodyweight of rat [11].

Sample Collection

Blood sample of all animals were collected at the start of the experiment and after 28 days of the treatment for the analysis of haematology, liver function tests, micronucleus assay and oxidative stress enzymes. At the end of the experimental period, animals were fastened overnight, anaesthetized the next day, by administering ketamine hydrochloride (30 mg/kg BW) and sacrificed. Blood samples were collected in heparinised tubes and plasma was separated by centrifugation at 2000xg for 10 minutes. The liver were collected, weighed with the help of sartorius weighing balance and were separately immersed in fixative sera for further process of histology (by haematoxylin–eosin staining method).

TiO₂ Nanoparticles

The titanium dioxide nanoparticles were synthesized using Sol-Gel method from Titanium Isopropoxide [TTIP; Ti(OC₃H₇)₄] and characterized using XRD, SEM and EDX. The details of the synthesis and characterization of the nanoparticles and the preparation of solutions have been described in our previously published article [11].

Preparation of Cinnamon Extracts

The cinnamon (*Cinnamomum cassia*) bark of the medicinal quality was purchased from a local homeo-store and its botanical position was confirmed from the botany department of Government College University Faisalabad. The cinnamon bark extract was prepared by following the method of plant extraction described by Khan et al. [37].

Body Weight

The body weights of the control group and all the experimental groups were observed and recorded weekly to note weekly changes in body weights.

Haematological Analysis

Blood samples were analysed using a haematology auto-analyser for the analysis of red blood cells (RBC) and white blood cells (WBC) counts, haemoglobin (Hb), platelets (PLT), neutrophils, monocytes, eosinophils, total leukocyte count (TLC), erythrocyte sedimentation rate (ESR), total protein, packed cell volume (PCV), erythrocyte indices like mean corpuscular value (MCV), mean corpuscular haemoglobin (MCH) and mean corpuscular haemoglobin concentration (MCHC).

Liver Function Tests

The liver function tests such as ALT (Alanine Aminotransaminase), AST (Aspartate Aminotransferase) and ALP (Alkaline Phosphatase) were performed to analyse the effects of TiO₂ or TiO₂ nanoparticles as well as the cinnamon extract on the liver using standard methods [11].

Micronucleus Assay

All the blood samples were coded and processed immediately after collection. Slides were prepared for the micronucleus test [11].

Analysis of Oxidative Stress

The activities of biomarkers of oxidative stress, i.e., catalase activity, superoxide dismutase activity, glutathione-S-transferase activity and lipid peroxidation, were measured to determine the oxidative stress [11].

Histological Examination

The animals were euthanized and fresh portions were cut rapidly from lateral lobes of the liver of each rat. The samples were prepared for light microscopic analysis for liver histology following the haematoxylin and eosin staining technique [11].

Statistical Analysis

Data were statistically analysed with the help of Minitab17 software using ANOVA in General Liner Model to determine the treatment effects on different parameters. The analysis compared the effect of the above-mentioned treatments on body weight, liver function tests, lipid peroxidation, oxidative stress and micronucleus assay parameters at $P < 0.05$. Tukey's test was used to compare treatments means at $P < 0.05$. Histology was assessed by recording visual observations of the photographic illustrations of liver tissues.

RESULTS

The ameliorating activities of *C. cassia* were assessed by analysing the haematology, micronucleus assay and the liver histology of the treated rats.

Characterization of the Synthesized TiO₂ Nanoparticles

XRD analysis of the nanoparticles showed the average crystallite size of the said nanoparticles as 32.12 nm. The SEM microphotograph for the synthesized TiO₂ nanoparticles showed the particles in the range of 30-80 nm [11].

Role of *C. cassia* as Antioxidant

The ameliorating activities of *C. cassia* against toxic effects caused by the sub-lethal doses of the subcutaneously administered TiO₂ nanoparticles as well as their bulk counterparts were demonstrated by analysing the haematology, liver function tests, micronucleus assay, oxidative stress enzymes as well as the histology of the treated rat.

Physiological Changes

Subcutaneous injection of 150 mg/kg TiO₂ or TiO₂ nanoparticles along with oral administration of 50 or 100 or 150 mg/kg *C. cassia* extract caused physiological changes. The rats exposed with 150 mg/kg of TiO₂ or TiO₂ NPs along with lower dose of extract (50 mg/kg) showed reduced weights and mortality of 2 animals during the last week of the experiment. The health status and behaviour of the animals with oral administration of 100 or 150 mg/kg of *C. cassia* extract along with 150 mg/kg TiO₂ or TiO₂ nanoparticles showed normal behaviour and their weights were increased. Table 2 shows significant increase in body weight in some groups and decreased weight in the other ones.

Haematological Analysis

Oral administration of *C. cassia* extract showed some ameliorative effects on the haematological alterations in rats caused by subcutaneous injection of TiO₂ or TiO₂ nanoparticles. At the start of the experiment, no significant changes were observed among all groups before administration of TiO₂ or TiO₂ NPs. Table 3 shows mean $\pm$ SD values of the haematological parameters among all groups at the start of the experiment.

Oral administration of *C. cassia* extract exhibited some ameliorative effects in rats after 28 days of exposure to TiO₂ or TiO₂ NPs as shown in Table 4.

Table 2. Weekly body weight (g) of control groups and treated groups exposed to TiO₂ nanoparticles along with oral administration of *C. cassia* extract for 28 days.

Groups Time interval	Group 1 (Control)	Group 1(a) (Placebo)	Group 2 (NPs + 50 mg/kg CE)	Group 3 (NPs + 100 mg/kg CE)	Group 4 (NPs + 150 mg/kg CE)	Group 5 (BS + 50 mg/kg CE)	Group 6 (BS + 100 mg/kg CE)	Group 7 (BS + 150 mg/kg CE)
Initial weight	56.0±0.6	54.6±0.9	57.3±0.9	54.0±0.6	47.0±0.6	67.0±0.6	53.0±0.6	45.0±1.2
1st week	56.0±0.6	58.0±1.2	51.0 ± 3.5	53.3±1.2	45.6±0.7	68.3±0.9	54.6±1.2	55.3±0.3
2nd week	64.3±2.6	62.2±1.3	43.6±1.2	54.0±1.2	50.3±1.8	77.0±1.2	60.3±0.9	60.3±1.8
3rd week	72.3±1.8	66.2±1.0	43.0±0.6	58.3±2.2	58.0±2.1	83.6±1.2	64.0±0.6	65.3±0.9
4th week	78.0±1.2	70.3±1.5	42.1±1.2	63.0±0.6	60.3±0.9	87.6±1.5	67.0±0.6	73.3±3.8

Values are as Mean±SD

Table 3 Mean±SD of hematological parameters among the control and the treated groups at the start of the experiment.

Parameters	Units	Group 1 (Control)	Group 1(a) (Placebo)	Group 2 (NPs + 50 mg/kg CE)	Group 3 (NPs + 100 mg/kg CE)	Group 4 (NPs + 150 mg/kg CE)	Group 5 (BS + 50 mg/kg CE)	Group 6 (BS + 100 mg/kg CE)	Group 7 (BS + 150 mg/kg CE)
MCV	(fl)	42.4±0.1	42.3±0.3	42.4±0.1	42.3±0.1	42.4±0.1	42.3±0.1	42.4±0.1	42.3±0.1
MCH	(pg)	13.5±0.3	13.4±0.2	13.6±0.4	13.2±0.2	13.4±0.2	13.7±0.2	13.5±0.2	13.5±0.4
MCHC	(g/dl)	17.4±0.2	17.3±0.1	17.5±0.2	17.4±0.2	17.7±0.2	17.4±0.2	17.7±0.2	17.3±0.2
HCT	%	21.5±0.1	21.6±0.2	21.5±0.1	21.4±0.1	21.5±0.1	21.4±0.1	21.4±0.1	21.4±0.1
Hb	(g/dl)	12.3±0.2	12.2±0.2	12.3±0.2	12.3±0.2	12.4±0.2	12.3±0.2	12.3±0.2	12.4±0.2
WBC	×10 ³ /μL	6.5±0.1	6.4±0.2	6.5±0.1	6.4±0.2	6.4±0.1	6.5±0.1	6.4±0.1	6.4±0.1
RBC	×10 ⁶ /Ml	6.5±0.02	6.6±0.1	6.4±0.02	6.4±0.1	6.4±0.02	6.5±0.02	6.4±0.01	6.4±0.01
LYM	%	42.3±0.1	42.1±0.3	42.3±0.2	42.3±0.02	42.3±0.1	42.2±0.1	42.3±0.2	42.3±0.2
PLT	×10 ³ /μL	178.2±0.1	178.3±0.02	178.2±0.1	178.2±0.2	178.2±0.1	178.2±0.1	178.2±0.1	178.2±0.1

Table 4 Mean±SD of haematological parameters among the control and the treated groups at the end of the experiment.

Parameters	Units	Group 1 (Control)	Group 1(a) (Placebo)	Group 2 (NPs + 50 mg/kg CE)	Group 3 (NPs + 100 mg/kg CE)	Group 4 (NPs + 150 mg/kg CE)	Group 5 (BS + 50 mg/kg CE)	Group 6 (BS + 100 mg/kg CE)	Group 7 (BS + 150 mg/kg CE)
MCV	(fl)	44.3±0.2 ^H	46.2±0.1 ^G	50.2±0.1 ^F	51.6±0.2 ^E	54.2±0.2 ^C	54.7±0.2 ^B	53.2±0.1 ^D	56.5±0.39 ^A
MCH	(pg)	15.5±0.2 ^H	16.2±0.1 ^G	17.5±0.2 ^F	18.3±0.2 ^E	19.2±0.1 ^C	19.6±0.2 ^B	18.5±0.39 ^D	20.6±0.2 ^A
MCHC	(g/dl)	28.5±0.2 ^F	27.5±0.4 ^G	33.23±0.2 ^E	35.5±0.2 ^C	35.2±0.1 ^D	35.5±0.4 ^C	35.6±0.2 ^B	36.3±0.2 ^A
HCT	%	30.2±0.1 ^G	28.5±0.2 ^H	33.47±0.3 ^E	35.3±0.2 ^D	36.6±0.2 ^B	35.7±0.2 ^C	37.2±0.1 ^A	33.6±0.2 ^F
Hb	(g/dl)	11.3±0.2 ^F	11.6±0.4 ^E	12.3±0.2 ^C	12.5±0.2 ^B	12.5±0.4 ^B	12.5±0.3 ^B	13.4±0.2 ^A	12.2±0.2 ^D
WBC	×10 ³ /μL	6.6±0.2 ^H	8.2±0.1 ^F	7.5±0.4 ^G	8.5±0.4 ^E	22.7±0.2 ^A	20.5±0.2 ^B	19.5±0.39 ^C	9.3±0.2 ^D
RBC	×10 ⁶ /Ml	6.8±0.02 ^B	6.6±0.02 ^C	5.34±0.02 ^F	6.8±0.02 ^B	6.8±0.02 ^B	6.2±0.6 ^D	6.98±0.01 ^A	5.9±0.02 ^E
LYM	%	38.2±0.1 ^E	37.1±0.1 ^F	31.2±0.1 ^H	35.2±0.1 ^G	66.5±0.2 ^D	72.2±0.2 ^B	68.5±0.2 ^C	96.5±0.3 ^A
PLT	×10 ³ /μL	177.0±0.1 ^G	170.0±0.1 ^H	1902.0±0.1 ^B	1920.0±0.1 ^A	1698.0±0.1 ^D	1592.0±2 ^E	1378.0±2 ^F	1817.0±2 ^C

Means that do not share a letter in the same row are significantly different

Liver Function Tests

Liver function tests such as ALT, AST and ALP were performed to study liver damage of control and treated groups. After 28 days of exposure the levels of ALT, AST and ALP were increased significantly at 150 mg/kg of TiO₂ nanoparticles and bulk salt TiO₂ along with low dose (50 mg/kg) of *C. cassia* extract while at highest dose (150 mg/kg) of *C. cassia* extract, the level of these enzymes gradually decreased. The oral administration of *C. cassia* significantly reduced the concentrations of ALT, AST and ALP. **Error! Reference source not found.** shows TiO₂ or TiO₂ nanoparticles induced detectable increase in ALT levels in rat's blood with oral administration of *C. cassia* extract at low dose but the higher dose of *C. cassia* extract reduced the ALT levels in blood. **Error! Reference source not found.** also indicate similar effects on the levels of AST in rat groups treated with TiO₂ or TiO₂ nanoparticles as well as with oral administration of *C. cassia* extract. The low doses of the *C. cassia* extract (50 mg/kg body weight of rats) was least effective in ameliorating the alterations in the concentrations of the AST induced due to the subcutaneously injected TiO₂ or TiO₂ nanoparticles, while the higher dose (150mg/kg body weight of rats) showed significant ameliorative effects on the blood levels of AST. It is also clear from the **Error! Reference source not found.** that the *C. cassia* extract showed decreasing effects on the blood levels of ALP in groups treated with high doses (150 mg/kg body weight of rat) of *C. cassia* extract, whereas, the low doses of *C. cassia* extract showed no obvious control on the TiO₂ or TiO₂ nanoparticles induced alterations in the ALP concentrations in blood.

Micronucleus Assay

Micronucleus assay was performed to study micronucleus (MN) frequency of control and treated groups. After 28 days of exposure the micronuclei frequency increased significantly at 150 mg/kg of TiO₂ nanoparticles or bulk-salt of TiO₂ with low dose of 50 mg/kg of *C. cassia* extract while at highest dose (150 mg/kg) of *C. cassia*, the micronuclei frequency was gradually decreased. Table 5 shows TiO₂ or TiO₂ nanoparticles induced detectable damage in rat's blood with oral administration of *C. cassia* extracts at low dose but the higher dose reduced the DNA damage due to the antioxidant property^H of *C. cassia*.

Table 5 Number of micronuclei after administration of different concentrations of *C. cassia* extract with TiO₂ or TiO₂ nanoparticles for 28 days.

Exposure Duration Groups	0 days	28 days
Group 1	0.21±0.02	0.30±0.57 ^H
Group 1(a)	0.23±0.02	0.41±0.21 ^G
Group 2	0.23±0.023	2.41±0.01 ^A
Group 3	0.29±0.05	2.12±1.01 ^C
Group 4	0.13±0.21	1.92±0.61 ^E
Group 5	0.21±0.02	2.20±0.11 ^B
Group 6	0.23±0.02	2.11±0.13 ^D
Group 7	0.23±0.02	1.80±0.09 ^F

Values as Mean ± SD for five replicates.

Histology

After the oral treatment of *C. cassia* extract to Sprague-Dawley rats along with the subcutaneous injections of TiO₂ or TiO₂ nanoparticles for 28 days, the livers of the test animals were evaluated for the histological studies.

Results showed that the extract had ameliorative effects on the histological damages caused by the TiO₂ bulk salt or the TiO₂ nanoparticles. The histological results are shown in **Error! Reference source not found.** for Group 1 (the control group receiving no treatments) and **Error! Reference source not found.** for Group 1(a) (the placebo group treated with only normal saline subcutaneously). **Error! Reference source not found.** to **Error! Reference source not found.** show the liver histology of rats treated with subcutaneous injections of sub-lethal dose of TiO₂ or TiO₂ nanoparticles and various concentrations of *C. cassia* extract.

Oxidative Stress Indicators

After the 28 days of treatment with TiO₂ or TiO₂ NPs, the level of CAT was found decreasing in liver and blood as compared to the control group (**Error! Reference source not found.**). However, the use of *C. cassia* increased the activity of CAT significantly ($P < 0.05$) as compared to the treated group of TiO₂ or TiO₂ NPs (**Error! Reference source not found.**).

TiO₂ or TiO₂ NPs significantly increased the activity of SOD after 28 days of treatment. However, the use of *C. cassia* ameliorated the oxidative stress and reduced the level of SOD in both liver and blood (**Error! Reference source not found.**).

The use of TiO₂ or TiO₂ NPs significantly increased the level of lipid peroxidation as compared to the control group. Oral administration of *C. cassia* significantly reduced the level of lipid peroxidation (**Error! Reference source not found.**).

DISCUSSION

Medicinal plants are being screened for the free radical scavenging properties based on the reports about their safety, efficacy and cost effectiveness [38-40] over the synthetic antioxidants that have been reported with side effects upon their consumption. As a result, a number of medicinal plants reported to have antioxidant properties are being recommended for pharmaceutical industries and traditional medicine to control and treat different types of ailments [41]. In recent times phytotherapy has gained popularity all over the world. Many plant parts possess phytochemical compounds which may be categorized into three major groups: the carotenoids, limonoids and flavonoids. Antioxidants interfere with oxidation reaction by reacting with free radicals, chelating catalytic metals, scavenging oxygen [42] and prevent lipid auto-oxidation [43, 44]. Researches have been performed worldwide for the potential of herbal medicine for their use in different fields of medicine. Use of herbal medicines exhibits positive effects in curing different diseases [45].

Cinnamon has been reported by a number of studies for having a reasonable amount of phenols and flavonoids. Polyphenolic and flavonoid compounds are important secondary metabolites in plants and are reported to be responsible for the variation in antioxidant activities in plants [46-48] and are capable of fighting against free radicals by inactivating lipid free radicals or preventing decomposition of hydrogen peroxide into free radicals due to their redox properties, chelate metal ions, quenching singlet and triplet oxygen [49-52].

This may undoubtedly suggest that plants with high quantity of polyphenols and flavonoids are good antioxidant sources.

Proximate analyses indicate that cinnamon powder has been found to be rich in fibre and carbohydrates and also contains appreciable amounts of protein and some minerals [53, 54]. Cinnamon powder has also been found to have high total phenolic content and antioxidant activity [53, 55]. Cinnamon contains complex chemical profile that includes phenolic compounds. The phenolic compound is general word and denotes to the compounds having at least one aromatic ring and one or extra attached hydroxyl group [56]. The phenolic compound possesses this capability due to the redox potential acting as reducing agent and reduces the singlet oxygen [57]. Flavonoid is considered one of most important phenolic compound with known properties and a significant role as free radical scavenging.

The current study investigated the ameliorative effects of *C. cassia* against the toxicity caused by the subcutaneously injected TiO₂ or TiO₂ nanoparticles in the body of albino rats. This study revealed the remedial or healing effects of *C. cassia* (Cinnamon) in rats exposed to TiO₂ sub-lethal doses. This study showed that TiO₂ and TiO₂ NPs exposure for 28days with oral administration of Cinnamon extract at dose of 100 or 150mg/kg was able to minimize the toxic effects of TiO₂ or TiO₂ NPs and induced healing effects in liver pathology, decreased the DNA damage and oxidative stress due to TiO₂ and TiO₂ NPs in albino rats. The findings of this study are in agreement with the results of many scientists who found that Cinnamon has anti-oxidant properties [33-35, 45, 58-67].

In this study, nearly normal haematological parameters were observed when cinnamon extract was administered along with the TiO₂ or TiO₂ nanoparticles. A number of other researches are in support to these findings. Cinnamon, a natural product with a long history of safety, is rich in polyphenolic components that have been revealed to improve the action of insulin *in vitro* [68], in animal researches [69, 70] and to possess *in vitro* antioxidant activity [71]. In this study results showed that the administration of cinnamon significantly improved SOD, CAT and GST and decreased lipid peroxidation in dose dependent manner. Similar findings were observed in a research by Khaki et al. [45] where the administration 75mg/kg *Cinnamon zeylanicum* caused significantly increase in serum SOD, CAT, GPX and decreasing MDA levels in rats. This results due to its flavonoids and related compounds. Researches confirmed that flavonoid especially Phytoestrogens have the potential to affect steroid biosynthesis and metabolism through a number of pathways [45, 72]. Markedly antioxidant activity of cinnamon was also shown

in liver tissue of the Wistar albino rats by [Noori et al. [33], Moselhy and Ali [61]] and Eidi et al. [73]. Normal and healing effects of cinnamon extracts have been demonstrated in a number of researches. Pretreatment of male Wistar rats with cinnamon extract considerably preserved the structure of hepatic cells and markedly reduced the hepatotoxicity caused by carbon tetrachloride (CCl₄) [61, 73] and oral treatment of cinnamon bark extract at the dose of 250 and 500 mg/kg BW to diabetic rats revealed completely repaired histology of the testis of the diabetic rats [74]. A recent study showed the ability of cinnamon to cure the tubular injury of kidney of female Wistar rats caused by Gentamicin apart from having phenolic compounds and antioxidant properties [75]. Oral administration of cinnamon oil significantly restored liver function and normalized the histological, histochemical and biochemical anomalies in alloxan treated diabetic rats, might be due to antioxidant properties [76].

In this study cinnamon treated rats also showed approximately normal values of some haematological parameters like RBC, Hb, HCT, MCH and MCHC supported by a number of studies. Ahmad et al. [77] reported recently no significant effects of 0.1 or 0.5 g/kg BW cinnamon extract on the behaviour, mortality, food consumption, water intake, weight gain, liver histology and haematological parameters in Sprague-Dawley rats. Cinnamon extract significantly protected the Hb, HCT, MCH and MCHC decrease in frog *Rana ridibunda* caused by CCl₄ [78]. Molla et al. [79] reported that 1 mL/litre cinnamon extract in drinking water caused no change in haematological parameters in broilers. Similarly 0.8% cinnamon increased the values of RBC and Hb concentrations [80]. However, the values WBC, LYM and PLT increased, that may be due to the provocation of the immune response. So, it is concluded that *Cinnamomum cassia* (Cinnamon) showed remedial or healing effects in rats exposed to TiO₂ sub-lethal doses as this plant is known for its anti-oxidative properties.

CONCLUSION

Oral administration of *C. cassia* (cinnamon) extract at doses of 100 or 150 mg/kg reduced the toxic effects of exposure with subcutaneous injection of TiO₂ or TiO₂ NPs in rat and reduced the liver injuries that was evident from the histological studies as well as the liver function enzymes concentrations. Therefore, cinnamon extract aided in the recovery of the antioxidant system as well as protective role in haematological and histological damages in the rat liver treated with TiO₂ or TiO₂ NPs.

Conflict of Interest

The authors declare that they have no conflict of interest.

Ethical Approval

All procedures performed in this study involving rats were in accordance with the research ethical standards of the Government College University Faisalabad, Pakistan.

REFERENCES

1. Shukla RK, Sharma V, Pandey AK, Singh S, Sultana S, Dhawan A (2011) ROS-mediated genotoxicity induced by titanium dioxide nanoparticles in human epidermal cells. *Toxicology in Vitro* 25(1):231-241. doi: <http://dx.doi.org/10.1016/j.tiv.2010.11.008>
2. Li N, Duan Y, Hong M, Zheng L, Fei M, Zhao X, Wang J, Cui Y, Liu H, Cai J, Gong S, Wang H, Hong F (2010) Spleen injury and apoptotic pathway in mice caused by titanium dioxide nanoparticles. *Toxicology Letters* 195(2–3):161-168. doi: <http://dx.doi.org/10.1016/j.toxlet.2010.03.1116>
3. Saman S, Moradhaseli S, Shokouhian A, Ghorbani M (2013) Histopathological effects of ZnO nanoparticles on liver and heart tissues in wistar rats. *Advances in BioResearch* 4(2):83-88.
4. Philbrook NA, Investigating the effects of nanoparticles on reproduction and development in *Drosophila melanogaster* and cd-1 mice. 2010, Queen's University, Kingston, Ontario, Canada. p. 91.
5. Pierzchala K, Oxidative stress on human cells in the presence of nano-sized titanium dioxide. 2010, École polytechnique fédérale de Lausanne EPFL.
6. Higarashi MM, Jardim WF (2002) Remediation of pesticide contaminated soil using TiO₂ mediated by solar light. *Catalysis Today* 76(2–4):201-207. doi: [http://dx.doi.org/10.1016/S0920-5861\(02\)00219-5](http://dx.doi.org/10.1016/S0920-5861(02)00219-5)
7. Balasubramanian G, Dionysiou DD, Suidan MT, Baudin I, Laine JM (2004) Evaluating the activities of immobilized TiO₂ powder films for the photocatalytic degradation of organic contaminants in water. *Applied Catalysis B: Environmental* 47(2):73-84. doi: <http://dx.doi.org/10.1016/j.apcatb.2003.04.002>
8. Konstantinou IK, Albanis TA (2004) TiO₂-assisted photocatalytic degradation of azo dyes in aqueous solution: kinetic and mechanistic investigations: A review. *Applied Catalysis B: Environmental* 49(1):1-14. doi: <http://dx.doi.org/10.1016/j.apcatb.2003.11.010>

9. Esterkin CR, Negro AC, Alfano OM, Cassano AE (2005) Air pollution remediation in a fixed bed photocatalytic reactor coated with TiO₂. *Journal of American Institute of Chemical Engineers* 51(8):2298-2310. doi: 10.1002/aic.10472
10. Zhang XD, Wu HY, Wu D, Wang YY, Chang JH, Zhai ZB, Meng AM, Liu PX, Zhang LA, Fan FY (2010) Toxicologic effects of gold nanoparticles *in vivo* by different administration routes. *International Journal of Nanomedicine* 5:771-781. doi: 10.2147/IJN.S8428
11. Shakeel M, Jabeen F, Qureshi NA, Fakhr-e-Alam M (2016) Toxic effects of titanium dioxide nanoparticles and titanium dioxide bulk salt in the liver and blood of male sprague-dawley rats assessed by different assays. *Biological Trace Element Research* 173(2):405-426. doi: 10.1007/s12011-016-0677-4
12. Shakeel M, Jabeen F, Shabbir S, Asghar MS, Khan MS, Chaudhry AS (2016) Toxicity of nano-titanium dioxide (TiO₂-NP) through various routes of exposure: a review. *Biological Trace Element Research* 172(1):1-36. doi: 10.1007/s12011-015-0550-x
13. Asif M (2012) Antimicrobial potential of *Azadirachta indica* against pathogenic bacteria and fungi. *Journal of Pharmacognosy and Phytochemistry* 1(4):78-83.
14. Nayak A, Nayak GNG, Soumya B, Bhat K, Kudalkar M (2011) Evaluation of antibacterial and anticandidial efficacy of aqueous and alcoholic extract of neem (*Azadirachta indica*) an *in vitro* study. *International Journal of Research in Ayurveda and Pharmacy* 2:230-235.
15. Bamidele FP, Ajibade AJ, Oyewo EB, Hannah AO (2013) A study of some effects of aqueous extract of neem (*Azadirachta indica*) leaves on the lead acetate induced neurotoxicity in the superficial layers of superior colliculus of adult wistar rats (*Rattus norvegicus*). *British Journal of Pharmaceutical Research* 3(2):217-231.
16. Biswas K, Chattopadhyay I, Banerjee RK, Bandyopadhyay U (2002) Biological activities and medicinal properties of neem (*Azadirachta indica*). *Current Science* 82(11):1336-1345.
17. Krentz A, Bailey C (2005) Oral antidiabetic agents. *Drugs* 65(3):385-411. doi: 10.2165/00003495-200565030-00005
18. Mousa MAA, El-Ashram AMM, Hamed M. Effect of neem leaf extract on freshwater fishes and zooplankton community. in *8th International Symposium on Tilapia in Aquaculture. The Central Laboratory for Aquaculture Research, Cairo, Egypt*. 2008.

19. Vasantharaj S, Sathiyavimal S, Hemashenpagam N (2013) Phytochemical analysis and antibacterial activity of *Azadirachta indica*. International Journal of Pharmaceutical Sciences Review and Research 22:59-61.
20. Mann A, Banso A, Clifford L (2008) An antifungal property of crude plant extracts from *Anogeissus leiocarpus* and *Terminalia avicennioides*. Tanzania Journal of Health Research 10(1):34-38. doi: <http://dx.doi.org/10.4314/thrb.v10i1.14339>
21. Elujoba AA, Odeleye OM, Ogunyemi CM (2005) Traditional medicine development for medical and dental primary health care delivery system in Africa. African Journal of Traditional, Complementary and Alternative Medicines 2(1):46-61.
22. Gupta YK, Briyal S (2004) Animal models of cerebral ischemia for evaluation of drugs. Indian Journal of Physiology and Pharmacology 48(4):379-394.
23. Khan N, Mukhtar H (2008) Multitargeted therapy of cancer by green tea polyphenols. Cancer Letters 269(2):269-280. doi: 10.1016/j.canlet.2008.04.014
24. Vinothini G, Manikandan P, Anandan R, Nagini S (2009) Chemoprevention of rat mammary carcinogenesis by *Azadirachta indica* leaf fractions: Modulation of hormone status, xenobiotic-metabolizing enzymes, oxidative stress, cell proliferation and apoptosis. Food and Chemical Toxicology 47(8):1852-1863. doi: <http://dx.doi.org/10.1016/j.fct.2009.04.045>
25. Abdel Moneim AE, Othman MS, Aref AM (2014) *Azadirachta indica* attenuates cisplatin-induced nephrotoxicity and oxidative stress. BioMed Research International 2014:11. doi: 10.1155/2014/647131
26. Vinoth B, Manivasagaperumal R, Rajaravindran M (2012) Phytochemical analysis and antibacterial activity of *Azadirachta indica* A. Juss. International Journal of Research in Plant Science 2(3):50-55.
27. Mahesh B, Satish S (2008) Antimicrobial activity of some important medicinal plant against plant and human pathogens. World Journal of Agricultural Sciences 4(5):839-843.
28. Raja RV, Ramanathan T, Savitha S (2009) Studies on wound healing property of coastal medicinal plants. Journal of Bioscience and Technology 1:39-44.
29. Shukia R, Sharma SB, Puri D, Prabhu KM, Murthy PS (2000) Medicinal plants for treatment of diabetes mellitus. Indian Journal of Clinical Biochemistry 15(Suppl 1):169-177. doi: 10.1007/BF02867556

30. Weisburger JH (2001) Antimutagenesis and anticarcinogenesis, from the past to the future. Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis 480–481:23-35. doi: [http://dx.doi.org/10.1016/S0027-5107\(01\)00166-X](http://dx.doi.org/10.1016/S0027-5107(01)00166-X)
31. Subapriya R, Nagini S (2003) Ethanolic neem leaf extract protects against N-methyl-N'-nitro-N-nitrosoguanidine-induced gastric carcinogenesis in Wistar rats. Asian Pac J Cancer Prev 4(3):215-23.
32. Paul R, Prasad M, Sah NK (2011) Anticancer biology of *Azadirachta indica* L (neem): A mini review. Cancer Biology & Therapy 12(6):467-476. doi: 10.4161/cbt.12.6.16850
33. Noori S, Azmat M, Mahboob T (2012) Study on antioxidant effects of cinnamon and garlic extract in liver, kidney and heart tissue of rat. Bioscience Research 9(1):17-22.
34. Qin B, Polansky M, Anderson R (2010) Cinnamon extract regulates plasma levels of adipose-derived factors and expression of multiple genes related to carbohydrate metabolism and lipogenesis in adipose tissue of fructose-fed rats. Hormone And Metabolic Research 42(3):187-193.
35. Ciftci M, Simsek UG, Yuce A, Yilmaz O, Dalkilic B (2010) Effects of dietary antibiotic and cinnamon oil supplementation on antioxidant enzyme activities, cholesterol levels and fatty acid compositions of serum and meat in broiler chickens. Acta Veterinaria Brno 79(1):33-40.
36. Peng X, Cheng KW, Ma J, Chen B, Ho CT, Lo C, Chen F, Wang M (2008) Cinnamon bark proanthocyanidins as reactive carbonyl scavengers to prevent the formation of advanced glycation endproducts. Journal of Agricultural and Food Chemistry 56(6):1907-1911. doi: 10.1021/jf073065v
37. Khan RA, Khan MR, Sahreen S (2012) Protective effect of *Sonchus asper* extracts against experimentally induced lung injuries in rats: A novel study. Experimental and Toxicologic Pathology 64(7–8):725-731. doi: <http://dx.doi.org/10.1016/j.etp.2011.01.007>
38. Ikpeme EV, Ekaluo UB, Kooffreh ME, Udensi O (2011) Phytochemistry and heamatological potential of ethanol seed leaf and pulp extracts of *Carica papaya* (Linn.). Pakistan Journal of Biological Sciences 14(6):408-411.

39. Ikpeme EV, Nta AI, Ekaluo UB, Udensi O (2012) Phytochemical screening and haematological evaluation of *Parkia biglobosa* and *Gongronema latifolium*. Journal of Basic and Applied Scientific Research 2(3):2599-2606.
40. Calixto BJB (2000) Efficacy, safety, quality control, marketing and regulatory guidelines for herbal medicines (phytotherapeutic agents). Brazilian Journal of Medical and Biological Research 33(2):179-189.
41. Ekaluo UB, Ikpeme EV, Udensi OU, Ekerette EE, Usen SO, Usoroh SF (2015) Comparative *in vitro* assessment of drumstick (*Moringa oleifera*) and neem (*azadiracta indica*) leaf extracts for antioxidant and free radical scavenging activities. Research Journal of Medicinal Plant 9(1):24-33.
42. Kelen M, Tepe B (2007) Screening of antioxidative properties and total phenolic compounds of various extracts of three different seed of grape varieties (*Vitis vinifera* L.) from Turkish flora. Pakistan Journal of Biological Sciences 10(3):403-408.
43. Brand-Williams W, Cuvelier ME, Berset C (1995) Use of a free radical method to evaluate antioxidant activity. Food Science and Technology 28(1):25-30. doi: [http://dx.doi.org/10.1016/S0023-6438\(95\)80008-5](http://dx.doi.org/10.1016/S0023-6438(95)80008-5)
44. Bondet V, Brand-Williams W, Berset C (1997) Kinetics and mechanisms of antioxidant activity using the DPPH^{*} free radical method. Food Science and Technology 30(6):609-615. doi: <http://dx.doi.org/10.1006/fstl.1997.0240>
45. Khaki A, Bayatmakoo R, Nouri M, Khaki AA (2013) Remedial effect of *Cinnamom zeylanicum* on serum anti-oxidants levels in male diabetic Rat. Life Science Journal 10(4s):103-107.
46. Basma AA, Zakaria Z, Latha LY, Sasidharan S (2011) Antioxidant activity and phytochemical screening of the methanol extracts of *Euphorbia hirta* L. Asian Pacific Journal of Tropical Medicine 4(5):386-390. doi: [http://dx.doi.org/10.1016/S1995-7645\(11\)60109-0](http://dx.doi.org/10.1016/S1995-7645(11)60109-0)
47. Uyoh EA, Chukwurah PN, David IA, Bassey AC (2013) Evaluation of antioxidant capacity of two ocimum species consumed locally as spices in nigeria as a justification for increased domestication. American Journal of Plant Sciences 4(2):222-230. doi: 10.4236/ajps.2013.42029
48. Demiray S, Pintado M, Castro P (2009) Evaluation of phenolic profiles and antioxidant activities of Turkish medicinal plants: *Tilia argentea*, *Crataegi folium*

- leaves and *Polygonum bistorta* roots. International Journal of Medical, Health, Biomedical, Bioengineering and Pharmaceutical Engineering 3(6):74-79.
49. Pokorný J, Yanishlieva N, Gordon M, Antioxidants in food: Practical applications. 2001, Cambridge England: CRC Press, Elsevier Science, Woodhead Publishing Limited.
 50. Maisuthisakul P, Suttajit M, Pongsawatmanit R (2007) Assessment of phenolic content and free radical-scavenging capacity of some Thai indigenous plants. Food Chemistry 100(4):1409-1418. doi: <http://dx.doi.org/10.1016/j.foodchem.2005.11.032>
 51. Balasundram N, Sundram K, Samman S (2006) Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses. Food Chemistry 99(1):191-203. doi: <http://dx.doi.org/10.1016/j.foodchem.2005.07.042>
 52. Javanmardi J, Stushnoff C, Locke E, Vivanco JM (2003) Antioxidant activity and total phenolic content of Iranian *Ocimum* accessions. Food Chemistry 83(4):547-550. doi: [http://dx.doi.org/10.1016/S0308-8146\(03\)00151-1](http://dx.doi.org/10.1016/S0308-8146(03)00151-1)
 53. Dhillon GK, Amarjeet K (2013) Quality evaluation of bread incorporated with different levels of cinnamon powder. International Journal of Food Science, Nutrition and Dietetics 2(7):70-74.
 54. Gul S, Safdar M (2009) Proximate composition and mineral analysis of cinnamon. Pakistan Journal of Nutrition 8(9):1456-1460.
 55. Shan B, Cai YZ, Sun M, Corke H (2005) Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. Journal of Agricultural and Food Chemistry 53(20):7749-7759.
 56. Mnayer D, Fabiano-Tixier AS, Petitcolas E, Hamieh T, Nehme N, Ferrant C, Fernandez X, Chemat F (2014) Chemical composition, antibacterial and antioxidant activities of six essential oils from the *Alliaceae* family. Molecules 19(12):20034-20053. doi: 10.3390/molecules191220034
 57. Atoui AK, Mansouri A, Boskou G, Kefalas P (2005) Tea and herbal infusions: Their antioxidant activity and phenolic profile. Food Chemistry 89(1):27-36. doi: <http://dx.doi.org/10.1016/j.foodchem.2004.01.075>
 58. Hashmat I, Azad H, Ahmed A (2012) Neem (*Azadirachta indica* A. Juss)-A nature's drugstore: An overview. International Research Journal of Biological Sciences 1:76-79.

59. Pankaj S, Lokeshwar T, Mukesh B, Vishnu B (2011) Review on neem (*Azadirachta indica*): thousand problems one solution. International Research Journal of Pharmacy 2(12):97-102.
60. Abubakar AA, Salka MN, Hassan FB (2011) Behaviour of male Wistar rats in response to different doses of extract of *Isobberlinia doka*. Asian Journal of Plant Science and Research 1(1):95-101.
61. Moselhy SS, Ali HKH (2009) Hepatoprotective effect of Cinnamon extracts against carbon tetrachloride induced oxidative stress and liver injury in rats. Biological Research 42:93-98.
62. Kim SH, Hyun SH, Choung SY (2006) Anti-diabetic effect of cinnamon extract on blood glucose in db/db mice. Journal of Ethnopharmacology 104(1–2):119-123. doi: <http://dx.doi.org/10.1016/j.jep.2005.08.059>
63. Kannappan S, Jayaraman T, Rajasekar P, Ravichandran M, Anuradha C (2006) Cinnamon bark extract improves glucose metabolism and lipid profile in the fructose-fed rat. Singapore Medical Journal 47(10):858-863.
64. Solomon TPJ, Blannin AK (2007) Effects of short-term cinnamon ingestion on *in vivo* glucose tolerance. Diabetes, Obesity and Metabolism 9(6):895-901. doi: 10.1111/j.1463-1326.2006.00694.x
65. Saglam O, Degirmenci I, Ustuner MC, Gunes HV (2012) The protective effects of cinnamon and sugar tea extract on diabetic rats with interrelationships between oxidative stress and DNA damage. African Journal of Pharmacy and Pharmacology 6(43):3012-3017.
66. Khaki A, Khaki AA, Hajhosseini L, Golzar FS, Ainehchi N (2014) The anti-oxidant effects of ginger and cinnamon on spermatogenesis dys-function of diabetes rats. African Journal of Traditional, Complementary and Alternative Medicines 11(4):1-8.
67. Elhassaneen YA, Saleh SA, El-Abd SF, El-Sayed MM, El-Nashar NN (2009) Dibutyl nitrosamine induces histopathological changes in rat: possible protective effects of cinnamon flavonoid extract. Journal of Pharmacology and Toxicology 4(2):58-69.
68. Anderson RA, Broadhurst CL, Polansky MM, Schmidt WF, Khan A, Flanagan VP, Schoene NW, Graves DJ (2004) Isolation and characterization of polyphenol type-A polymers from cinnamon with insulin-like biological activity. Journal of Agricultural and Food Chemistry 52(1):65-70. doi: 10.1021/jf034916b

69. Qin B, Nagasaki M, Ren M, Bajotto G, Oshida Y, Sato Y (2003) Cinnamon extract (traditional herb) potentiates *in vivo* insulin-regulated glucose utilization via enhancing insulin signaling in rats. *Diabetes Research and Clinical Practice* 62(3):139-148. doi: [http://dx.doi.org/10.1016/S0168-8227\(03\)00173-6](http://dx.doi.org/10.1016/S0168-8227(03)00173-6)
70. Qin B, Nagasaki M, Ren M, Bajotto G, Oshida Y, Sato Y (2004) Cinnamon extract prevents the insulin resistance induced by a high-fructose diet. *Hormone and Metabolic Research* 36(2):119-125.
71. Shobana S, Naidu KA (2000) Antioxidant activity of selected Indian spices. Prostaglandins, Leukotrienes and Essential Fatty Acids 62(2):107-110. doi: <http://dx.doi.org/10.1054/plef.1999.0128>
72. Whitten PL, Patisaul HB (2001) Cross-species and interassay comparisons of phytoestrogen action. *Environmental Health Perspectives* 109(Suppl 1):5-20.
73. Eidi A, Mortazavi P, Bazargan M, Zaringhalam J (2012) Hepatoprotective activity of cinnamon ethanolic extract against CCl₄-induced liver injury in rats. *Experimental and Clinical Sciences Journal* 11:495-507.
74. Shalaby MA, Mouneir SM (2010) Effect of *Zingiber officinale* roots and *Cinnamon zeylanicum* bark on fertility of male diabetic rats. *Global Veterinaria* 5(6):341-347.
75. Jahromi NA, Jahromi AR, Jahromi ZA, Jamali H, Tanomand S, Jahromi MA (2014) Cinnamon effect on kidney histological changes induced by gentamicin in rats. *International Journal of Advanced Biological and Biomedical Research* 2(4):1262-1267.
76. Mahera N, Shaikh A (2010) Histological and histochemical changes in diabetic male rat liver and intestine and protective effect of cinnamon oil. *Journal of the Faculty of Medicine Baghdad University* 52:366-371.
77. Ahmad RA, Serati-Nouri H, Abdul Majid FA, Sarmidi MR, Abdul Aziz R (2015) Assessment of potential toxicological effects of cinnamon bark aqueous extract in rats. *International Journal of Bioscience, Biochemistry and Bioinformatics* 5(1):36-44.
78. Al-Attar AM (2007) Chemopreventive effect of cinnamon extract on carbon tetrachloride-induced physiological changes in the frog, *Rana ridibunda*. *Journal of Applied Sciences* 7(1):79-90.
79. Molla M, Rahman M, Akter F, Mostofa M (2013) Effects of *Nishyinda*, black pepper and cinnamon extract as growth promoter in broilers. *Bangladesh Veterinarian* 29(2):69-77.

80. Najafi S, Taherpour K (2014) Effects of dietary ginger (*Zingiber officinale*), cinnamon (*Cinnamomum*), synbiotic and antibiotic supplementation on performance of broilers. Journal of Animal Science Advances 4(1):658-667.