Flynn SL, Gao Q, Robbins LJ, Warchola TJ, Weston JNJ, Alam MS, Liu Y,
Konhauser KO, Alessi DS.

[Measurements of bacterial mat metal binding capacity in alkaline and
carbonate-rich systems.](#)

Chemical Geology 2017, 451, 17-24.

Copyright:

© 2017. This manuscript version is made available under the [CC-BY-NC-ND 4.0 license](#)

DOI link to article:

<https://doi.org/10.1016/j.chemgeo.2017.01.002>

Date deposited:

19/10/2017

Embargo release date:

06 January 2018

This work is licensed under a
[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International licence](#)

Shannon L. Flynn^{1*}, Qiyang Gao^{1,2}, Leslie J. Robbins¹, Tyler J. Warchola¹, Johanna N.J. Weston^{1,3}, Md. Samrat Alam¹, Yuxia Liu¹, Kurt O. Konhauser¹, and Daniel S. Alessi¹

University of Alberta, Edmonton, AB T6G 2E3, Canada

University of Toronto, Toronto, ON M5S 3E5, Canada

University of Alberta Augustana Faculty, Camrose, AB T6B 1T3

Measuring the metal binding potential and reactivity of bacterial mats is challenging in alkaline and carbonate-rich systems. Traditional methods used to measure these parameters, such as potentiometric titrations and metal adsorption pH edges, are difficult to implement due to the presence of the carbonate minerals that buffer pH and prevent assessment of mat surface reactivity. Additionally, under alkaline conditions metals may form hydroxide and/or carbonate precipitates. In this study we examined the metal binding capacity of four distinct bacterial mats collected from Fairmont Hot Springs, BC, Canada. To prevent metal precipitation, the bacterial mat concentration was varied under a constant initial cadmium (Cd) concentration of 8.89 μM and at pH 8. In addition to the intact bacterial mats, a carbonate mineral sample and two bacterial mats in which the carbonate mineral was removed via acid-treatment, were used as end-members to assess the mechanisms of reactivity in the whole system. Freundlich adsorption isotherms were used to fit

metal adsorption data and directly compare surface reactivity amongst intact mats and mat components. Two of the intact mats exhibited a higher affinity for Cd compared to the mineral at metal equilibrium concentrations above 2.5 μM , while the other two intact mats had lower affinities under all experimental conditions. Generally, we found the acid-treated mats had higher Cd adsorption capacities than the carbonate mineral. When compared to their equivalent intact mats, only one acid-treated mat had a higher affinity for Cd. Further, we modelled whether metal adsorption in the intact mats, containing microbes and carbonate mineral, could be explained by a linear combination of the observed metal uptake by the organic and inorganic components through end-member experiments. Metal adsorption additivity results were mixed. Metal uptake by one intact mat was found to be additive, while for the other mat the additive model significantly underestimated the observed Cd accumulation. Our study demonstrates the potential, as well as the limitations, of using modified metal adsorption edges to determine the metal binding affinity and surface reactivity of bacterial mats in alkaline and carbonate-rich systems.

Introduction:

Bacteria are found in nearly every environment on Earth, and because of their ubiquity, they play an integral part in the cycling of most elements (see Konhauser, 2007 for details). The initial step in many biogeochemical processes is the adsorption of metal cations to the negatively-charged cellular surfaces (Beveridge and Murray, 1976). The adsorbed cations subsequently facilitate numerous chemical processes including metal reduction (e.g., Newman et al., 1997), oxidation (e.g., Nealson et al., 1988), intercellular accumulation (Southam and Beveridge, 1994), and in many cases, biomineralization (e.g., Ferris et al., 1986).

48 Significant work has been undertaken to determine the mechanisms by which metal cations
49 bind to bacteria (Mullen et al., 1989; Fein et al., 1997; Lalonde et al., 2007; Alessi and Fein, 2010).
50 Bacterial cell envelopes contain multiple surface functional groups that have distinct proton and
51 metal binding capacities (Beveridge and Murray, 1980; Fein et al., 1997). Typically,
52 potentiometric titrations of bacterial cells combined with data from bulk metal adsorption
53 experiments to the same bacteria have been used to develop protonation models and metal surface
54 complexation models (SCM), respectively (Cox et al., 1999; Fowle and Fein, 1999; Ngwenya et
55 al., 2003). For the SCM approach, the proton reactivity of a suspension of bacteria is tested by
56 potentiometric titration; that is, by adding precisely-measured aliquots of acid or base to the
57 suspension, and measuring the pH change due to the addition of that aliquot after equilibrium has
58 been achieved, across a wide pH range. The buffering capacity measured may then be ascribed to
59 a number of surface functional groups with discrete binding constants (pKa) and site
60 concentrations. Once the surface functional group concentrations and pKa value(s) have been
61 determined, bulk metal sorption experimental data from either metal adsorption pH edges (fixed
62 metal concentration with pH varied) or metal adsorption isotherms (fixed pH with metal
63 concentration varied) are fitted to determine the metal binding (equilibrium) constants onto the
64 proton-active functional groups calculated from the potentiometric titration data (Fein et al., 1997;
65 Cox et al., 1999; Haas et al., 2001; Ngwenya et al., 2003).

66 While this body of work demonstrates the potential of bacteria to bind and sequester metal
67 cations, most of these studies have focused on pure cultures of bacteria grown as suspensions under
68 carefully controlled laboratory conditions (e.g., Teitzel and Parsek, 2003). However, most bacteria
69 found in natural environments grow in dynamic multispecies communities, referred to as either
70 biofilms, or their thicker versions, bacterial mats (Kolter and Greenberg, 2006; Elias and Banin,

2012). Mats are also composed of dead cells and exopolymeric substances (EPS) which have been shown increase the number surface reactive sites, measured as mmol/g, compared to the intact bacterium itself (Baker et al., 2009). The secreted EPS have metal binding properties distinct from the bacterial cells and, therefore, they also impact the adsorption of metal cations from solution by bacterial mats. For example, Spath et al. (1998) measured the distribution of adsorbed cadmium (Cd^{2+}) and zinc (Zn^{2+}) on mats collected from wastewater treatment sequencing batch biofilm reactors. The EPS was found to retain 20% of the total sorbed metal cations. This indicates that there is a substantial difference between in the behavior of bacterial mats in nature, and the individual cells that make up the mats.

In addition to being composed of bacteria and EPS, mats often incorporate detrital and authigenic or allogenic minerals into their structure (e.g., Konhauser et al., 1998; Jones et al., 2004). These minerals have their own reactivity, which can alter the overall reactivity of the mats (Konhauser and Urrutia, 1999). For instance, Lalonde et al. (2007) applied a SCM approach to model the surface reactivity of intact mats from an alkaline hydrothermal spring in Yellowstone National Park. To determine the mat reactivity they performed acid titrations from pH 10 to 4, on both intact and acid treated mats. They found that the surface reactivity of the intact mats was dominated by a single surface site with a modeled pKa of approximately 7, which they attributed to the presence of incorporated carbonate precipitates. With the majority of carbonate precipitates removed, the acid treated mats exhibited an order of magnitude less buffering capacity (measured as mmol protons consumed per g mat) distributed across the tested pH range of 4 to 10. While that study was able to determine the relative buffering capacity of the mats, the authors never removed all of the carbonate minerals, meaning that even in their acid treated mat the carbonate minerals likely made a substantial contribution to the overall buffering capacity measured. Additionally,

94 since titrations were only preformed in a single direction, the reversibility of the system was
95 undetermined. If carbonate dissolution occurred during acid titration, a reverse titration (with base)
96 would have shown hysteresis in the buffering capacity between the forward and reverse titration
97 curves due to this loss of carbonate mineral. The study by Lalonde et al. (2007) highlights the
98 difficulties in determining reactivity in alkaline and carbonate systems and our inability to use
99 potentiometric titrations in the presence of carbonate precipitates or other pH sensitive precipitates.

100 In this study, we quantify the reactivity of four different bacterial mats collected from the
101 outflow of an alkaline hot spring located at Fairmont Hot Springs, British Columbia. Because of
102 the reasons described above, traditional methods to probe reactivity, such as potentiometric
103 titrations, adsorption isotherms, and adsorption pH edges, are likely to be incompatible due to
104 potential interferences caused by the rich abundance of carbonate minerals and the alkaline
105 conditions of Fairmont Hot Springs (Rainey and Jones, 2009). Based on the results of Lalonde et
106 al. (2007), the dissolution of carbonate minerals intertwined with the bacterial component of the
107 intact mats during acid titrations would likely dominate the proton buffering capacity and thereby
108 obscure direct measurement of the proton reactivity of the bacteria in the mat (Warchola et al., in
109 review), which is ultimately key in quantifying the contribution of mat bacteria to overall observed
110 metal removal from solution. Additionally, during metal adsorption pH edge experiments, the
111 presence of buffering carbonates would limit the pH range able to be tested. Methods such as metal
112 sorption edges, which are conducted under alkaline conditions, often create conditions in which
113 formation and precipitation of carbonate and hydroxide minerals are favorable. For example, in a
114 system in equilibrium with otavite (CdCO_3), the total Cd in solution decreases exponentially as a
115 function of increased pH (Figure 1). In this case Cd and otavite are used as proxies for divalent
116 metals and their carbonate precipitates, and clearly illustrate that under increasingly alkaline

conditions greater divalent cation precipitation can be expected. Accordingly, this precludes the investigation of metal adsorption using traditional metal adsorption isotherms for alkaline conditions.

As metal adsorption and the reactivity of bacterial mats are understudied in carbonate rich and alkaline systems, in this study, we propose a method to bypass the issues posed by traditional methods. To address these issues, modified metal adsorption experiments (in which Cd was used as a proxy for environmentally relevant divalent metals) were used to quantify the reactivity of the bacterial mats from a carbonate-rich system under metabolically inactive conditions. To minimize Cd precipitation, the concentration of sorbent (the bacterial mat or mineral) was varied with a constant initial Cd concentration in lieu of varying the metal concentration. Adsorption is thought to be a fast process occurring on a time scale of seconds to minutes (Xue et al., 1988; Kuyucak and Voelsky, 1989; Matheickal et al., 1999), whereas precipitation of mineral phases is the slower step on a time scale of hours to days (Sadiq, 1992). Therefore, the length of Cd exposure was limited to 4 h to allow for adsorption to reach equilibrium and to exclude significant precipitation of Cd as carbonates or hydroxides. In addition to determining the reactivity of intact bacterial mats, reactivity was also determined for travertine (a lithified carbonate sediment associated with microbial mats in hot springs) and two of the bacterial mats that were acid treated to remove the incorporated carbonate minerals. The reactivity of the intact bacterial mats, endmember mineral component, and pure bacterial components (acid treated mats) was quantified through the Cd sequestration potential. By quantifying the potential of not just the intact mats but each endmember component, the contribution of each endmember to the overall observed Cd sequestration in intact mat samples was assessed.

2.0 Methods:

2.1 Handling and Collection of Samples

Bacterial mats and travertine samples were collected from an alkaline hot spring located at Fairmont Hot Springs, BC, Canada on September 2, 2015. Material was collected from four distinct bacterial mats, subsequently referred to as: black mat, green mat, spongy green mat, and orange mat (Figure SI.1). The mats were removed from the underlying travertine and placed in polypropylene tubes submerged in water from the hot spring. In addition to bacterial mat and travertine (mineral) samples, water samples were collected for metals analysis. The pH and temperature were measured during sample collection using a Fisher Science Accumet AP71 portable dual pH and temperature meter along the hot spring flow path (locations shown in Figure SI.1). Water samples were filtered through 0.22 μm nylon membranes and acidified with three drops of 12.1 M HCl for preservation. All samples were placed on ice before transport to the University of Alberta. Samples were stored at 2° C prior to analyses.

2.2 Preparation of Travertine Sample

A travertine mineral sample, which was overlain by the bacterial mats in the field, was collected for reactivity/adsorption experiments because it represents the mineral endmember associated with the bacterial mats. Following collection, the travertine was dried in an oven at 60° C for 24 h, then crushed and ground into a fine powder using a mortar and pestle. The powdered travertine was treated to remove any trace organics that may have been incorporated in the travertine using a modified version of the separation method described in Schofield et al. (2008). In summary, the powdered travertine was soaked in 1 M NaOH for 24 h before being repeatedly washed with 18.2 M Ω ultrapure water and centrifuged at 7,000 g; this process was repeated five times. Following

washing, the travertine-water slurry was placed in a separatory funnel with hexane, shaken vigorously and left to separate overnight. The following morning the travertine powder and water was separated from the organics and hexanes, as the latter floated on top as a separate layer. The organics-free travertine was washed with 18.2 MΩ ultrapure water once more before being centrifuged and decanted twice at 7,000 g for 5 min, and a final time for 30 min to form a wet pellet. These wet pellets were subsequently used for metal adsorption experiments and a portion of the organics-free travertine was dried. The dried travertine, with a wet to dry ratio of 1.45 (SI Table 1), was analyzed for mineral composition and surface area using X-ray powder diffraction (XRD) and Brunauer-Emmett-Teller surface area analyzer (BET), respectively.

2.3 XRD

XRD analysis of the powdered mineral sample was performed using a Rigaku Geigerflex Powder X-ray diffractometer equipped with a cobalt tube, and graphite monochromator and scintillation detector. Data were analyzed using the JADE 9.1 software, with peak locations being referenced using the International Centre for Diffraction Data (ICDD) and Inorganic Crystal Structure Databases (ICSD).

2.4 Preparation of Bacterial Mat

All four bacterial mat samples were washed and centrifuged 3 times for 5 min at 7,000 g using a prepared stock a solution of 10 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) buffer and 10 mM NaCl adjusted to pH 8. The bacteria were then suspended in approximately 100 ml of the previously described solution on a magnetic stir plate and pH adjusted using small

185 aliquots of 1 M NaOH until the solution pH stabilized at 8 (the approximate ambient pH of the hot
186 spring). Bacterial mats were then pelleted by centrifuging and decanting two times for 5 min and
187 two times for 30 min at 7,000 g to produce a compressed wet pellet from each mat. A subsection
188 of each mat was subsequently acid treated with a solution of 0.1 M nitric acid and 10 mM NaCl to
189 remove carbonate minerals that may have been associated with the bacterial mats. Acid treated
190 bacterial cells have been found to remain intact and viable without any visible difference from
191 their untreated counterparts in previous studies (Fein et al., 1997; Borrok et al., 2004). The acid
192 treatment was performed on a stir plate with continual additions of the 0.1 M nitric acid plus 10
193 mM NaCl solution, until the pH of the bacterial mat suspension had stabilized at or below pH 4.5.
194 The acid treated mats were then pelleted using the method described above (Section 2.2). Two of
195 the acid treated mats (green and orange) were further treated with the previously described HEPES
196 stock solution until the pH of the bacterial mat suspension stabilized at 8. The acid treated mats
197 were then pelleted using the method described above, and are referred to here as ‘acid treated green
198 mat’ and ‘acid treated orange mat’. The mass difference between the pre- and post-acid treated
199 bacterial mat was used to determine the relative bacterial and mineral content of the intact bacterial
200 mats.

202 2.5 Cd Adsorption Experiments

203 Batch Cd adsorption experiments were conducted at room temperature (20°C) and pressure in a
204 matrix of 10 mM HEPES and 10 mM NaCl, adjusted to pH 8. As no nutrients or electron donors
205 were added to the experimental solutions, and the contact time in metal adsorption experiments
206 was short (4 h), the role of potential bacterial metabolism was assumed to be minimal. In contrast
207 to traditional adsorption experiments where the concentration of metal is varied, the concentration

of Cd added was held constant at 8.89 μM (1 ppm) while the sorbent concentration was varied from 0 to 8 g/L for the travertine and 0 to 25 g/L for the bacterial mats, based on wet weight from the pelleted material. Duplicate experiments were conducted on the travertine, four intact bacterial mats, and two acid treated bacterial mats. In preparation for the experiments, bacterial mat or travertine samples were first added to the solution in polypropylene test tubes; the pH was then checked and adjusted if needed before the solution was spiked with Cd. Test tubes were rotated end-over-end at 100 rpm for 4 h, adequate time for Cd adsorption and equilibrium to be achieved but short enough to prevent significant precipitation (<1.5%, see control experiment in Figure SI.3). The pH of the solution was measured and adjusted as needed over the first hours of the experiment. At the end of the adsorption experiment, the pH of each sample was measured and recorded, the tubes centrifuged at 5,000 g for 5 min, the resulting supernatant subsequently filtered through a 0.2 μm nylon membrane, and then diluted 1:10 with 2% nitric acid before Cd analysis.

2.6 ICP-MS Analysis

Metal analyses of hot spring water and Cd from adsorption experiments were conducted at the Environmental Geochemistry Laboratory at the University of Alberta using an Agilent 8800 ICP-MS/MS. A detailed table of the analysis conditions and dilution factors used is provided in Table SI.2.

3. Results

3.1 Hot Spring Water, Travertine and Bacteria Mat Mineral Analysis

Water samples along the flow path were found to show no systematic variation in measured chemical composition, while the pH increased 1.04 units along the sampled transect and the

temperature decreased by 1.9 °C (Table SI.3). In total 14 cationic and 1 anionic elements were found to be in quantifiable concentration in hot spring water by ICP-MS/MS analysis. The primary cations in the hot spring water were Ca and Mg, with Na, K, Si, and Sr also being present in the ppm range; similar to concentrations reported by Warchola et al. (in review) for the same transect. The remaining 8 elemental concentrations were found to be in the ppb range (Table SI.3). Anions were not measured but were assumed to be similar to previous measurements from the same site by Warchola et al. (in review), in which the anions were dominated by alkalinity in the form of carbonate (380-400 ppm), sulphate (230 ppm), and chloride (26 ppm) with a range of ionic strengths between 12-14 mM.

The XRD analysis of powdered travertine mineral from the hot springs was found to be composed of calcite, quartz, dolomite and rutile (Figure SI.2). The calcite likely precipitated in response to the supersaturated state of the solution with respect to $p\text{CO}_2$ in water emerging from the subsurface. The dolomite is likely the product of alteration and Mg exchange in a portion of the calcite. The presence of quartz is likely the result of detrital sand grains entrapped in the carbonate travertine. Rutile, a common paint additive, could be from the inclusion of paint chips washed from the thermal pools at the resort above the sampling site.

The carbonate mineral content of each bacterial mat was determined through acid treatment (Table 1.). The orange mat had the lowest carbonate mineral content, with 32.2% carbonate minerals by mass. The black and spongy green mats were found to have similar carbonate mineral contents to the orange mat with 36.5% and 39.8%, respectively. The green mat was composed of significantly more carbonate minerals with 72.8% of the mat being composed of carbonates, nearly double the content of the other bacterial mats.

253

254 3.2 Cd Adsorption Experiments

255 3.2.1 Travertine

256 The amount of Cd adsorbed from solution increased with increasing concentration of travertine
257 particles until complete removal of Cd was achieved at approximately 7.7 g/L (Figure 2). From a
258 concentration of 0.06 g/L of travertine to 2.0 g/L, a rapid and near linear increase in the adsorption
259 of Cd was observed. The extent of Cd adsorbed per g of travertine decreased from 2.0 g/L until
260 7.7 g/L of travertine where less than 1% of Cd was measured to be remaining in solution. Above
261 7.7 g/L of travertine, near-complete removal of solution Cd was observed.

262

263 3.2.2 Bacterial Mats

264 3.2.2.1 *Green Mat*

265 Cd adsorption increased with increasing concentration of green mat until the percent adsorbed
266 plateaued above 15.8 g/L of bacterial mat (Figure 3.A). At this concentration or greater, the percent
267 Cd adsorbed was found to be 90-95% of the total Cd regardless of increasing concentration of
268 bacterial mat, potentially due to the presence of organic acids in solution that complex Cd. Initially
269 the extent of Cd adsorption, from 0.041 to 1.2 g/L of bacterial mat, was rapid and linear in increase
270 relative to the increase in bacterial mat added. Once the green mat concentration was greater than
271 1.2 g/L, and until 15.8 g/L, the extent decreased before the percent adsorbed plateaued.

272

273 3.2.2.2 *Black Mat*

274 A linear and rapid increase in adsorbed Cd was observed with increasing black mat concentrations
275 from 0.069 to 1.13 g/L (Figure 3.B). Above 1.13 g/L of bacterial mat the extent of Cd adsorption

per g/L decreased with increasing bacterial mat until 15.9 g/L of black mat. Above 15.9 g/L of bacterial mat, the percent of Cd adsorbed plateaus at 82-92% despite increasing bacterial mat concentration.

3.2.2.3 Spongy Green Mat

The spongy green mat showed a sharp and linear increase in Cd adsorbed with increasing bacterial mat concentration from 0.062 to 1.06 g/L (Figure 3.C). The extent of adsorption decreased with increasing spongy green mat concentration from 1.06 to 5.11 g/L bacterial mat before the percent adsorbed plateaus at 81-92% even with increasing bacterial mat concentrations.

3.2.2.4 Orange Mat

Cd adsorption to the orange mat increased with increasing mat concentrations until plateauing above 11.00 g/L (Figure 3.D). The maximum adsorbed Cd was found to be between 95-98% of the initial concentration. At lower concentrations of the orange mat, from 0.048 to 1.12 g/L, the extent of adsorption is rapid and linear. From 1.12 to 11.00 g/L the extent of Cd adsorption decreases with increasing mat concentrations.

3.2.3 Acid Treated Bacterial Mats

3.2.3.1 Acid Treated Green Mat

A linear increase in Cd adsorption was observed with increasing acid treated green mat from 0.069 to 0.48 g/L (Figure 4.A). Above 0.48 to 0.99 g/L of bacterial mat, the extent of Cd adsorption decreased with increasing bacterial mat concentration. After 0.99 g/L, the percent of Cd adsorbed plateaus at 97 to 100% even with continuing additions of bacterial mat.

299

300 3.2.3.4 Acid Treated Orange Mat

301 The acid treated orange mat showed a rapid and linear increase in the percent of Cd adsorbed with
302 increasing mat concentrations from 0.034 to 0.98 g/L (Figure 4.B). The extent of increased
303 adsorption per g/L of bacterial mat decreased from 0.98 to 8.10 g/L of bacterial mat. Once the
304 concentration of bacterial mat reached 8.10 g/L the adsorption of Cd ranged between 91-100% of
305 the initial Cd concentration.

306

307 4. Discussion

308 For this study, metal adsorption to travertine, bacterial mats, and acid treated bacterial mats was
309 modeled using; (1) Freundlich isotherms as a means of determining the relative metal binding
310 affinity, and (2) linear combination fits of the endmember component contributions to the intact
311 mat Cd adsorption capacity as a means of testing the additivity of metal adsorption in the intact
312 bacterial mats. Freundlich isotherms have been successfully employed to quantify the adsorption
313 capacity of numerous biosorbents, such as marine algae (Kuyucak and Volesky, 1989), fungi
314 (Mullens et al., 1992), and bacteria (Mullens et al., 1989; Volesky and Holan, 1995). The combined
315 experimental data for the duplicate experiments for each sorbent was fitted together using the
316 linearized Freundlich isotherm equation:

$$317 \quad \text{Log}(C^*) = \text{Log}(K) + N * \text{Log}(C) \quad (1)$$

318 in which C and C* represent the equilibrium Cd concentration in solution (μm) and the adsorbed
319 Cd concentration ($\mu\text{m/g}$), respectively, and N and K are Freundlich constants. In this form, the K
320 constant represents the affinity of the sorbent for Cd in $\mu\text{m/g}$ at an equilibrium solution
321 concentration of 1 μm . The linearized Freundlich equation fits the Cd adsorption data for

environmental samples well (Table 2), with coefficient of determination values ranging from 0.86 to 0.96 (insets in Figures 2, 3A-D, and 4A-B).

At low metal concentrations, those below 1 μM equilibrium Cd in solution, travertine shows the highest modeled affinity for Cd (See Figure 5). The second highest affinity in this range was predicted to be for the acid treated green mat, which along with travertine at 1 μM , had an affinity that was double that of all other sorbents. Above 1 μM the affinity of travertine for Cd increases more slowly relative to the other sorbents, as indicated by the corresponding N values (Table 2). Once the equilibrium Cd solution concentration reaches 7.5 μM , the affinity of travertine for Cd is only greater than that of the intact black mat, and at or above 10 μM equilibrium Cd in solution travertine has the lowest overall Cd affinity.

Among the intact mats, which include both the microbial biomass and carbonate solids, the orange mat has the highest Cd affinity at equilibrium concentration below 3 μM ; above 3 μM the green mat has the highest affinity with the orange mat having the second highest. The spongy green and black mats are predicted by the models to have lower affinity for Cd than the green and orange mats throughout the entire examined Cd concentration range. Of these two intact mats, the black mat is predicted to have the higher affinity for Cd when the metal concentration is below 4.5 μM , and vice versa above 4.5 μM .

Of the two acid treated mats tested, the green mat has a higher predicted affinity for Cd over the entire range examined. When compared to the intact green mat, the acid treated green mat was found to have approximately twice the affinity for Cd. In contrast the acid treated orange mat exhibited lower Cd affinity, approximately half that of the intact orange mat.

To determine whether metal adsorption by the intact mats is additive (that is, if it can be modelled as the sum of its constituent components), a component additivity modeling approach utilizing acid treated mat (biomass only) and travertine-only endmembers was applied to measured Cd adsorption data from the corresponding intact mats. Using this approach, it can be determined whether Cd adsorption to the components of the intact mats is additive, or if when combined, the sum of the Cd adsorption to the individual components show synergistic or antagonistic relationships that impact overall Cd removal from solution (Alessi and Fein, 2010). A composite model composed of the sum of the adsorption capacity of individual mat components, similar to that used by Small et al. (1999) for bacteria-Fe(III) composites, was used to determine the additivity of the bacterial mats. The composite Cd removal from solution ($C^*_{\text{Comp.}}$), which represents the predicted Cd removal by intact mats, was calculated as the sum of the intact mat components, according to:

$$\% \text{Bac.} * C^*_{\text{Bac.}} + \% \text{Min.} * C^*_{\text{Min.}} = C^*_{\text{Comp.}} \quad (2)$$

in which the sum of the predicted component Cd affinity, $C^*_{\text{Bac.}}$ (acid treated bacterial mat) and $C^*_{\text{Min.}}$ (travertine) are multiplied by their respective mass percent in the intact bacterial mat. The calculated Cd removal ($C^*_{\text{Comp.}}$) was then substituted for C^* in Equation 1 to calculate the hypothetical composite model Cd removal across a range of Cd concentrations. Overall, the composite model for the green mat reasonably matches the Freundlich fit to the intact green mat Cd adsorption data (Table 2 and Figure SI.4.A,B), but overestimates affinity for Cd at low concentrations (below 6 μM) and underestimates mat Cd affinity at higher equilibrium Cd concentrations in solution above 6.5 μM . The composite orange mat underestimates the adsorbed Cd concentration once the equilibrium solution Cd concentration is above 0.95 μM , or across the majority of the experimental range.

As a second test of the component additivity of the system, a subtractive model (equation 3), was created to assess if the adsorption behavior of travertine and an intact mat could be used to calculate the adsorption behavior of the bacteria-only component of that mat, as represented by the acid treated mat. In this model, the total concentration of Cd adsorbed to the intact mat ($C^*_{Tot.}$) is subtracted by the product of the percent travertine by mass (%Min.) in the mat and the concentration adsorbed to the travertine ($C^*_{Min.}$), and then divided by the percent mass of bacteria in the intact mat determined through acid treatment (%Bac.), giving the concentration of Cd that should be adsorbed on the bacterial component of the mat ($C^*_{Bac.}$), according to:

$$\frac{C^*_{Tot.} - \%Min. * C^*_{Min.}}{\%Bac.} = C^*_{Bac.} \quad (3)$$

The subtractive models fitting parameters are shown in 2 and Figure SI.5. The subtractive model for the green mat overestimates the affinity of the acid treated green mat for Cd, once the equilibrium solution Cd concentration is above 3.0 μ M. In contrast, the subtractive model for the orange mat overestimates Cd adsorption onto the bacterial component of acid treated orange mat over the entire experimental range. The overestimation for both the subtractive green and orange mats could be caused by a difference in the size of reactive mineral particles used as the inorganic endmember and particle size found in the mats. If the particles found in the mats were smaller than those used as the mineral endmember, then the mineral contribution would have likely been underestimated due to the difference in the reactive surface area. Another possibility is that Ca and Mg are released during travertine dissolution in the intact mats. Then, the Ca and Mg compete with Cd for adsorption to surface sites within the bacterial components of the mat. The competition is expected to be minimal as binding constants for Ca^{2+} and Mg^{2+} to the model bacterium *Bacillus subtilis* are two orders of magnitude lower than that for the binding of Cd^{2+} (Flynn et al., 2014). This underestimation would then result in the overestimation of the bacterial component

contribution, as our models are too simplistic to represent the true complexity of the mat. The subtractive models for the spongy green and black mats were found to have higher N values than those for the green and orange mats, however, the K values were significantly lower. Over the range examined, the bacterial component of the green mat is modeled to have the highest Cd metal sorption followed by the bacterial component in the orange, spongy green, and black mats.

In alkaline and carbonate systems, metals can be incorporated into carbonate precipitates by either abiotic or biotic processes. Abiotic partitioning of alkali earth elements into carbonates has been found to be more significant into aragonite than calcite, with incorporation increasing with decreased ionic radius (Okumura and Yasushi, 1986). Additionally, the partitioning of divalent metals into carbonate minerals is affected by their depth in the water column (Elderfield et al., 1996) and divalent metals have been shown to coordinate at Ca sites on calcite, forming mononuclear inner-sphere adsorption complexes (Elzinga and Reeder, 2002). At present it is not known whether the same mechanism of trace metal incorporation into carbonates minerals occurs via biotic processes. However, the first step is likely adsorption of Ca^{2+} ions to the negatively charged bacterial surface sites or negatively charged bacterial mat surface sites (a combination of bacterial surface sites and EPS surface sites) (Thompson and Ferris, 1990; Braissant et al., 2007). The surface sites act as nucleation points from which carbonate minerals can grow using the bacterial surface as a template. Indeed, a number of different carbonate phases can form by simply replacing Ca for other divalent cations in the growth media, e.g., Fe^{2+} to form siderite (Mortimer et al., 1997) and Sr^{2+} to form strontianite (Schultze-Lam and Beveridge, 1994). In addition to the Ca being incorporated into the biogenic carbonate precipitates, other metals (i.e., Cd, Cu, Zn, Pb, REEs, Sr) bound to the bacterial mats could also be incorporated into the carbonate precipitates nucleating on the bacterial mat. For instance, previous studies have documented that cyanobacteria

can partition of up to 1.0 wt% strontium in calcite (Ferris et al., 1995). This has important implications for trace metal incorporation into biogenic minerals in the environment because passive carbonate biomineralization may provide a useful strategy for contamination issues in Ca carbonate-rich aquifers (Warren et al., 2001).

This study demonstrates the possibility to probe and measure metal reactivity in alkaline- and carbonate-rich systems by altering traditional metal adsorption experiments – using a low and consistent initial metal concentration, varying the bacterial mat concentration, and shortening the exposure time. While this method may be expanded upon or altered to elucidate additional information about the bacterial mat reactivity, our modified metal adsorption experiments have shown the ability to determine the relative Cd affinity of several bacterial mats from a hot spring under varied metal loading conditions, and provides a first step in determining which of the metal immobilization mechanisms discussed above may be important in alkaline and carbonate-rich systems. Differences in the metal binding between the mats likely was due to the differences in the ratio of biological material to travertine/mineral phases and the ratio of bacterial cells to EPS present in the mats. The reactivity of the bacteria within the mats is not thought to be strongly influenced by the species of bacteria present, as previous work by Borrok et al. (2005) found remarkable similarities in the reactivity of 36 separate bacterial species and consortia. Our method showed mixed results in determining the relative additivity of mat biological and inorganic components toward the total reactivity of the intact mat. The mixed results with regards to system additivity may have stemmed from a variety of differences in the mats, including density of the biological material, diffusion of Cd through the mat, and detrital inorganic grain size. Further work will be needed to determine whether adsorption of metals to bacterial surface functional groups is the precursory step for metal incorporation into biogenic carbonates. Future studies are also needed

to explore how the adsorption of metal onto bacterial mats affects metal incorporation to microbially-induced carbonates, and to assess the relative contribution of biotic or abiotic mechanisms to the formation of microbial mat carbonates. By continuing to study bacterial mat reactivity in alkaline- and carbonate-rich systems, we may be able to determine these controls on metal partitioning into biogenic carbonates, and thereby further our understanding of metal cycling and sequestration in relevant modern environments, ranging from hot springs to oceans.

5. Conclusions

This study demonstrates the potential and limitations of using modified metal adsorption edges to determine the metal binding affinity and surface reactivity of metabolically inactive bacteria in alkaline and carbonate-rich systems, such as in bacterial mats that have authigenic carbonate minerals. The empirical models were effective in determining which of the Fairmont Hot Springs' bacterial mats had the highest metal adsorption potential. However, the results were mixed when the additivity of the bacterial mat components, mineral, and bacteria, were measured separately and the sum of their relative contributions used to create additive models. This study highlights the potential as well as limitations of using modified metal adsorption edges to determine the metal binding affinity and surface reactivity of natural bacteria in alkaline and carbonate-rich systems.

Models such as the ones presented in this paper expand our understanding of bacterial mat reactivity in alkaline and carbonate rich systems. Further studies are required to transition from modeling reactivity in these complex systems with empirical models, such as Freundlich or Langmuir isotherms, to thermodynamic models, such as surface complexation models. The current limit in transitioning to thermodynamically driven models is our inability to measure proton

457 reactivity in the presence of carbonate minerals and in turn to quantify the number of surface sites
458 could adsorb metals. While metals can be used as a probe for overall mat reactivity, the reactivity
459 of bacterial mats will vary based on the metal tested (depending on, i.e., charge and hydrodynamic
460 radius), interactions between individual sorbents in the intact mat, and potential diffusional or
461 kinetics effects. Therefore, further work is required to constrain proton reactivity of the biomass
462 component of intact mats, and to provide improved methods for predicting total reactivity of mats
463 toward metals in alkaline and carbonate rich systems.

464 **Acknowledgements:**

465 This work was supported by Discovery Grants from the Natural Sciences and Engineering
466 Research Council of Canada (NSERC) to Daniel Alessi and Kurt Konhauser. Leslie Robbins
467 gratefully acknowledges the support of a NSERC Vanier CGS, and Tyler Warchola a NSERC
468 CGS-M. We also would like to acknowledge the staff of the Fairmont Hot Spring Resort for
469 providing us with access to the site, as well as working space to set up a temporary lab. In addition,
470 we thank the two reviewers for their insightful and pointed reviews which greatly aided in the
471 improvement of this manuscript.

472

473

References:

- Alessi DS, and Fein JB (2010) Cadmium adsorption to mixtures of soil components: Testing the component additivity approach, *Chem. Geol.*, **270**, 186-195.
- Baker MG, Lalonde Sv, Konhauser KO, and Foght JM (2009) Role of extracellular polymeric substances in the surface chemical reactivity of *Hymenobacter aerophilus*, a psychrotolerant bacterium, *Appl. Environ. Microbiol.*, **76**, 102-109.
- Beveridge TJ, and Murray RGE (1976) Uptake and retention of metals by cell-walls of *Bacillus subtilis*, *J. Bacteriol.*, **127**, 1502-1518.
- Beveridge TJ, and Murray RGE (1980) Sites of metal-deposition in the cell wall of *Bacillus subtilis*, *J. Bacteriol.*, **141**, 876-887.
- Braissant O, Decho AW, Dupraz C, Glunk C, Przekop KM, and Visscher PT (2007) Exopolymeric substances of sulfate-reducing bacteria: interactions with calcium at alkaline pH and implication for formation of carbonate minerals, *Geology*, **5**, 401-411.
- Borrok D, Fein JB, Tischler M, O'loughlin E, Meyer H, Liss M, and Kemner KM (2004) The effects of acidic solutions and growth conditions on the adsorptive properties of bacterial surfaces, *Chem. Geol.*, **209**, 107-119.
- Borrok D, Turner BF, and Fein JB (2005) A universal surface complexation framework for modeling proton binding onto bacterial surfaces in geologic settings, *Am. J. Sci.*, **305**, 826-853.
- Cox JS, Smith DS, and Ferris FG (1999) Characterizing heterogeneous bacterial surface functional groups using discrete affinity spectra for proton binding, *Environ. Sci. Technol.*, **33**, 4514-4521.
- Elderfield H, Bertram CJ, and Eraz J (1996) A biomineralization model for the incorporation of trace elements into foraminiferal calcium carbonate, *Earth Planet. Sci. Lett.*, **142**, 490-423.
- Elias S and Banin E (2003) Multi-species biofilms: living with friendly neighbors, *FEMS Microbiol. Reviews*, **36**, 990-1004.
- Elzinga EJ, and Reeder RJ (2002) X-ray absorption spectroscopy study of Cu^{2+} and Zn^{2+} adsorption complexes at the calcite surface: Implications for site-specific metal incorporation preferences during calcite crystal growth, *Geochim. Cosmochim. Acta*, **66**, 3943-3954.
- Fein JB, Daughney CJ, Yee N, and Davis TA (1997) A chemical equilibrium model for metal adsorption onto bacterial surfaces, *Geochim. Cosmochim. Acta*, **16**, 3319-3228.
- Ferris FG, Beveridge TJ, and Fyfe WS (1986) Iron-silica crystallite nucleation by bacteria in a geothermal sediment, *Nature*, **320**, 609-611.

506 Ferris, F.G., Frattton, C.M., Gerits, J.P., Schultze-Lam, S., and Sherwood Lollar, B. (1995)
 507 Microbial precipitation of a strontium calcite phase at a groundwater discharge zone near Rock
 508 Creek, British Columbia, Canada. *Geomicrobiol. J.*, **13**, 57-67.

509 Flynn SL, Szymanowski JES, and Fein JB (2014) Modeling bacterial metal toxicity using a
 510 surface complexation approach, *Chem. Geol.*, **374**, 110-116.

511 Fowle DA and Fein JB (1999) Competitive adsorption of metal cations onto two gram positive
 512 bacteria: testing the chemical equilibrium model, *Geochim. Cosmochim. Acta*, **63**, 3059–3067.

513 Haas JR, Dichristina TJ, and Wade R (2001) Thermodynamics of U(VI) sorption onto
 514 *Shewanella putrefaciens*, *Chem. Geol.*, **49**, 609-611.

515 Jones, B., Konhauser, K., Renaut, R.W. and Wheeler, R. (2004) Microbial silicification in Iodine
 516 Pool, Waimangu geothermal area, North Island, New Zealand: Implications for recognition and
 517 identification of ancient silicified microbes. *Geological Society of London*, **161**, 983-993.

518 Kolter R, and Greenberg PE (2006) Microbial sciences: the superficial life of microbes, *Nature*,
 519 **441**, 300-302.

520 Konhauser KO (2007) *Introduction to Geomicrobiology*, Blackwell Publishing, Oxford, UK.

521 Konhauser, K.O., Fisher, Q.J., Fyfe, W.S., Longstaffe, F.J., and Powell, M.A. (1998) Authigenic
 522 mineralization and detrital clay binding by freshwater biofilms: The Brahmani River, India.
 523 *Geomicrobiol. J.*, **15**, 209-222.

524 Konhauser KO, and Urrutia MM (1999) Bacterial clay authigenesis: a common bacterial process,
 525 *Chem. Geol.*, **161**, 399-413.

526 Kuyucak N and Volesky B (1989) Accumulation of cobalt by marine alga, *Biotechnol. Eng.*, **33**,
 527 809-814.

528 Lalonde SV, Amskold L, Warren LA and Konhauser KO (2007) Surface chemical reactivity and
 529 metal adsorptive properties of natural cyanobacterial mats from an alkaline hydrothermal spring,
 530 Yellowstone National Park, *Chem. Geol.*, **243**, 36-52.

531 Matheichal JT, Yu Q, and Woodburn G.M. (1999) Biosorption of cadmium (II) from aqueous
 532 solution by pre-treated biomass of marine alga *Durvillaea potatorum*, *Water Res.*, **33**, 335-342.

533 Mortimer, R.J.G., Coleman, M.L., and Rae, J.E. (1997) Effect of bacteria on the elemental
 534 composition of early diagenetic siderite: Implications for palaeoenvironmental interpretations.
 535 *Sedimentology*, **44**, 759-765.

536 Mullens MD, Wolf DC, Ferris FG, Beveridge TJ, Flemming CA, and Bailey GW (1989)
 537 Bacterial sorption of heavy metals, *Appl. Environ. Microbiol.*, **55**, 3143-3149.

538 Mullens MD, Wolf DC, Beveridge TJ, and Bailey GW (1992) Sorption of heavy metals by soil
 539 fungi *Aspergillus Niger* and *Mucor Rouxii*, *Soil Biol. Biochem.*, **24**, 129-135.

540 Nealson, K.H., Tebo, B.M., and Rosson, R.A. (1988) Occurrence and mechanisms of microbial
 541 oxidation of manganese. *Adv. Appl. Microbiol.*, **33**, 279-318.

542 Newman, D.K., Beveridge, T.J., and Morel, F.M.M. (1997) Precipitation of arsenic trisulfide by
543 *Desulfotomaculum auripigmentum*. Appl. Environ. Microbiol., **63**,2022-2028.

544 Ngwenya BT, Sutherland IW, and Kennedy L (2003) Comparison of the acid–base behavior and
545 metal adsorption characteristics of gram-negative bacterium with other strains, Appl. Geochem.,
546 **18**, 527–538.

547 Okumura M and Yasushi M (1986) Coprecipitation of alkali metal ions with calcium carbonate,
548 Geochim. Cosmochim. Acta, **50**, 49-58.

549 Rainey RK, and Jones B (2009) Abiotic versus biotic controls on the development of the
550 Fairmont Hot Spring carbonate deposit, British Columbia, Canada, Sedimentology, **56**, 1832-
551 1857.

552 Sadiq M (1992) Basic concepts in Marine Chemistry, In: Toxic metal chemistry in marine
553 environments, Marcel Dekker Inc., New York, pp 5-29.

554 Schofield EJ, Veeramani H, Sharp JO, Suvorova E, Latmani RB, Mehta A, Stahlman J, Webb
555 SM, Clark DL, Conradson SD, Ilton ES, and Bargar JR (2008) Structure of biogenic uraninite
556 produced by *Shewanella onedensis* strain MR-1, Environ. Sci. Technol., **42**, 7898-7904.

557 Schultze-Lam, S. and Beveridge, T.J. (1994) Nucleation of celestite and strontionite on a
558 cyanobacterial S-layer. Appl. Environ. Microbiol., **60**, 447-453.

559 Small TD, Warren LA, Roden EE, and Ferris FG (1999) Sorption by bacteria, Fe(III) oxide, and
560 Bacteria-Fe(III) oxide composites, Environ. Sci. Technol., **33**, 4465-4470.

561 Spath R, Flemming HC, and Wuerzt S (1998) Sorption properties of biofilms, Wat. Sci. Tech.,
562 **37**, 207-210.

563 Southam, G. and Beveridge, T.J. (1994) The in vitro formation of placer gold by bacteria.
564 Geochim. Cosmochim. Acta, **58**,4527-4530.

565 Teitzel GM, and Parsek MR (2003) Heavy metal resistance of biofilm and planktonic
566 *Pseudomonas aeruginosa*, Appl. Environ. Microbiol., **69**, 2313-2320.

567 Thompson, J.G. and Ferris, F.G. (1990) Cyanobacterial precipitation of gypsum, calcite, and
568 magnesite from natural alkaline lake water. Geology, **18**,995-998.

569 Warchola TJ, Flynn SL, Robbins LJ, Liu Y, Meyers R, Gauger T, Kovalchuk O, Alam MS, Wei
570 S, Lalonde SV, Kappler A, Alessi DS, and Konhauser KO (in review) Field- and lab-based
571 potentiometric titrations of microbial mats from the Fairmont Hot Springs, Canada,
572 Geomicrobiol. J.

573 Warren, L.A., Maurice, P.A., Parmar, N., and Ferris, F.G. (2001) Microbially mediated calcium
574 carbonate precipitation: Implications for interpreting calcite precipitation and for solid phase
575 capture of inorganic contaminants. Geomicrobiol. J.,**18**, 93-115.

576 Volesky B and Holan ZR (1995) Biosorption of heavy metals, Biotechnol. Prog., **11**, 235-250.

577 Xue HB, Stumm W, and Sigg L (1988) The binding of metals to algal surfaces, *Water Res.*, **22**,
578 917-926.

579

580 Tables and Figures:

581

582 Figure 1. Total Cd in solution in equilibrium with otavite as a function of pH.

583 Table 1. Partitioning of mass between bacteria and carbonate mineral in the intact bacterial mats.

	% Mineral	% Bacteria
Spongy Green Mat	39.8	60.2
Black Mat	36.5	63.5
Orange Mat	32.1	67.9
Green Mat	72.8	27.2

584

585

586

587

588 Figure 2. Results of duplicate experiments measuring Cd adsorption onto the travertine from
 589 Fairmont Hot Springs plotted as the percentage of Cd adsorbed to the travertine as a function of
 590 Travertine concentration with the fit of the linear form of the Freundlich isotherm equation inset.

591

592

593

594

Figure 3. Cd adsorption to each of the bacterial mats plotted as percent Cd adsorbed as a function of the bacterial mat concentration with the respective fits of their linear form of the Freundlich isotherm equation inset (A. green mat, B. black mat, C. spongy green mat, and D. orange mat).

Figure 4. Cd adsorption to the acid treated bacterial mats plotted as percent Cd adsorbed as a function of the bacterial mat concentration with the respective fits of their linear form of the Freundlich isotherm equation inset (A. acid treated green mat, B. acid treated orange mat).

Table 2. Freundlich isotherm fitting parameters for each sorbent with the coefficient of determination for the linear regressions for the linear fit of the experimental data, and Freundlich fitting parameters for composite and subtractive models.

	N	K	R ²
Travertine	0.45	2.41	0.91
Green Mat	1.25	1.10	0.95
Orange Mat	0.94	1.52	0.95
Spongy Green Mat	1.57	0.27	0.86
Black Mat	1.19	0.48	0.96
Acid Treated Green	1.29	2.42	0.90
Acid Treated Orange	1.08	0.87	0.89
Composite Green Mat	0.77	2.60	
Composite Orange Mat	0.75	1.51	
Subtractive Green Mat	1.64	1.63	
Subtractive Orange Mat	1.26	1.03	
Subtractive Spongy Green Mat	3.63	0.005	
Subtractive Black Mat	2.51	0.027

Figure 5. Models of the predicted adsorbed Cd concentrations (C^*) based on the Freundlich isotherm, including travertine mineral (grey line), green mat (dark green line), black mat (black line), Spongy green mat (light green line), orange mat (orange line), acid treated green mat (dashed green line), and acid treated orange mat (dashed orange line).