Visual evidence of reduced seafloor conditions and indications of a cold seep ecosystem from the Hatton–Rockall basin (NE Atlantic).

Francis C. Neat^{*1}, Alan J. Jamieson², Heather A. Stewart³, Bhavani E. Narayanaswamy⁴, Neil Collie¹, Michael Stewart¹, Thomas D. Linley²

***Corresponding author: francis.neat@msc.org**

¹Marine Scotland Science, Marine Laboratory, Aberdeen, U.K. AB11 9DB

²School of Marine Science and Technology, Newcastle University, Newcastle Upon Tyne, U.K. NE1 7RU

³British Geological Survey, Lyell Centre, Research Avenue South, Edinburgh, U.K. EH14 4AP

⁴Scottish Association for Marine Science, Scottish Marine Institute, Oban, Argyll, U.K. PA37 1QA

Abstract

High definition video from a towed camera system was used to describe the deep sea benthic habitats within an elongate depression located at the western margin of Rockall Bank in the Hatton–Rockall Basin. At depths greater than 1190 m, an extensive area (10 km long by 1.5 km wide) of what appeared to be reduced sediments, bacterial mats and flocculent matter indicated possible cold seep habitat. Plumes of sediment rich fluid were observed alongside raised elongate features that gave topographic relief to the otherwise flat seafloor. In the deepest section of the depression (1215 m) dense flocculent matter was observed suspended in the water column, in places obscuring the seabed. Away from the bacterial mats, the habitat changed rapidly to sediments dominated by tube-dwelling polychaete worms and then to deep-sea sedimentary habitats more typical for the water depth (sponges and burrowing megafauna in areas of gentle slopes, and coral gardens on steeper slopes).

Keywords: reduced sediment environment, chemosynthetic, bacterial mats

Running Head: Camera survey of putative cold seep

Introduction

Reduced environments are found throughout the world's ocean seafloor and sediments and typically arise from a lack of dissolved oxygen or excessive organic inputs to the seabed, such as particulate organic matter (marine snow) or large animal carcasses (Tunnicliffe *et al.* 2002). They also arise from geological conditions, for example, cold seeps that are typically located along continental margins and associated with sub-seabed reservoirs of hydrocarbons (often methane) and sulphide rich fluid expelled to the seafloor by gravitational and tectonic forces (Levin, 2005). Methane seeps, mud volcanoes, pock marks, carbonate slabs and brine pools are all examples of cold-seep chemosynthetic habitats (Cordes *et al.*, 2010). At cold seeps, microbial consortia anaerobically oxidise methane, reduce sulphate to sulphide and chemically acquire the necessary energy for metabolism, growth and replication (Boetius *et al.*, 2000; Orphan *et al.*, 2002; Joye *et al.*, 2004; Dubilier *et al.*, 2008; Boetius & Wenzhöfer, 2013). The biochemical reactions result in the precipitation of authigenic carbonate which can lead to the formation of solid crusts surrounding seeps and vents that in turn provide substrate for other fauna (Bayon *et al.*, 2013).

The bacteria occur either as chemosynthetic symbionts of other organisms (Dubilier *et al.*, 2008) or as free-living bacteria. The free-living bacteria often form extensive filamentous mats across the seafloor (Boetius & Wenzhöfer, 2013). Only a few higher marine taxa have evolved the specialised adaptations to tolerate high concentrations of sulphide and live symbiotically with the bacteria (Dubilier *et al.*, 2008). These include, among others; mussels of the subfamily Bathymodiolinae; bivalve clams of the family Vesicomidae, Thyasiridae, Lucinidae and Solemyidae; annelid worms of the family Siboglinidae; and sponges of the family Cladorhizidae.

The species and communities of cold-seep sediments are often dominated by endemic fauna (Olu *et al.*, 1997; Levin, 2005; Vanreusel *et al.*, 2009). Cold seep habitats are highly heterogeneous (Cordes *et al.*, 2010) and contrast starkly with adjacent deep-sea habitats (Bowles *et al.*, 2016). The biomass of cold-seep communities can be orders of magnitude greater than the surrounding deep-sea habitat (Tunnicliffe, 1992).

Although world-wide less than 100 active seeps have been described (German *et al.* 2011), they are thought to be much more widespread along continental margins than currently documented. This is likely to be the case in the deep north-east Atlantic, where as far as we know, they are few and far between. Only three confirmed sites have so far been described; at approximately 72 °N lies the Haakon Mosby mud volcano in the Barents Sea (Niemann *et al.*, 2006; Jerosch *et al.* 2007), at around 64 °N, offshore from Norway, is found the Nyegga pock-mark region (Krylova *et al.*, 2011), and south of Spain, at 36 °N, are found the mud volcanoes of the Gulf of Cadiz (Cunha *et al.* 2013; Rodrigues *et al.*, 2010). This apparent rarity of cold seeps in the northeast Atlantic is unlikely to be real and highlights a major gap in our knowledge of deep-water benthic ecosystems. Most of the sedimentary basins of the North Atlantic margin contain potential hydrocarbon source rocks and this includes the Hatton–Rockall basin, a large expanse of submerged continental crust that lies approximately 500 km west of the British Isles (Hitchen, 2004). The underlying sediments of the Hatton–Rockall Basin are characterised by an extensive polygonal fault system which has been suggested to be a currently active process resulting in fluid expulsion (Berndt *et al.*, 2012). As seabed fluid expulsion is often associated with chemosynthetic ecosystems (Sibuet & Olu-Le Roy, 1998), the Hatton–Rockall Basin is a candidate area for harbouring cold-seep ecosystems.

87

88 In 2012 a benthic sampling net deployed at the deepest point (1200 m) of the
89 western margin of Rockall Bank recovered two new species of chemosymbiotic bivalve
90 of the families Vesicomyidae and Thyasiridae (Oliver & Drewery 2014). These
91 specimens suggested a cold seep ecosystem (hereafter referred to as the *Scotia*
92 *Seep*) within the UK continental shelf claim. However, the occurrence of these
93 'indicator' species is not itself proof of a cold-seep, as they may be associated with
94 reduced sediments arising from biotic inputs. In order to assess the seabed habitats
95 in the vicinity of where these species were found, a visual survey of the area was made
96 using a towed camera to produce the first account of the site and map the seabed
97 habitats present.

98

99 **Materials and Methods**

100

101 **Study Site**

102 The Hatton–Rockall Basin is a large, sedimentary deep-water habitat (1000-
103 1500 m water depth). Bounded by the Rockall Bank to the east and the Hatton Bank
104 to the west and north, it is located around 500 km west of the United Kingdom in
105 international waters, but within the extended continental shelf claim of the United
106 Kingdom (Figure 1, inset map). The study site (provisionally named the *Scotia Seep*
107 after MRV Scotia) was located at approximately 57° 57' N and 15° 33' W at the bottom
108 of the western slope of the Rockall Bank at depths of between 1100 and 1216 m.

109

110 **Visual survey**

Eight towed camera transects were made aboard *MRV Scotia* totalling approximately 12.5 hours and 40 km of video footage (Figure 1, main map). Transects spanned a depth range of 1040-1216 m. Seabed imagery was obtained using a towed body, consisting of an aluminium frame fitted with a Kongsberg Maritime colour HD video camera and six high-intensity undersea lamps and 30 cm spaced lasers for scale (McIntyre *et al.*, 2016). The camera also had the option to take strobe-lit stills which was done to obtain images for close inspection. In addition, the frame housed sensors for pressure (depth), bearing, altitude from seabed, pitch and heave, all recording at 1 s intervals throughout. The towed body was attached to the research vessel with a 4000 m 'Netsonde' co-axial cable and towed at a speed of between 1.5 – 2.5 knots at an elevation of between 2 to 5 m above the seafloor. An ultra-short baseline beacon (USBL) provided positional information on the towed camera, calculating the position of the body relative to the vessel. Transmission from the USBL was, however, intermittent and for many video sequences the position of the camera had to be estimated from the length of wire out and the seabed depth using standard spherical geometry (assuming the body to lie directly behind the vessel). Comparison of estimated positions with known positions from the USBL indicated estimated positions were on average within 164 m (+/- 22 m) of the known position. In addition to the towed video five point observations were made from a free fall baited lander system with time-lapse cameras that took horizontal digital stills of the seabed (Linley 2016). These were used to supplement the towed camera data.

Habitat classification

The identification of species from digital stills images and HD video is frequently problematic in the absence of physical samples; however visible species were identified to the lowest possible taxonomic level. Following detailed analysis of the HD video that involved species identification when possible and an assessment of the seabed type, six broad habitat types were evident that appeared to be generally exclusive of one another. These were; bacterial mats covering flat seabed and raised areas; sponges in bioturbated sediment; coral gardens; sand-rich sediment; sediment with tube-dwelling worms; zero visibility (when the seabed was obscured by dense flocculent matter). Segments of video were classified to one of these habitat types and using time referenced positions were then mapped in ArcGIS.

Bathymetry mapping

The ship-board Olex seabed mapping system (www.olex.no) generated a bathymetric map of the area based on echo-sounder readings. This was used to generate 10 m depth contours from 1100 to 1210 m water depth in ArcGIS on which to results are presented.

Results

Location and topography

The putative cold seep habitat was revealed within a distinct narrow depression (Figure 1) beneath a steep escarpment to the east (Rockall Bank) and a more gradual ascent to the west (the Hatton–Rockall Basin). At its deepest point the depression reached 1216 m. There were two distinct basins along the north-south axis of the basin

where water depth exceeded 1200 m, separated by a ridge that rose to around 1180 m water depth. To the south and north of the trench, the seabed gradually shallowed before levelling out at around 1100 m water depth.

Habitats and distribution

1. Bacterial mats. Large expanses of the seafloor appeared to be covered in dense bacterial mats. The bacterial mats appeared as patches on the seabed (Figure 2a and Supplementary video) between areas of fine sediment (Figure 2b) and areas where flocculent matter had settled to form a fluffy lighter brown layer (Figure 2a, b). The bacterial mats were mainly devoid of any emergent epifauna with the exception of sporadic, small unidentified anemones (supplementary video). The bacterial mats were encountered mainly at depths of 1190 m and deeper (Table 1). At times the bacterial mats covered distinct three-dimensional features referred to as 'green mounds' in Figure 2c and Table 1. These mounds rose an estimated 1 m above the surrounding seafloor, forming elongate ridges and mounds. At least four separate sediment rich fluid plumes emanating from the seafloor were observed (Figure 2d, e and supplementary video). Small concentric features of the seabed were sometimes evident (Figure 2f) that resembled holes or burrows. Visibility was variable along transects a result of often dense flocculent matter in the water column. Fish (e.g. *Corephaeneoides rupestris*, *Chimaera opalescence*, *Synaphobranchus kaupi*) were seen swimming above the bacterial mats.

2. Sponges in bioturbated sediment. This habitat was characterised by soft, bioturbated sediment (Figure 3a). Megafauna included sponges, of which the

hexactinellid *Pheronema carpenteri*, the encrusting yellow sponge *Hexadella* sp. and a stalked sponge of the genus *Hyalonema* were identified. What appeared to be Xenophyophores were evident, although at times it was difficult to distinguish these from species of sponge. The squat lobster *Munida* sp. was seen frequently, sometimes partially burrowed in the sediment. The biotope was further characterised by cerianthid anemones, of which *Pachycerianthus* sp. was identified. Other commonly observed taxa included holothurians of the genus *Benthogone* and various large echinoids including *Calveriosoma* sp. This habitat was characteristic of the shallower western slopes (1130-1180 m) of the study area (Table 1). Fish were observed including *Lepidon eques*. This habitat rapidly disappeared at depths of around 1180 m and greater, transitioning into either the bacterial mat or tubeworm dominated habitats.

3. Coral gardens. This habitat (Figure 3b) was typified by the presence of gorgonian corals (sea fans) of the genus *Placogorgia*, stony coral (*Madrepora oculata*), alcyonaceans (soft corals) and the anemone *Bolocera* sp. The slope was steep and the seabed composed of outcrops of rock and areas of gravel with intermittent patches of fine-grained sediments. Fish were observed including *Lepidon eques*. This habitat was only observed in the eastern reaches of the study area, coincident with the steep flank of Rockall Bank along a narrow depth range of between 1140 and 1167 m (Table 1).

4. Sand-rich sediment. In this habitat the sediment comprised muddy sand with occasional gravel sized clasts. It was predominantly not bioturbated and lacked sponges. Occasional boulders were colonised by an unidentified species of barnacle.

Other benthic fauna included small cerianthid anemones and holothurians including *Benthogone* sp. This habitat was observed on the ridge between the two basins, but also extended into the basins in places and was characteristic of the southern limit of the basin as the depth decreased.

5. Tube-dwelling polychaete worm sediment. This habitat was characterised by the presence of high densities of small unidentified tube-dwelling polychaete worms (Figure 3 c, d). At times there appeared also to be a light covering of bacterial mat. There was no evidence of bioturbation and no sponges or other benthic megafauna. This habitat was typically seen adjacent to and at marginally shallower water depths than the bacterial mats habitat (Table 1).

6. Zero-visibility. This category was reserved for sections of transects where the seabed was completely obscured by dense flocculent matter in the water column (see supplementary video). The location of this cloud of flocculent matter was predominantly encountered in the deepest sections of the study area (Table 1), although in two transects areas of zero-visibility was encountered at shallower water depths and in one instance it was encountered where the seabed had been visible on another transect.

Discussion

Extensive areas of what appeared to be bacterial mats smothering the seafloor were observed in this study. The localised nature of the mats and their absence from

adjacent shallower areas suggest a focal source of reduced sediments, hydrogen sulphide production and/or methane. The bacterial mats could be indicative of a chemosynthetically active cold seep ecosystem as has been reported from numerous sites across the world's oceans (Boetius & Wenzhöfer, 2013). Alternatively they could be sustained through excessive accumulation of photosynthetically derived particulate organic matter (Gooday & Lamshead, 1989). This would fit with the bacterial mats being found predominantly in the deepest part of the study area (below 1190 m) and, also with the survey being undertaken in mid-summer following the spring phytoplankton bloom in the NE Atlantic. The presence of bacterial mats together with the initial discoveries of chemosynthetic clams (Oliver & Drewery, 2014) provides strong evidence for the presence of a reduced sediment conditions and possibly the first cold-seep ecosystem to have been found within the continental shelf limits of the UK. In places the seafloor was obscured due to dense flocculent matter in the water column, but again it is not clear at this stage if this is bacterially derived flocculants from the seabed or is pelagic derived phytodetritus. Bacterial mats and dense particulate matter in the water column have been previously reported from cold-seeps from the Barents Sea (Grunke *et al.* 2012), the Gulf of Mexico (Sassen *et al.*, 1993), the Californian slope (Levin, 2003) and the New Zealand margin (Baco *et al.*, 2009). The near-bottom water throughout the area was clearly not toxic to fish as numerous species were observed swimming directly above the seabed.

Assuming that most of the seabed below 1190 m water depth is covered in bacterial mats this would equate to an area of at least 10 km² similar in scale to cold seeps found on the slope of the Gulf of Mexico (Sibuet & Olu 1998). While there was no evidence for gas bubbling or 'shimmering water' from the seabed, as is often

reported from cold-seeps and mud volcanoes (Neimann *et al.*, 2006; Skarke *et al.*, 2014), sediment rich fluid plumes were observed that might suggest an active geological process. Water currents, fish motion and pressure waves from the towed camera could have caused sediment disturbance, but the plumes were sporadic and very localised, occurring both in absence and presence of fish (see supplementary video). The fluid plumes were seen in association with unusual elongate three-dimensional features on the seabed. Directed sampling of the sediments, mounds and plumes is needed to confirm their origin and nature.

The video footage did not reveal any obvious aggregations of bivalves in the bacterial mats as has been reported from other cold seep sites (Barry *et al.* 1996; Sibuet & Olu, 1998; Baco *et al.*, 2009), but this may simply reflect the small size and burrowing habits of the Vesicomidae and Thyasiridae clams described by Oliver & Drewery (2014). Overall there was a scarcity of benthic epifauna associated with the bacterial mats, with the exception of small, dark red anemones that appeared sporadically. These remain to be identified and it is not known if they belong to the several genera of cerianthid anemones that are known to associate with reduced deep-sea habitats (Rodriguez & Daly 2012). Dense fields of what appeared to be small (2-3 cm in height) tube-dwelling polychaete worms were observed in areas adjacent to the bacterial mats. They clearly were not large enough and did not have the characteristic bright red tentacles to suggest they were species of the family Siboglinidae (that are well known from cold seeps) and is not clear whether they are chemosynthetic or not. It is interesting however that they appeared in a transition zone between the bacterial mats and the sponges and bioturbated sediment habitat observed at shallower depths.

278 In the area beyond the reduced sediments, on the shallower flanks of the
279 depression, numerous sessile and more typical deep-sea species were observed.
280 These included hexactinellid sponges *Hyalonema* sp. (stalked sponge) and
281 *Pheronema carpenteri*, the ophiuroid *Ophiecten gracilis* and possibly
282 Xenophyophores. This habitat occurred primarily on the shallower western slope of
283 the study area and was characterised by bioturbation of the sediment and the
284 presence of the squat lobster *Munida* sp. and the holothurian, *Benthogone* sp. These
285 species and similar biotopes have been reported from numerous sites nearby in the
286 Rockall Trough and Hatton–Rockall Basin (Hughes & Gage, 2004; Narayanaswamy
287 *et al.*, 2013; McIntyre *et al.*, 2016). The coral garden habitat occurred on the steep
288 eastern slope of the study area and again contrasted sharply with the seabed in the
289 depression. Various gorgonians, the anemone *Phelliactis* sp. and soft corals
290 (alcyonaceans) were seen in this area. Coral gardens are frequently associated with
291 escarpment features with steep gradients, boulders and exposed bedrock (Davies *et*
292 *al.*, 2015).

293
294 The habitats observed at the *Scotia Seep* contrasted starkly from the
295 surrounding deep seafloor and slopes. A concentric pattern of habitats was apparent
296 with a central area located in the deepest part of the depression covered with dense
297 bacterial mats, elongate raised features and sediment rich fluid plumes. Away from
298 the bacterial mats there appears to be transition into sediments with tube-dwelling
299 worms, and in turn to more typical deep-sea bioturbated sediments and communities.
300 Concentric patterns of chemosynthetic habitat radiating out from active seep areas
301 have been previously described (Olu *et al.*, 1997; Levin *et al.*, 2003, 2005, 2016).

302

303 Chemosynthetic ecosystems are currently understood to be rare in the NE
304 Atlantic with only the three sites previously mentioned confirmed to date. Given that
305 these are thousands of kilometres away from the current study site, it is perhaps not
306 surprising that species from the *Scotia Seep* have turned out to be new to science
307 (Oliver & Drewery 2014). Interestingly, west of the *Scotia Seep* there is a network of
308 kilometre-scale polygonal faults that extend across the Hatton–Rockall Basin (Berndt
309 *et al.*, 2012). It is possible the *Scotia Seep* is connected in some way to this geological
310 fault system. The observation of flocculent matter above the seabed suggests it could
311 have a significant effect on the overlying water column. Methane plumes from cold
312 seeps in the Gulf of Mexico, at depths similar to this study are thought to support
313 planktonic microbial communities from the seabed into the surface water (Rakowski
314 *et al.*, 2015).

315

316 In conclusion while the habitats described are clearly atypical of the seabed
317 sediments in the north Atlantic for these depths, it is not clear whether it is a geological
318 driven cold seep ecosystem or a biologically driven reduced environment. It might
319 even arise from both if geological conditions (emissions of hydrogen sulphide rich
320 fluids) give rise to toxic seafloor conditions for detritivores, the absence of which allows
321 phytodetritus to build-up. Geochemical analyses of sediment samples and genetic
322 bar-coding of the microbial and invertebrate communities are currently underway and
323 should better define the biogeochemical nature of the *Scotia Seep*.

324

325 **Acknowledgments**

326

327 The MRV Scotia 0915S expedition was awarded to the Marine Alliance for Science
328 and Technology Scotland (MASTS) Deep-Sea Forum through an open call for ship
329 time by Marine Scotland Science (Scottish Government). The MASTS Deep-Sea
330 Forum received support from Scottish Funding Council (grant reference HR09011).
331 We are grateful for help from the other scientists aboard survey 0915S and the officers
332 and crew of MRV Scotia for their professional assistance at sea. H.A.S. publishes with
333 permission of the Executive Director of the British Geological Survey (Natural
334 Environment Research Council).

335

336 **References**

337 **Baco, A., Rowden, A., Levin, L.A., Smith, C.R. and Bowden, D.A.** (2009) Initial
338 characterization of cold seep faunal communities on the New Zealand Hikurangi
339 margin. *Marine Geology* 272, 251-259

340

341 **Barry, J.P., Greene, H.G., Orange, D.L., Baxter, C.H., Robison, B.H., Kochevar,**
342 **R.E., Nybakken, J.W. and McHugh, C.M.** (1996) Biologic and geologic
343 characteristics of cold seeps in Monterey Bay, California. *Deep Sea Research Part I*,
344 43, 1739-1762.

345

346 **Bayon, G., Dupre, S., Ponzevera, E., Etoubleau, J., Cheron, S., Mascle, J.,**
347 **Boetius, A. and de Lange, G.T.** (2013). Formation of carbonate chimneys in the

348 Mediterranean Sea linked to deep-water oxygen depletion. *Nature Geosciences* 6,
349 755-760.

350

351 **Berndt, C., Jacobs, C., Evans, A., Gay, A., Elliott, G., Long, D. and Hitchen, K.**
352 (2012). Kilometre-scale polygonal seabed depressions in the Hatton Basin, NE
353 Atlantic Ocean: Constraints on the origin of polygonal faulting. *Marine Geology*, 332,
354 126-133.

355

356 **Boetius, A., Ravensschlag, K., Schubert, C.J., Rickert, D., Widdel, F., Gieseke,**
357 **A., Amann, R., Barker Jørgensen, B., Witte, U. and Pfannkuche, O.** (2000) A
358 marine microbial consortium apparently mediating anaerobic oxidation of methane.
359 *Nature* 427, 623-626.

360

361 **Boetius, A. and Wenzhöfer, F.** (2013). Seafloor oxygen consumption fuelled by
362 methane from cold seeps. *Nature Geosciences* 6, 725–734.

363

364 **Bowles, M., Hunter, K.S., Samarkin, V. and Joye, S.** (2016). Patterns and
365 variability in geochemical signatures and microbial activity within and between
366 diverse cold seep habitats along the lower continental slope, Northern Gulf of
367 Mexico. *Deep Sea Research Part II*, 129, 31-40.

368

Cordes, E.E., Cunha, M.R., Galeron, J., Mora, C., Olu-Le Roy, K., Sibuet, M., Van Gaever, S., Vanreusal, A. and Levin, L. (2010) The influence of geological, geochemical and biogenic habitat heterogeneity on seep biodiversity. *Marine Ecology*, 31, 51-65.

Cunha, M. R., Rodrigues, C. F., Genio, L., Hilario, A., Ravara, A., and Pfannkuche, O. (2013). Macrofaunal assemblages from mud volcanoes in the Gulf of Cadiz: abundance, biodiversity and diversity partitioning across spatial scales, *Biogeosciences*, 10, 2553–2568.

Davies, J.S., Stewart, H.A., Narayanaswamy, B.E., Jacobs, C., Spicer, J., Golding, N., and Howell, K.L. (2015). Benthic Assemblages of the Anton Dohrn Seamount (NE Atlantic): Defining deep-sea biotopes to support habitat mapping and management efforts with a focus on vulnerable marine ecosystems. *PLoS ONE*, 10: e0124815.

Dubilier, N., Bergin, C. and Lott, C. (2008). Symbiotic diversity in marine animals: the art of harnessing chemosynthesis. *Nature Reviews in Microbiology*, 6, 725-740.

German, C.R., Ramirez-Llodra, E., Baker, M.C. and Tyler, P.A. (2011) Deep-water chemosynthetic ecosystem research during the Census of Marine Life decade and beyond: A proposed deep-ocean road map. *PLOS One* 6, 1-16.

392 **Gooday, A.J. and Lambshead, P.J.D.** (1989). Influence of seasonally deposited
393 phytodetritus on benthic foraminiferal populations in the bathyal northeast Atlantic:
394 the species response. *Marine Ecology Progress Series*. 58, 53-67.

395

396 **Grunke, S., Lichtschlag, A., de Beer, D., Felden, J., Salman, V., Ramette, A.,**
397 **Schulz-Vogt, H.N. and Boetius, A.** (2012). Mats of psychrophilic thiotrophic
398 bacteria associated with cold seeps of the Barents Sea. *Biogeosciences*, 9, 2947–
399 2960.

400

401 **Hitchen, K.** (2004). The geology of the UK Hatton–Rockall margin. *Marine*
402 *Petroleum Geology*, 21, 993-1012.

403

404 **Hughes, D.J. and Gage, J.D.** (2004). Benthic metazoan biomass, community
405 structure and bioturbation at three contrasting deep-water sites on the northwest
406 European continental margin. *Progress in Oceanography*, 63, 29-55.

407

408 **Jerosch K., Schluter M., Foucher J.P., Allais A.G., Klages M., Edy C.** (2007).
409 Spatial distribution of mud flows, chemoautotrophic communities, and
410 biogeochemical habitats at Hakon Mosby Mud Volcano. *Marine Geology*, 243, 1–17.

411

Joye, S.B., Boetius, A., Orcutt, B.N., Montoya, J.P., Schulz, H.N., Erickson, M.J. and Lugo, S.K. (2004). The anaerobic oxidation of methane and sulfate reduction in sediments from Gulf of Mexico cold seeps. *Chemical Geology*, 205, 219-238.

Juniper, S.K. and Brinkhurst, R.O. (1986). Water-column dark CO₂ fixation and bacterial-mat growth in intermittently anoxic Saanich Inlet, British Columbia. *Marine Ecology Progress Series*, 33, 41–50.

Krylova, E.M., Gebruk, A.V., Portnova, D.A., Todt, C., Haflidason, H. (2011) New species of the genus *Isorropodon* (Bivalvia: Vesicomidae: Pliocardiinae) from cold methane seeps at Nyegga (Norwegian Sea, Vøring Plateau, Storrega Slide). *Journal of the Marine Biological Association of the United Kingdom*, 91, 1135-1144.

Levin, L.A. (2003). Spatial heterogeneity of macrofauna at northern California methane seeps: influence of sulphide concentration and fluid flow. *Marine Ecology Progress Series*, 265, 123-139.

Levin, L.A. (2005). Ecology of cold-seep sediments: Interactions of fauna with flow, chemistry and microbes. *Oceanography and Marine Biology*, 43, 1-46.

Levin L.A., Baco, A.R., Bowden, D.A., Colaco, A, Cordes, EE, Cunha, M.R., Demopoulos, A.W.J., Gobin, J., Grupe, B.M., Le, J., Metaxas, A., Netburn, A.N.,

**Rouse,G.W., Thurber, A.R., Tunnicliffe, V., Van Dover, C.L., Vanreusel, A.,
Watling L.** (2016). Hydrothermal Vents and Methane Seeps: Rethinking the Sphere
of Influence. *Frontiers in Marine Science*, 3.DOI:10.3389/fmars.2016.00072

Linley,T.D. (2016). *Fishes of the pacific abyssal and hadal zones: new technology
and advances in geographic and bathymetric resolution*. PHD thesis. University of
Aberdeen.

McIntyre, F.D., Drewery, J., Eerks-Medrano, D. and Neat, F. (2016) Diversity and
distribution of deep-sea sponge grounds on the Rosemary Bank Seamount, NE
Atlantic. *Marine Biology*, 163, 143-153.

Narayanaswamy, B.E., Hughes, D.J., Howell, K.L., Davies, J. and Jacobs, C.
(2013). First observations of megafaunal communities inhabiting George Bligh Bank,
northeast Atlantic. *Deep Sea Research Part II* 92, 79-86.

**Niemann,H., Lösekann, T., de Beer, D., Elvert, M., Nadalig, T., Knittel, K.,
Sauter, E.J., Schluter, M., Klages, M., Foucher, J.P. and Boetius, A.** (2006) Novel
microbial communities of the Haakon Mosby mud volcano and their role as a
methane sink. *Nature* 443, 854-8.

455 **Oliver, P.G. and Drewery, J.** (2014). New species of chemosymbiotic clams
456 (Bivalvia: Vesicomidae and Thyasiridae) from a putative 'seep' in the Hatton–
457 Rockall Basin, north-east Atlantic. *Journal of the Marine Biological Association of the*
458 *United Kingdom* 94, 389-403.

459

460 **Olu, K., Lance, S., Sibuet, M., Henry, P., Fiala-Médioni, A. and Diné, A.** (1997).
461 Cold seep communities as indicators of fluid expulsion patterns through mud
462 volcanoes seaward of the Barbados accretionary prism. *Deep Sea Research Part I*,
463 44, 811-841.

464

465 **Orphan, V.J., House, C.H., Hinrichs, K., McKeegan, K.D. and DeLong, E.F.**
466 (2002). Multiple archaeal groups mediate methane oxidation in anoxic cold seep
467 sediments. *Proceeding of the National Academy of Sciences USA*, 99, 7663–7668.

468

469 **Rakowski, C., Magen, C., Bosman, S., Gillies, L., Rogers, K. and Chanton, J.**
470 (2015). Methane and microbial dynamics in the Gulf of Mexico water column.
471 *Frontiers in Marine Science*, 2, 69.doi:10.3389/fmars.2015.00069

472

473 **Rodríguez, E. and Daly, M.** (2010) Phylogenetic relationships among deep-sea and
474 chemosynthetic sea anemones: Actinoscyphiidae and Actinostolidae (Actiniaria:
475 Mesomyaria). *PLoS ONE* 5(6), e10958.

476

477 **Rodrigues, C.F., Webster, G., Cunha, M.R., Duperron, S. and Weightman, A.J.**
478 (2010). Chemosynthetic bacteria found in bivalve species from mud volcanoes of the
479 Gulf of Cadiz. *FEMS Microbiology and Ecology*, 73, 486-499.

480

481 **Sassen, R., Roberts, H.H., Aharon, P., Larkin, J., Chinn, E. and Carney, R.**
482 (1993) Chemosynthetic bacterial mats at cold hydrocarbon seeps, Gulf of Mexico
483 continental slope. *Organic Geochemistry*, 20, 77-89.

484

485 **Sibuet, M. and Olu, K.** (1998) Biogeography, biodiversity and fluid dependence of
486 deep-sea cold-seep communities at active and passive margins. *Deep Sea*
487 *Research Part II*, 45, 517-567.

488

489 **Skarke, A., Ruppel, C., Kodis, M., Brothers, D., Lobecker, E.** (2014) Widespread
490 methane leakage from the sea floor on the northern US Atlantic margin. *Nature*
491 *Geoscience*, 7, 657–661.

492

493 **Tunnicliffe, V.** (1992). Hydrothermal-vent communities of the deep sea. *American*
494 *Scientist*, 80, 336-349.

495

496 **Tunnicliffe, V., Juniper, S. K. and Sibuet, M.** (2003). Reducing environments of the
497 deepsea floor. In (Tyler, P. A. ed.) *Ecosystems of the World: The Deep Sea*. Chapter
498 4, pp. 81 - 110. London, Elsevier Press.

499

500 **Van Dover, C.L.** (2000). *The ecology of deep-sea hydrothermal vents*. Princeton
501 University Press, 448 pp.

502

503 **Vanreusel, A., Andersen, A.C., Boetius, A., Connelly, D., Cunha, M.R., Decker,**
504 **C., Hilario, A., Kormas, K.A., Maignien, L., Olu, K., Pachiadaki, M.** (2009)
505 Biodiversity of cold seep ecosystems along the European margins. *Oceanography*
506 22, 110-127.

507

508 **Figure Captions**

509

510 **Figure 1.** Map of the study site and the habitats found along the towed camera
511 transects. Baited camera deployments are also indicated (circles) as are the sites of
512 active vents. Bathymetric contours are plotted every 10 m from 1100 m to 1210m.
513 The 1190 m isobaths is shown in black to indicate the area in which the majority of
514 cold seep habitats were observed. Inset map gives location of the putative cold seep
515 (red dot).

516

517 **Figure 2.** Seabed imagery of cold seep habitats. (a) and (b) The bacterial mats
518 shrouding the seabed. The paler areas are flocculent matter that has settled from the
519 water column onto the mats. (c) An example of the raised elongate mounds on the
520 seabed covered with bacterial mats. (d) and (e) Active fluid seepage in association

521 with the raised mounds. (f) An example of what appears to be a hole in the seabed
522 between the bacterial mats. Red laser points are 30 cm apart

523

524 **Figure 3.** Seabed imagery of habitats on slopes adjacent to the cold seep. (a)
525 Bioturbated sediments with sponges/and or xenophyophores. (b) Coral gardens with
526 gorgonian corals, stony corals and anenomes. (c) Tube-dwelling worm dominated
527 sediments. (d) Close up of the tube-dwelling worm dominated sediments (image
528 acquired from a baited lander deployment).

Table 1. Depth ranges (derived from pressure sensor on towed camera) for each of the habitat types observed in the study.

habitat	average depth (m)	max depth (m)	min depth (m)
bacterial mats	1193.3	1211.2	1070.8
green mounds	1201.4	1208.1	1188.2
tube worms	1174.1	1200.0	1149.7
sand	1191.2	1210.8	1117.5
sponges	1132.5	1197.4	1068.9
coral garden	1157.0	1167.7	1140.4
zero visibility	1201.8	1216.6	1140.1

Figure 1.