Nonomuraea insulae sp. nov., isolated from forest soil

Salih Saricaoglu^{1, 2}, Imen Nouioui³, Hilal Ay⁴, Hayrettin Saygin², Kadriye Inan Bektas⁵,
Kiymet Guven⁶, Demet Cetin⁷, Hans-Peter Klenk³, Kamil Isik², Nevzat Sahin^{2†}

¹Department of Nursing, School of Health, Ahi Evran University, 40100, Kırşehir, Turkey

²Department of Biology, Faculty of Science and Arts, Ondokuz Mayıs University, 55139, Samsun, Turkey

³School of Natural and Environmental Sciences, Newcastle University, Ridley Building 2, Newcastle upon Tyne NE1 7RU, UK

⁴Department of Molecular Biology and Genetics, Faculty of Sciences and Arts, Ondokuz Mayıs University, 55139, Samsun, Turkey

⁵Department of Molecular Biology and Genetics, Faculty of Science, Karadeniz Technical University, 61080, Trabzon, Turkey

⁶Department of Biology, Faculty of Science, Anadolu University, 26470, Eskisehir, Turkey

⁷Division of Science Education, Department of Mathematics and Science Education, Gazi University, 06500, Ankara, Turkey

[†]**Author for correspondence:** Nevzat Sahin

Tel : +90 362 312 19 19 - Ext: 5452

Fax: +90 362 457 60 81

E-mail: nsahin@omu.edu.tr

Abstract

Strain H2R21^T, a novel actinobacterium, isolated from a forest soil sample collected from Heybeliada, Istanbul, Turkey, and a polyphasic approach was used for characterisation of the strain. Chemotaxonomic and morphological characterisation of strain H2R21^T indicated that it belongs to the genus *Nonomuraea*. 16S rRNA gene sequence similarity showed that the strain is closely related to *Nonomuraea purpurea* 1SM4-01^T (99.1 %) and *Nonomuraea solani* CGMCC 4.7037^T (98.4 %). DNA-DNA relatedness values were found to be lower than 70 % between the isolate and its phylogenetic neighbours *N. purpurea* 1SM4-01^T, *N. solani* CGMCC 4.7037^T and *Nonomuraea rhizophila* YIM 67092^T. The whole cell hydrolysates of strain H2R21^T were found to contain *meso*-diaminopimelic acid as the diagnostic diamino acid and glucose, madurose, mannose and ribose as the cell sugars. The polar lipids were identified as phosphatidylglycerol, diphosphatidylglycerol, phosphatidylmethylethanolamine, phosphatidylethanolamine, hydroxy-phosphatidylethanolamine, dihydroxy-phosphatidylethanolamine, phosphatidylinositol, glycoposphatidylinositol, two glycopospholipids and two unidentified lipids. The predominant menaquinones were identified as MK-9(H₄) and MK-9(H₆). The major fatty acids were found to be *iso*-C_{16:0}, *iso*-C_{16:0} 2OH and C_{17:0} 10-*methyl*. On the basis of DNA-DNA relatedness data and some phenotypic characteristics, it is evident that strain H2R21^T can be distinguished from the closely related species in the genus *Nonomuraea*. Thus, it is concluded that strain H2R21^T represents a novel species of the genus *Nonomuraea*, for which the name *Nonomuraea insulae* sp. nov. is proposed. The type strain is H2R21^T (=DSM 102915^T =CGMCC 4.7338^T =KCTC 39769^T).

Key words: *Actinobacteria*, *Nonomuraea insulae* sp. nov., Polyphasic taxonomy, 16S rRNA gene

Introduction

The genus *Nonomuraea*, containing aerobic, Gram-stain positive and non-acid-fast bacteria was proposed by Zhang et al. (1998) and belongs to the family *Streptosporangiaceae*. Members of the genus *Nonomuraea* form extensively branched substrate and aerial mycelia, which differentiate into hooked, spiral, or straight chains of spores (Kämpfer et al. 2005; Fang et al. 2017). Currently, the genus consists nearly fifty and 2 subspecies with validly published names (<http://www.bacterio.net/nonomuraea.html>). The members of the genus are characterised by the presence of *meso*-diaminopimelic as the diamino acid and madurose as the diagnostic sugar in whole cell sugar patterns (chemotype IIIB sensu Lechevalier and Lechevalier 1970). Characteristically, members of the genus are rich in iso-C_{16:0} and C_{17:0} 10-methyl branched chain fatty acids; contain MK-9(H₄), MK-9(H₂) and MK-9(H₀) as predominant menaquinones; and have diphosphatidylglycerol, phosphatidylethanolamine and hydroxylated phosphatidylethanolamine as predominant phospholipids (Kämpfer 2012). Members of the genus *Nonomuraea* have been recently isolated from various habitats such as soil (Wang et al. 2014; Sriprechasak et al. 2017), rhizosphere (Shen et al. 2016; Wang et al. 2017), cave (Fang et al. 2017), root (Rachniyom et al. 2015; Li et al. 2017), coastal sediment (Xi et al. 2011), plant tissues (Li et al. 2011; Niemhom et al. 2017) and mangrove sediment (Suksard et al. 2016). Additionally, it is known that some secondary metabolites such as maduramycin, pyralomicin, the glycopeptide antibiotic A40926 and brartemicin are produced by members of the genus *Nonomuraea* (Sungthong and Nakaew 2015; Dalmastri et al. 2016).

In this study, we aimed to isolate novel actinobacteria from soil samples collected from Heybeliada island. The results of a polyphasic characterisation of strain H2R21^T, one of the isolated strains, indicated that it represents a novel species in the genus *Nonomuraea*, here named *Nonomuraea insulae* sp. nov.

Materials and methods

Isolation and maintenance of strains

A forest soil sample was collected from Heybeliada (GPS coordinates 40°52'22.30"N and 29°04'33.80"E), İstanbul, Turkey. The sample was air-dried at room temperature for 14 days, heated at 60 °C for 20 min and suspended with Ringer's solutions (Oxoid) to prepare serial dilutions. The suspension was spread on humic acid-vitamin (HV) agar (Hayakawa and Nonomuraea 1987) supplemented with filter sterilised cycloheximide (50 µg/ml), nalidixic acid (10 µg/ml) and rifampicin (0.5 µg/ml). After incubation at 28 °C for two weeks, strain H2R21^T was purified on N-Z Amine agar (DSMZ Medium 554) for further study and stored in glycerol (20 %, v/v) at -80 °C for long term preservation. The reference strains *Nonomuraea solani* DSM 45729^T and *Nonomuraea rhizophila* DSM 45382^T were purchased from DSMZ (German Collection of Microorganisms and Cell Cultures) and *Nonomuraea purpurea* 1SM4-01^T was obtained from Dr. Kannika Duangmal (Kasetsart University).

Morphological, cultural and physiological characteristics

The micromorphological properties of the strain were observed by examining gold coated dehydrated specimens grown on ISP 3 agar medium for 4 weeks using a JEOL JSM 6060 instrument. Cultural characteristics including growth, colours of the substrate mycelia and aerial mycelia, and soluble pigmentation were determined on ISP media 2-7 (Shirling and Gottlieb 1966), modified Bennett's agar (MBA; Jones 1949), Czapek's agar (Waksman 1967), nutrient agar (Waksman 1961) and tryptic soy agar (TSA; Difco) after 14 days of incubation at 28 °C using ISCC-NBS colour charts (Kelly 1964). Growth was determined at various temperatures (4, 10, 20, 28, 37, 40, 45, 50 and 55 °C), different pH (4.0-12.0) and tolerance of NaCl (up to 10 %, w/v, at intervals of 1.0 % unit) on yeast extract-malt extract (ISP 2) agar. The incubation time for temperature tests was 14 days, while the time for pH tests and tolerance of NaCl was 28 days. Degradations of Tweens 20, 40 and 80 were evaluated using the methods described by Nash and Krent (1991) while the other degradation tests (adenine,

hypoxanthine, guanine, starch, xylan and xanthine) were carried out as described by Williams et al. (1983). Utilisation of carbohydrates as sole carbon sources at a final concentration of 1 % (w/v) was performed according to the methods proposed by Shirling and Gottlieb (1966). Nitrogen source utilisation was examined according to Williams et al. (1983) with a final concentration of 0.1 % (w/v) for each nitrogen source.

Chemotaxonomic characterisation

Biomass for chemotaxonomic studies of the isolate was prepared by growing cultures in N-Z Amine broth for 2 weeks at 28 °C with shaking at 150 rpm, harvested and washed three times with distilled water before freeze-drying. The diaminopimelic acid isomers and whole cell sugars of strain H2R21^T were analysed according to Stanneck and Roberts (1974) and Lechevalier and Lechevalier (1970) by thin layer chromatography (TLC), respectively. Polar lipids were extracted and analysed using the method of Minnikin et al. (1984) by two-dimensional TLC. Isoprenoid quinones were extracted and purified from freeze-dried cells by following the procedure of Collins et al. (1985) and analysed by high performance liquid chromatography (HPLC) (Kroppenstedt 1982). The cellular fatty acids were extracted, methylated and analysed by gas chromatography using an Agilent Technologies 6890 N instrument, fitted with an autosampler and a 6783 injector, following the instructions for the Sherlock Microbial identification (MIDI) system (Sasser 1990; Kämpfer and Kroppenstedt 1996); the fatty acid methyl ester peaks were quantified using the TSBA 5.0 database.

DNA preparation, amplification and determination of 16S rRNA gene sequence

Genomic DNA extraction, PCR-mediated amplification and 16S rRNA gene sequencing were carried out as described by Chun and Goodfellow (1995) using an ABI PRISM 3730 XL automatic sequencer. An almost complete 16S rRNA gene sequence (1463 bp) was deposited in the GenBank database and aligned with corresponding sequences of representative type strains of the genus *Nonomuraea* retrieved from the EzBioCloud server (Yoon et al. 2017).

Phylogenetic trees were constructed with the neighbour-joining (Saitou and Nei 1987), maximum-likelihood (Felsenstein 1981) and maximum-parsimony (Kluge and Farris 1969) algorithms in MEGA version 7.0 software (Kumar et al. 2016). Evolutionary distances were calculated using model of Jukes and Cantor (1969) and topologies of the resultant trees evaluated by bootstrap analysis (Felsenstein 1985) based on 1000 resamplings. *Thermopolyspora flexuosa* DSM 43186^T (AY039253) was used as outgroup.

DNA base composition and DNA-DNA hybridization

In order to determine G+C content, genomic DNA of strain H2R21^T was extracted and purified by thermal denaturation (*T_m*) method as described by Mandel and Marmur (1968) using *Escherichia coli* JM109 as the reference strain. DNA–DNA hybridization was performed spectrophotometrically by DNA reassociation kinetics as described by De Ley et al. (1970) under consideration of the modifications described by Huss et al. (1983). Genomic DNAs from *N. purpurea* 1SM4-01^T, *N. solani* DSM 45729^T, *N. rhizophila* DSM 45382^T and strain H2R21^T were prepared according to the method of Marmur (1961); the purity and quality of each DNA preparation were checked by determination of the absorbance ratios at 260/280 nm (1.8 to 1.9) and at 260/230 nm (2.0 to 2.3) (according to Marmur, 1961) and by the melting curves, which include hyperchromicity values from 30 to 40% (Mandel & Marmur, 1968). Purified genomic DNAs were sonicated using a Sartorius Labsonic M to generate DNA fragments of between 400 and 800 bp in size. Sheared genomic DNA (about 100 mg) from each sample was added to 2 X SSC buffer and then denatured by raising the temperature to 100 °C and cooled to the melting temperature in the spectrophotometer. The samples were kept at the optimal renaturation temperature for 3 min and the absorbance at 260 nm was recorded at 30 s intervals for a total of 30 min. The initial reassociation kinetics were determined by using linear regression analysis. The DNA hybridization values (%) for these microorganisms were calculated using the equation described by De Ley et al. (1970). Hybridizations were replicated

five times for each sample. The highest and lowest values for each sample were excluded and the remaining three values were used for calculation of the mean level of DNA–DNA relatedness.

Results and discussion

Strain H2R21^T was observed to show good growth on ISP 2, ISP 3, ISP 6, TSA, MBA and nutrient agar media, moderate growth on ISP 7, weak growth on ISP 5 and Czapek's agar media but no growth on ISP 4 medium. The colour of the aerial mycelia was white on ISP 2, ISP 3 and Czapek's agar media (Table S1). It was determined that strain H2R21^T produces short spore chains with smooth surfaces on aerial mycelia (Fig. 1). The extensively branched substrate mycelium was observed to be in the cream to brown colour series on all tested media. No soluble or melanin pigments were observed on any of the tested media. The strain was found to grow at a temperature range of 20-37 °C and optimally at 28 °C. Growth was observed in the pH range of 6.0-9.0 (optimum at pH 7.0). Strain H2R21^T was found to tolerate NaCl up to 2 % (w/v). Detailed physiological and biochemical properties are given in Table 1 and the species description.

The chemotaxonomic properties of strain H2R21^T were found to be consistent with the characteristics of members of the genus *Nonomuraea* (Lechevalier and Lechevalier, 1970, Kämpfer 2012). The novel strain contains *meso*-diaminopimelic acid in the cell wall peptidoglycan and glucose, mannose, ribose and madurose in whole cell hydrolysates. The main menaquinones were identified as MK-9(H₄) (58.4 %), MK-9(H₆) (9.7 %) and MK-9(H₂) (5.5 %), whilst other components could not be definitively identified. The polar lipids were identified as phosphatidylglycerol, diphosphatidylglycerol, phosphatidylmethylethanolamine, phosphatidylethanolamine, hydroxy-phosphatidylethanolamine, dihydroxy-

174 phosphatidylethanolamine, phosphatidylinositol, glycerophosphatidylinositol, two
175 glycerophospholipids and two unidentified lipids (Fig. S1). The major cellular fatty acids (≥ 10
176 %) were identified as iso-C_{16:0} (26.2 %), iso-C_{16:0} 2OH (24.9 %) and C_{17:0} 10-methyl (12.5 %);
177 the detailed fatty acid profile is listed in Table S2 in comparison with those of closely related
178 type strains. Mycolic acids were absent.

179 Phylogenetic analysis based on 16S rRNA gene sequences of strain H2R21^T and closely related
180 type strains showed that the strain forms a subclade with *N. purpurea* 1SM4-01^T (Fig. 1),
181 which shares high 16S rRNA gene sequence similarity with the strain (99.1 % similarity; 13 nt
182 differences at 1441 sites). The 16S rRNA gene sequence similarity between strain H2R21^T and
183 *N. solani* CGMCC 4.7037^T is 98.4 % (23 nt differences at 1441 sites), and with both
184 *Nonomuraea stahlianthi* SC1-1^T and *N. rhizophila* YIM 67092^T is 98.3 % (24 nt differences at
185 1441 sites and 25 nt differences at 1443 sites, respectively). The association with *N. purpurea*
186 1SM4-01^T was also supported by the maximum-likelihood and maximum-parsimony
187 algorithms (Figs. S2 and S3). DNA-DNA hybridization experiments were carried out between
188 strain H2R21^T and its phylogenetic neighbours, *N. purpurea* 1SM4-01^T, *N. solani* DSM 45729^T
189 and *N. rhizophila* DSM 45382^T. The values between them were 28.0 $\pm$ 2.4 %, 38.1 $\pm$ 3.1 % and
190 40.4 $\pm$ 2.9 %, respectively, which are lower than the cut-off value of 70 %, the recommended
191 threshold value for delineation of same prokaryotic genomic species (Wayne et al. 1987). The
192 DNA G+C content of strain H2R21^T was found to be 69.6 mol %.

193 Both phylogenetic and chemotaxonomic analyses, as well as phenotypic characterisation,
194 indicated that the strain is a member of the genus *Nonomuraea*. However, strain H2R21^T differs
195 from its closely related neighbours in terms of some chemotaxonomic properties, such as the
196 lack of C_{17:1} cis9 in its fatty acid profile, a major cellular fatty acid of *N. purpurea* 1SM4-01^T,
197 and having glucose and mannose in whole cell hydrolysates unlike *N. purpurea* 1SM4-01^T and
198 *N. solani* DSM 45729^T, respectively. In addition, the strain can be differentiated from closely
199 related species in terms of several phenotypic characteristics, such as inability to utilise D-

melezitose, D-melibiose, dextran, L-arabinose, L-glutamine and maltose as sole carbon source, along with other phenotypic characters as given in Table 1.

It is evident on the basis of genotypic and phenotypic properties that strain H2R21^T can be distinguished from its close neighbours *N. purpurea*, *N. solani* and *N. rhizophila*. Therefore, it is proposed that strain H2R21^T be classified in the genus *Nonomuraea* as *Nonomuraea insulae* sp. nov. The Digital Protologue database TaxoNumber for strain H2R21^T is TA00414.

Description of *Nonomuraea insulae* sp. nov.

Nonomuraea insulae sp. nov. (in'su.lae. L. gen. n. *insulae*, belonging to the island of Heybeliada in the Marmara Sea, from where the type strain was isolated).

Aerobic, Gram-stain positive, non-acid-fast, non-motile actinobacterium that forms an extensively branched substrate mycelia. The aerial mycelium is white and forms short spore (0.6-0.8 x 1.2-1.4 µm) chains with smooth surfaces. Good growth occurs on ISP 2, ISP 3, ISP 6, modified Bennett's agar, tryptic soy agar and nutrient agar media, moderate growth on ISP 7 medium and weak growth on ISP 5 and Czapek's agar media; does not grow on ISP 4 medium. No diffusible pigment is observed on any of the above media and melanoid pigments are not produced on ISP 6 or ISP 7 media. Growth occurs at 20–37 °C (optimum at 30 °C), at pH 6.0-9.0 (optimum 7.0) and in the presence of 0-2 % (w/v) NaCl. Allantoin and arbutin are hydrolysed, but not nitrate. Guanine, hypoxanthine, starch and xylan are degraded, but not adenine, xanthine or Tweens 20, 40 and 80. D-Arabinose, D-cellobiose, dextrin, glucose, myo-inositol, D-mannose, mannitol, L-rhamnose, sucrose and xylose are utilised as sole carbon and energy sources, but not L-arabinose, dextran, D-galactose, L-glutamine, D-fructose, inulin, lactose, maltose, D-melezitose, D-melibiose, D-ribose, D-sorbitol or xylitol. L-alanine, L-cysteine, glycine, L-histidine, L-hydroxyproline, Alfa-isoleucine, L-lysine, L-methionine, DL-phenylalanine, L-phenylalanine, L-proline, L-threonine and L-tyrosine are utilised as sole

nitrogen sources, but not glycine, L-arginine, L-asparagine or L-serine. The predominant menaquinone is MK-9(H₄), and MK-9(H₆) and MK-9(H₂) are also present as minor menaquinones. The polar lipids profile includes phosphatidylglycerol, diphosphatidylglycerol, phosphatidylmethylethanolamine, phosphatidylethanolamine, hydroxy-phosphatidylethanolamine, dihydroxy-phosphatidylethanolamine, phosphatidylinositol, glycerophosphatidylinositol, two glycerophospholipids and two unidentified lipids. The whole cell sugars are glucose, mannose, ribose and madurose. The major fatty acids are *iso*-C_{16:0}, *iso*-C_{16:0} 2OH and C_{17:0} 10-*methyl*. The G+C content of the genomic DNA of the type strain is 69.6 mol %.

The type strain, H2R21^T (=DSM 102915^T =CGMCC 4.7338^T =KCTC 39769^T), was isolated from a soil sample from Heybeliada, an island in the Marmara Sea, Turkey. The GenBank accession number for the 16S rRNA gene sequence of the strain is KP027414.

Acknowledgements

This research was supported by Ondokuz Mayıs University (OMU), project no. PYO.FEN.1904.13.004. We gratefully acknowledge Dr. Kannika Duangmal from Kasetsart University in Thailand for help to us in obtaining one of the type species.

Conflicts of Interest

The authors declare that they have no conflict of interest.

References

- Chun J, Goodfellow M (1995) A phylogenetic analysis of the genus *Nocardia* with 16S rRNA gene sequences. *Int J Syst Bacteriol* 45:240–245
- Collins MD (1985) Isoprenoid quinone analysis in bacteria classification and identification. In: Goodfellow M, Minnikin DE (eds) *Chemical methods in bacterial systematics*. Academic Press, London, pp 267–287

250 Dalmastri C, Gastaldo L, Marcone GL, Binda E, Congiu T, Marinelli F (2016) Classification of
 251 *Nonomuraea* sp. ATCC 39727, an actinomycete that produces the glycopeptide antibiotic
 252 A40926, as *Nonomuraea gerenzanensis* sp. nov. Int J Syst Evol Microbiol 66:912–921
 253 De Ley J, Cattoir H, Reynaerts A (1970) The quantitative measurement of DNA hybridization
 254 from renaturation rates. Eur J Biochem 12:143–153
 255 Fang BZ, Hua ZS, Han MX, Zhang ZT, Wang YH, Yang ZW, Zhang WQ, Xiao M, Li WJ
 256 (2017) *Nonomuraea cavernae* sp. nov., a novel actinobacterium isolated from a karst cave
 257 sample Int J Syst Evol Microbiol 67:4692–4697
 258 Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach.
 259 J Mol Evol 17:368–376
 260 Felsenstein J (1985) Confidence limits on phylogeny: an approach using the bootstrap.
 261 Evolution 39:783–791
 262 Hayakawa M, Nonomura H (1987) Humic acid-vitamin agar, a new medium for the selective
 263 isolation of soil actinomycetes. J Ferment Technol 65:501–509
 264 Huss VAR, Festl H, Schleifer KH (1983) Studies on the spectrophotometric determination of
 265 DNA hybridisation from renaturation rates. Syst Appl Microbiol 4:184–192
 266 Jones KL (1949) Fresh isolates of actinomycetes in which the presence of sporogenous aerial
 267 mycelia is a fluctuating characteristic. J Bacteriol 57:141–145
 268 Jukes TH, Cantor CR (1969) Evolution of protein molecules. In: Munro HN (ed) Mammalian
 269 protein metabolism. Academic Press, New York, pp 21–132
 270 Kämpfer P, Kroppenstedt RM (1996) Numerical analysis of fatty acid patterns of coryneform
 271 bacteria and related taxa. Can J Microbiol 42:989–1005
 272 Kämpfer P (2012) Genus VI. *Nonomuraea*. In: Whitman WB, Goodfellow M, Kämpfer P,
 273 Busse H-J, Trujillo ME, Ludwig W, Suzuki KI, Parte A (eds) Bergey's Manual of Systematic
 274 Bacteriology. vol 5: The Actinobacteria, 2nd edn, Springer, New York, pp 1844–1861

275 Kämpfer P, Kroppenstedt RM, Grün-Wollny L (2005) *Nonomuraea kuesteri* sp. nov. Int J Syst
 276 Evol Microbiol 55:847–851
 277 Kelly KL (1964) Inter-society color council-national bureau of standards color-name charts
 278 illustrated with centroid colors. US Government Printing Office, Washington, DC
 279 Kroppenstedt RM (1982) Separation of bacterial menaquinones by HPLC using reverse phase
 280 (RP18) and a silver loaded ion exchanger. J Liquid Chromatogr 5:2359–2387
 281 Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis
 282 version 7.0 for bigger datasets. Mol Biol Evol 33:1870–1874
 283 Lechevalier MP, Lechevalier HA (1970) Chemical composition as a criterion in the
 284 classification of aerobic actinomycetes. Int J Syst Bacteriol 20:435–443
 285 Li Z, Song W, Zhao JW, Zhuang XX, Zhao Y, Xiang WS, Wang XJ (2017) *Nonomuraea*
 286 *glycinis* sp. nov., a novel actinomycete isolated from the root of Black soya bean [*Glycine max*
 287 (L.) Merr]. Int J Syst Evol Microbiol 67:5026–5031
 288 Li J, Zhao GZ, Huang HY, Zhu WY, Lee JC, Xu LH, Kim CJ, Li WJ (2011) *Nonomuraea*
 289 *endophytica* sp. nov., an endophytic actinomycete isolated from *Artemisia annua* L. Int J Syst
 290 Evol Microbiol 61:757–761
 291 Marmur J (1961) A procedure for the isolation of deoxyribonucleic acid from microorganisms.
 292 J Mol Biol 3:208–218
 293 Mandel M, Marmur J (1968) Use of ultraviolet absorbance temperature profile for determining
 294 the guanine plus cytosine content of DNA. Methods Enzymol 12B:195–206
 295 Minnikin DE, O'Donnell AG, Goodfellow M, Alderson G, Athalye M, Schaal A, Parlett JH
 296 (1984) An integrated procedure for the extraction of bacterial isoprenoid quinones and polar
 297 lipids. J Microbiol Methods 2:233–241
 298 Nakaew N, Sungthong R, Yokota A, Lumyong S (2012) *Nonomuraea monospora* sp. nov., an
 299 actinomycete isolated from cave soil in Thailand, and emended description of the genus
 300 *Nonomuraea*. Int J Syst Evol Microbiol 62:3007–3012

301 Nash P, Krent MM (1991) Culture media. In: Balows A, Hausser WJ, Herrmann KL, Isenberg
302 HD and Shadomy HJ (eds) Manual of clinical microbiology, 3rd Edition, American Society for
303 Microbiology, Washington DC, pp 1268–1270

304 Niemhom N, Chutrakul C, Suriyachadkun C, Thawai C (2017) *Nonomuraea stahlianthi* sp.
305 nov., an endophytic actinomycete isolated from the stem of *Stahlianthus campanulatus*. Int J
306 Syst Evol Microbiol 67:2879–2884

307 Rachniyom H, Matsumoto A, Indananda C, Duangmal K, Takahashi Y, Thamchaipenet A
308 (2015) *Nonomuraea syzygii* sp. nov., an endophytic actinomycete isolated from the roots of a
309 jambolan plum tree (*Syzygium cumini* L. Skeels). Int J Syst Evol Microbiol 65:1234–1240

310 Saitou N, Nei M (1987) The neighbor-joining method. A new method for reconstructing
311 phylogenetic trees. Mol Biol Evol 4:406–425

312 Sasser M (1990) Identification of bacteria by gas chromatography of cellular fatty acids.
313 Technical Note 101. Newark, DE: MIDI Inc

314 Shen Y, Jia FY, Liu CX, Li JS, Guo SY, Zhou SY, Wang XJ, Xiang WS (2016) *Nonomuraea*
315 *zeae* sp. nov., isolated from the rhizosphere of corn (*Zea mays* L.). Int J Syst Evol Microbiol
316 66:2259–2264

317 Shirling EB, Gottlieb D (1966) Methods for characterization of *Streptomyces* species. Int J Syst
318 Bacteriol 16:313–340

319 Sripreechasak P, Phongsopitanun W, Supong K, Pittayakhajonwut P, Kudo T, Ohkuma M,
320 Tanasupawat S (2017) *Nonomuraea rhodomycinica* sp. nov., isolated from peat swamp forest
321 soil in Thailand. Int J Syst Evol Microbiol 67:1683–1687

322 Staneck JL, Roberts GD (1974) Simplified approach to identification of aerobic actinomycetes
323 by thin-layer chromatography. Appl Microbiol 28:226–231

324 Suksaard P, Mingma R, Srisuk N, Matsumoto A, Takahashi Y, Duangmal K (2016)
325 *Nonomuraea purpurea* sp. nov., an actinomycete isolated from mangrove sediment. Int J Syst
326 Evol Microbiol 66:4987–4992.

327 Sungthong R, Nakaew N (2015) The genus *Nonomuraea*: a review of a rare actinomycete taxon
 328 for novel metabolites. *J Basic Microbiol* 55:554-565.

329 Waksman SA (1961) *The Actinomycetes, Classification, Identification and Description of*
 330 *Genera and Species*, vol. 2. Baltimore: Williams & Wilkins

331 Waksman SA (1967) *The Actinomycetes. A Summary of Current Knowledge*. New York:
 332 Ronald Press

333 Wang F, Shi JD, Huang YJ, Wu YY, Deng XM (2017) *Nonomuraea ceibae* sp. nov., a new
 334 actinobacterium isolated from *Ceiba speciosa* rhizosphere. *Int J Syst Evol Microbiol* 67:1158–
 335 1162

336 Wang S, Liu C, Zhang Y, Zhao J, Zhang X, Yang L, Wang X, Xiang W (2014) *Nonomuraea*
 337 *guangzhouensis* sp. nov., and *Nonomuraea harbinensis* sp. nov., two novel actinomycetes
 338 isolated from soil. *Antonie van Leeuwenhoek* 105:109–118

339 Wayne LG, Brenner DJ, Colwell RR, Grimont PAD, Kandler O, Krichevsky MI, Moore LH,
 340 Moore WEC, Murray RGE, Stackebrandt E, Starr MP, Trüper HG (1987) Report of the *ad hoc*
 341 committee on reconciliation of approaches to bacterial systematics. *Int J Syst Bacteriol*
 342 37:463–464

343 Williams ST, Goodfellow M, Alderson G, Wellington EMH, Sneath PHA, Sackin MJ (1983)
 344 Numerical classification of *Streptomyces* and related genera. *J Gen Microbiol* 129:1743–1813

345 Xi L, Zhang L, Ruan J, Huang Y (2011) *Nonomuraea maritima* sp. nov., isolated from coastal
 346 sediment. *Int J Syst Evol Microbiol* 61:2740–2744

347 Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J (2017) Introducing EzBioCloud: a
 348 taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int*
 349 *J Syst Evol Microbiol* 67:1613–1617

350 Zhang Z, Wang Y, Ruan J (1998) Reclassification of *Thermomonospora* and *Microtetraspora*.
 351 *Int J Syst Bacteriol* 48:411–422

352

Table 1. Differential characteristics of strain H2R21^T and type strains of closely related species of the genus *Nonomuraeae*. Strains: **1**, H2R21^T; **2**, *N. purpurea* 1SM4-01^T; **3**, *N. solani* DSM 45729^T; **4**, *N. rhizophila* DSM 45382^T. Symbols: +, positive; -, negative.

Characteristics	1	2	3	4
Maximum NaCl tolerance (% w/v)	2	2	2	7
pH tolerance	6.0-9.0	7.0-9.0	7.0-9.0	6.0-8.0
Biochemical Tests				
Allantoin	+	-	+	+
Degradation of (% w/v)				
Hypoxanthine (0.4 %)	+	-	-	+
Starch (1%)	+	-	+	-
Guanine (0.05 %)	+	-	+	-
Carbon source utilization (1.0 %, w/v)				
D-Arabinose	+	+	+	-
D-Fructose	-	+	+	+
D-Sorbitol	-	+	+	-
D-Galactose	-	+	+	-
D-Mannose	+	-	+	+
D-Melezitose	-	+	+	+
D-Melibiose	-	+	+	+
D-Ribose	-	+	-	-
Dextran	-	+	+	+
Inulin	-	-	+	+
L-Arabinose	-	+	+	+
Lactose	-	-	+	+
L-Glutamine	-	+	+	+
Maltose	-	+	+	+
Sucrose	+	+	+	-
Xylitol	-	-	+	+
Xylose	+	+	+	-
Use of sole nitrogen sources 0.1 % (w/v)				
Glycine	+	-	+	+
L-Alanine	+	-	+	+
L-Arginine	-	-	+	+
L-Asparagine	-	-	+	+
L-Cysteine	+	-	+	+
L-Hydroxyproline	+	-	+	+
L-Serine	-	-	+	+
Major menaquinones (> 10%)	MK-9(H ₄)	MK-9(H ₄) ^a	MK-9(H ₄), MK-9(H ₆) ^b	MK-9(H ₄), MK-9(H ₆) ^c
Polar lipid profile	DPG, PME, PE, OH-PE, 2OH-PE, PI, GPI	DPG, OH-PE, OH-PME, PE, PG, PI, PIM ^a	DPG, PME, PE, OH-PME, OH-PE, PI, PIM ^b	DPG, PME, PE, PG, PI ^c
Major fatty acids (> 10%)	<i>iso</i> -C _{16:0} , <i>iso</i> -C _{16:0} 2OH	C _{17:1 cis} 9, <i>iso</i> -C _{16:0} 2OH, <i>iso</i> -C _{16:0}	<i>iso</i> -C _{16:0} , C _{17:0} 10-methyl, <i>iso</i> -C _{16:1} G, <i>iso</i> -C _{16:0} 2OH	C _{17:0} 10-methyl, <i>iso</i> -C _{16:0} 2OH
Whole-cell sugars	Glu, Man, Mad, Rib	Mad, Man, Rib ^a	Mad, Glu ^b	Mad, Glu, Man, Rib, Gal ^c

Abbreviations: DPG, diphosphatidylglycerol; PME, phosphatidylmethylethanolamine; PE, phosphatidylethanolamine; OH-PE, hydroxy-phosphatidylethanolamine; 2OH-PE, dihydroxy-phosphatidylethanolamine; PI, phosphatidylinositol; GPI,

358 glycophosphatidylinositol; OH-PME; hydroxy-phosphatidylmonomethylethanolamine; PG, phosphatidylglycerol; PIM,
359 phosphatidylinositol mannoside; Glu, glucose; Man, mannose; Mad, madurose; Rib, ribose; gal, galactose.
360 Data taken from: ^aSuksaard et al. (2016); ^bWang et al. (2013); ^cZhao et al. (2011).

361

362

Fig. 1

Fig. 2.

Figure captions

Fig. 1. Scanning electron micrograph of strain H2R21^T grown on ISP 3 agar after 4 weeks at 28°C showing aerial mycelium differentiating into smooth surfaces oval-shaped spore chains. Bar, 2 µm.

Fig. 2. Neighbour-joining tree based on almost complete 16S rRNA gene sequences showing the position of strain H2R21^T within members of the genus *Nonomuraea*. Asterisks indicate branches of the tree that were also recovered using the maximum-likelihood and maximum-parsimony tree-making algorithms. Numbers at the nodes indicate levels of bootstrap support (%); only values ≥ 50% are shown. *Thermopolyspora flexuosa* DSM 43186^T (AY039253) was used as the outgroup. The GenBank accession numbers are given in parentheses. Bar, 0.005 substitutions per site.

Supplementary Fig. S1. Two-dimensional TLC of polar lipids from strain H2R21^T. For detection of polar lipids, molybdophosphoric acid was applied. DPG, diphosphatidylglycerol; PME, phosphatidylmethylethanolamine; PE, phosphatidylethanolamine; OH-PE, hydroxyphosphatidylethanolamine; 2OH-PE, dihydroxyphosphatidylethanolamine; PI, phosphatidylinositol; GPI, glycerophosphatidylinositol; 2UL, two unidentified lipids; 2PL, two unidentified phospholipids.

Supplementary Fig. S2. Maximum-likelihood tree based on almost complete 16S rRNA gene sequences showing the position of strain H2R21^T with the *Nonomuraea* gene tree. Numbers at the nodes indicate levels of bootstrap support (%), only values ≥50 % are shown. Bar, 0.005 substitutions per nucleotide position.

Supplementary Fig. S3. Maximum-parsimony tree based on almost complete 16S rRNA gene sequences showing the position of strain H2R21^T with the *Nonomuraea* gene tree. Numbers at the nodes indicate levels of bootstrap support (%), only values ≥50 % are shown.

389 **Supplementary Table S1.** Cultural characteristics of the novel strain H2R21^T and closely
390 related type strains grown at 28 °C for 14 days.

391 **Supplementary Table S2.** Cellular fatty acids profiles of strain H2R21^T and type strains of
392 closely related species.