Cochrane
Library

Cochrane Database of Systematic Reviews

Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders (Review)

Pennington L, Akor WA, Laws K, Goldbart J

Pennington L, Akor WA, Laws K, Goldbart J.

Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders.

Cochrane Database of Systematic Reviews 2018, Issue 7. Art. No.: CD012507.

DOI: 10.1002/14651858.CD012507.pub2.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	4
BACKGROUND	7
OBJECTIVES	9
METHODS	9
RESULTS	12
Figure 1.	14
Figure 2.	17
DISCUSSION	19
AUTHORS' CONCLUSIONS	20
ACKNOWLEDGEMENTS	21
REFERENCES	21
CHARACTERISTICS OF STUDIES	27
DATA AND ANALYSES	36
Analysis 1.1. Comparison 1 Parent-mediated intervention versus no intervention: child communicative participation outcomes, Outcome 1 Initiating conversation.	37
Analysis 1.2. Comparison 1 Parent-mediated intervention versus no intervention: child communicative participation outcomes, Outcome 2 Joint attention.	37
Analysis 2.1. Comparison 2 Parent-mediated intervention versus no intervention: parents' communication and interaction strategies, Outcome 1 Maternal Behavior Rating Scale (MBRS) Responsiveness.	38
Analysis 2.2. Comparison 2 Parent-mediated intervention versus no intervention: parents' communication and interaction strategies, Outcome 2 MBRS Directiveness.	38
Analysis 3.1. Comparison 3 Parent-mediated intervention versus no intervention: family stress and coping, Outcome 1 Parent Stress Index: child temperament.	39
Analysis 3.2. Comparison 3 Parent-mediated intervention versus no intervention: family stress and coping, Outcome 2 Parent Stress Index: parent-child relationship.	39
Analysis 3.3. Comparison 3 Parent-mediated intervention versus no intervention: family stress and coping, Outcome 3 Parent Stress Index: learning expectation.	40
ADDITIONAL TABLES	40
APPENDICES	42
CONTRIBUTIONS OF AUTHORS	59
DECLARATIONS OF INTEREST	60
SOURCES OF SUPPORT	60
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	61

Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders

Lindsay Pennington¹, Wanwuri A Akor¹, Kate Laws², Juliet Goldbart³

¹Institute of Health and Society, Newcastle University, Newcastle upon Tyne, UK. ²Regional Communication Aid Service, Northumberland, Tyne and Wear NHS Foundation Trust, Walkergate Park Centre for Neuro-Rehabilitation and Neuro-Psychiatry, Newcastle upon Tyne, UK. ³Faculty of Health, Psychology & Social Care, Manchester Metropolitan University, Manchester, UK

Contact address: Lindsay Pennington, Institute of Health and Society, Newcastle University, Sir James Spence Institute - Royal Victoria Infirmary, Queen Victoria Road, Newcastle upon Tyne, NE1 4LP, UK. lindsay.pennington@ncl.ac.uk.

Editorial group: Cochrane Developmental, Psychosocial and Learning Problems Group.

Publication status and date: New, published in Issue 7, 2018.

Citation: Pennington L, Akor WA, Laws K, Goldbart J. Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders. *Cochrane Database of Systematic Reviews* 2018, Issue 7. Art. No.: CD012507. DOI: 10.1002/14651858.CD012507.pub2.

Copyright © 2018 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Children with motor disorders can have difficulties in producing accurate and consistent movements for speech, gesture or facial expression (or a combination of these), making their communication difficult to understand. Parents may be offered training to help recognise and interpret their child's signals and to stimulate their children's development of new communication skills.

Objectives

To assess the effectiveness of parent-mediated communication interventions, compared to no intervention, treatment as usual or clinician-mediated interventions, for improving the communication skills of preschool children up to five years of age who have non-progressive motor disorders.

Search methods

We searched CENTRAL, MEDLINE, Embase, CINAHL, PsycINFO, 12 other databases and three trials registers in July 2017. We also searched the reference lists of relevant papers and reviews, and contacted experts working in the field to find unpublished studies.

Selection criteria

We included studies that used randomised or quasi-randomised designs; compared a parent-mediated communication intervention with no treatment, treatment as usual or clinician-mediated therapy; and included children with non-progressive motor disorders up to five years of age.

Data collection and analysis

We used the standard methodological procedures expected by Cochrane.

Main results

This review included two randomised controlled trials involving 38 children (20 boys, 18 girls), aged 15 to 96 months, and their mothers. All children had developmental disabilities; 10 had motor disorders, but it was unclear if these motor disorders affected their gestural, vocal or verbal communication. Mothers attended eight group training sessions over 11 to 12 weeks and received two or three home visits. Outcomes were assessed immediately after training. We found no report of longer-term follow-up. One study took place at an intervention centre in Canada and the other in South Korea.

Both studies recruited small numbers of participants from single centres. Since it is not possible to blind participants attending or therapists providing training to group allocation, we considered both studies to be at high risk of performance bias. We also rated one study at high risk of attrition bias, and both studies at low risk of reporting bias.

There was very low-quality evidence for all outcomes assessed. There was no evidence of an effect of training for children's initiation of conversation or engagement in joint attention during interaction with their mothers. Mothers who received training became more responsive to their children's communication, but there were no differences in the extent to which they controlled conversation by directing their children. Missing data meant that we were unable to evaluate the effects of training on children's frequency of communication, frequency of spoken language in conversation, speech production, or receptive or expressive language development. There were no effects on maternal stress. We found no reports of the effects of parent training on children's use of individual communication skills, such as asking questions or providing information, on their generic participation or adverse outcomes. Neither did we find reports of mothers' satisfaction with treatment, its acceptability or their compliance with it.

Authors' conclusions

There is only limited, very low quality evidence that parent-mediated communication interventions may be associated with improvements in interaction between mothers and their preschool children who have motor disorders. The indirectness of the study samples and high risk of bias in the included studies significantly limits our confidence in the evidence, as do issues with study design and lack of detail in results. It is not clear if training has been tested with children whose motor disorders limit the consistency and accuracy of movements underpinning spoken or gestural communication. Some speech and language therapists currently provide communication training for parents. Further research, with larger numbers of children whose movement disorders affect their speech and gestures, coupled with detailed reporting of children's baseline skills, is needed to test whether communication training for parents can help them to promote the communication development of their young children with movement disorders.

PLAIN LANGUAGE SUMMARY

Communication training for parents of preschool children with motor disorders

Background

Children with movement disorders, such as cerebral palsy, often have difficulty producing speech and gesture. This can make their communication difficult to understand. In the preschool years, speech and language therapy often involves training parents to recognise their child's communication signals and promote communication development.

Review question

Does communication training for parents (parent-mediated communication intervention) of preschool children with movement disorders improve the communication between children and parents? We were also interested in whether the training had any unintended consequences, whether it had an effect on parents' levels of stress and coping, and whether parents were satisfied with the training and complied with it.

Study characteristics

We searched for studies published up to July 2017. We found only two studies that reported the effects of parent communication training; one study took place at an intervention centre in Canada, the other in South Korea. The studies involved 38 children (20 boys, 18 girls), aged 15 to 96 months, and their mothers. Both studies compared parent communication training with no intervention for communication problems. Mothers attended eight group training sessions 11 to 12 weeks with two or three home visits. The studies involved children with a range of developmental difficulties; most had intellectual disability, 10 had movement disorders (cerebral palsy). However, the extent to which children's movement disorder affected their communication was not clear; all children appeared to have good use of their hands for gesture and pointing, and impairment of speech was not reported.

Results were assessed immediately after training. We found no report of results at a later date (longer-term follow-up).

Key results and the quality of the evidence

In the two small studies, it appears that mothers may have responded more frequently to their child's interaction following parent-mediated communication training. However, there was no associated reduction in mothers' directiveness (such as their use of commands) in conversation and no change in maternal stress. For the children, we found no evidence for change in children's initiation of conversation or of joint attention in interaction with others. Studies did not report any negative effects of training, mothers' adherence to guidance within the training or the acceptability of the programmes.

We were not able to evaluate the effects of parent-mediated communication intervention and frequency of children's communication, their use of spoken language in conversation with their parents, their speech production or their language development because the data were not available. We have no reports of children's development of individual communication skills, such as learning to ask questions, and no reports of defects of the intervention on their generic participation or harms arising from the intervention. Finally, we found no reports of maternal satisfaction with the treatment.

We judged the evidence from the included studies to be of very low quality because of issues with study design and a lack of detail in the results presented, and because it was not clear whether children's movement disorders affected their communication.

Research with larger numbers of families of children whose movement disorders affect their speech and gesture is needed, to test whether communication training for parents can help them to promote the communication development of their young children with movement disorders.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Parent-mediated communication intervention compared with no intervention for improving the communication skills of preschool children with non-progressive motor disorder				
Patient or population: children up to 5 years of age with a communication difficulty associated with any non-progressive motor disorder acquired before 2 years of age Settings: speech and language therapy clinic Intervention: parent-mediated communication intervention on promoting communication development Comparison: no intervention				
Outcomes	Impact	Number of participants (studies)	Quality of the evidence (GRADE)	Comments
Children’s ability to communicate effectively in everyday life: taking turns in conversation, initiating conversation and responding to others’ conversational gambits Assessed with: observational scales of child interaction - Child Behavior Rating Scale (CBRS; Mahoney 1999b) (therapist rating of child engagement with parent in play in 7 domains (interest, attention to activity,persistence,initiation, affect, joint attention, co-operation). Items scored 1 to 5 on Likert scale; lower scores represent lower levels of coded behaviours) Follow-up: postintervention (time not specified)	1 study found no evidence of effect of parent-mediated intervention on children’s initiation of conversation (MD −0.25 points, 95% CI −1.01 to 0.51; P = 0.54), or joint attention during interaction (MD 0.05 points, 95% CI −0.52 to 0.62; P = 0.87)	18 (1 study)	⊕○○○ Very low ^{a,b,c}	The CBRS involved a subjective rating of child’s engagement in interaction. For this review, we were interested in the initiation and joint attention domains of the CBRS, as these relate directly to communication
Adverse events	Not measured			
Child outcomes				

Speech and language function Assessed with: Sequenced Inventory of Communication Development (Hedrick 1975) (age-normed assessment (4-48 months); lower score represents lower levels of communication development) Follow-up: postintervention (time not specified)	1 study measured the effect of 20 (1) parent-mediated communication intervention on children's speech and language function but missing data prevented analysis	⊕○○○ Very low ^{a,b}	1 included study reported a general communication ability measure
Children's generic participation: social	Not measured		
<i>Parent outcomes</i>			
Parents' communication and interaction strategies: directiveness and responsiveness Assessed with: Maternal Behavior Rating Scale (MBRS; Mahoney 1999a) (therapist rating of maternal behaviour on 12 items in 4 domains (responsiveness, affect, achievement orientation and directiveness). Items scored 1-5 on Likert scale; lower scores represent lower levels of coded behaviours) Follow-up: postintervention (time not specified)	1 study found increases in parents' responsiveness (MD 1.02 points, 95% CI 0.34 to 1.70; P = 0.01), but no effect on their directiveness (MD 0.49 points, 95% CI -0.02 to 1.00; P = 0.07)	⊕○○○ Very low ^{a,b,c}	The MBRS involves a subjective rating of parent's engagement in, and control of, interaction with the child. For this review, we were interested in the MBRS domains of responsiveness and directiveness, as these relate directly to the communication opportunities provided to children by parents in conversation
Family stress and coping: parent stress Assessed with: Korean version of the Parent Stress Index (Shin 1997) (parent-reported measure. Total Stress score calculated	1 study found no effect of parent-mediated training in reducing parental stress related to their child's temperament (MD -6.50 points, 95% CI -11.64 to 1.36), parent-child relationship (MD 0.	⊕○○○ Very low ^{a,c}	1 included study reported outcomes regarding the potential impact of communication difficulties on parents' psychological well-being. Prior to intervention, parents in the training group ap-

from 20 items in each of 3 scales (child temperament, parent-child relationship, learning expectation). Each item rated on 5-point Likert scale to indicate the extent to which items disturbed the parent in last week; lower scores indicate lower levels of disturbance) Follow-up: postintervention (time not specified)	25 points, 95% CI -4.70 to 5.20; P = 0.92), or learning expectation (MD -0.58 points, 95% CI -2.87 to -1.71)	peared to have higher levels of stress regarding their relationship with their child and lower levels of stress regarding their child's temperament than parents in the control group
---	--	---

Satisfaction of child and family with treatment	Not measured
Compliance with treatment	Not measured

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI)
CI: confidence interval; **CBRS**: Child Behavior Rating Scale; **MD**: mean difference

GRADE Working Group grades of evidence
High quality: we are very confident that the true effect lies close to that of the estimate of the effect.
Moderate quality: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.
Low quality: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.
Very low quality: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect

^aDowngraded by two levels due to risk of bias, and one level for indirectness. Risk of selection bias was unclear. No information was given on the allocation of participants to groups, although in one study, Kim 2005, parents and children in the parent-mediated intervention appeared to have higher levels of interaction difficulties than those in the control group prior to intervention. All children in the studies had developmental delays. Four children in Girolametto 1988 and six children in Kim 2005 had motor disorders (e.g. cerebral palsy). Participants in both studies were able to use their hands. It was not clear how far the results extended to children with motor disorder that impairs gesture and speech.

^bRisk of bias. Girolametto 1988 did not state if outcome assessors were blind to group or timing of data collection.

^cRisk of bias. Kim 2005 blinded outcome assessors to group and timing of data collection.

BACKGROUND

Description of the condition

Non-progressive motor disorders in childhood arise from a variety of conditions, including cerebral palsy, acquired brain injury, global developmental delay, Down's syndrome and genetic mutations. Exactly how many children are affected is currently unknown due to sparse population-level data. The most comprehensive data come from international surveillance of cerebral palsy. Cerebral palsy is defined as "a group of permanent disorders of the development of movement or posture, causing activity limitation, that are attributed to non-progressive disturbances that occurred in the developing fetal or infant brain" (Rosenbaum 2007). Registries have shown that cerebral palsy affects two or three children per 1000 children in high-income countries (Cans 2008; Kirby 2011; Reid 2011). Prevalence is likely to be greater in low- and middle-income countries where health care is less abundant, but cerebral palsy rates in these regions have not yet been ascertained. One Dutch study estimated that, in children under 14 years of age, there are about 3.6 new cases of severe acquired brain injury per 100,000 children per annum (De Kloet 2013). However, it is unclear how many of these children have ongoing motor disorders. Again, the incidence may be greater in low- and middle-income countries with less access to maternal and child health services. Global developmental delay and Down's syndrome lead to low motor tone and slow acquisition of motor skills, and global developmental delay affects approximately 39 per 1000 children and Down's syndrome affects 1.4 per 1000 children (Boyle 2011; Parker 2010). Genetic mutations that cause motor disorders include the PRRT2 (proline-rich transmembrane protein 2) mutation and the GLUT-1 (glucose transporter type 1) syndrome (Blackburn 2012), but their prevalence is unclear. Many developmental disabilities, including those causing non-progressive motor disorders included in this review, are more common in boys and in families in who live in poverty (Boyle 2011). Disorders are diagnosed by paediatricians, paediatric neurologists and geneticists. Differential diagnosis may take some years due to the slowly evolving nature of some conditions.

Motor disorders impair the range, speed, strength and consistency of movements. When disorders affect the movements underpinning vocalisation, speech, gesture or facial expression (or a combination of these), parents and other carers find it difficult to recognise and interpret children's attempts to communicate, and this can lead to a breakdown in interaction (Hanzlik 1990; Light 1985; Pennington 2001). To promote effective interaction, parents may structure conversations around the children's communication signals that are easy to understand (Dunst 1985; Tannock 1992). However, this can lead to asymmetrical interaction, with parents introducing topics, asking forced, choice questions and then acknowledging their child's response. Such an uneven, parent-led

pattern of conversation can make it difficult for children with motor disorders to learn new communication skills.

It is estimated that around 20% to 30% of children with cerebral palsy have no functional speech and a further 22% are able to speak but have intelligibility limitations due to their motor disorders (ACPR 2016; Nordberg 2013; Parkes 2010). Speech disorder is most likely to occur in bilateral spastic-type cerebral palsy (where muscle tone is increased on both sides of the body), in dyskinetic forms (where muscle tone fluctuates) and in ataxic forms (characterised by tremor and difficulties with muscular co-ordination) (Bax 2006; Parkes 2010). The prevalence of speech disorders in other conditions leading to non-progressive motor disorders is currently unknown.

Children with motor disorders who also have a cognitive impairment may take longer to reach milestones, such as intentionality and engaging in joint attention with another person, which are vital for interaction, and the development of linguistic understanding may be delayed. About 49% of children with cerebral palsy have an intellectual disability (IQ less than 70) and 28% have a severe intellectual disability (IQ less than 50) (Novak 2012). Current research suggests that receptive language is largely commensurate with cognitive development in cerebral palsy (Pirila 2007), but further epidemiological studies are needed to confirm this. Approximately 12% of children with cerebral palsy have hearing impairment and 35% have some visual impairment (ACPR 2016), both of which can influence communication development.

Communication difficulties have a profound impact on children's family, social and educational life. Children with communication and motor disorders are more at risk of lower quality of life and restricted social participation than their peers with and without motor disorders (Dickinson 2007; Fauconnier 2009). The impact of communication breakdown is felt throughout families, and parents report high levels of stress (Parkes 2011; Pousada 2013).

As differential diagnosis may not be possible in early childhood, and all motor disorders affecting speech and gesture can lead to intelligibility limitations, this review will be inclusive of all causes of non-progressive motor disorders in the preschool years. One exception to this is children with Down's syndrome, who will be considered in a separate review (O'Toole 2016). Thus, we will exclude studies examining only children with Down's syndrome, but will include studies in which Down's syndrome is one of a range of disorders causing motor impairment. Degenerative disorders, such as muscular dystrophies and metabolic disorders, may also be associated with motor impairment and may become apparent after a period of healthy development. As these disorders lead to a loss of skills rather than development following an atypical pattern, as is the case for children with non-progressive disorders, they were excluded from this review. Also, children with severe hearing or visual impairments, or both, have specific difficulties acquiring early interaction skills arising from their differences in processing communication signals, which are beyond the scope of this review.

Description of the intervention

Communication skills are developed in interaction, and children's most frequent communication partners are their parents, so therapy involves training parents to adapt their communication style. This is referred to as parent-mediated or indirect therapy. It aims to help parents of children with motor disorders to recognise and interpret their children's idiosyncratic attempts to communicate and to stimulate their child's development of new skills (e.g. Bruno 1998; Kaiser 1987; Kent-Walsh 2015; Mahoney 1988; Pepper 2004; Yoder 2002). Training is most often provided by speech and language therapists and other personnel with an interest in interaction (e.g. psychologists and early-years educators). It can be delivered to individuals or groups of parents and may take place in parents' homes or in health, education or social care settings. Training often teaches parents about how communication develops, from preintentional communication through to non-verbal, intentional communication and then on to linguistic communication (Girolametto 1986; Hemmeter 1994; Pepper 2004). It covers the purposes for which communication is used, how communication involves communication partners taking turns in expressing and receiving signals (Hemmeter 1994; Mahoney 1988; Pepper 2004), and how messages are coconstructed by communication partners (Clarke 2017). Techniques to aid children's language and communication development are introduced, including how to create simulating environments, promote a need to communicate, allow sufficient time for children to join or start conversations, respond contingently to children's messages, and repeat language to aid association between spoken words and their referents. Parents are encouraged to apply this information to their interaction with their children (Fey 2006; Gibbard 2004; Girolametto 1996; Kaiser 2001; Mahoney 1988; Pepper 2004). Training often includes one-to-one coaching, whereby therapists watch the interaction between a parent and a child (either in real time or on video) and highlight which behaviours prompted the child to communicate so the parent can repeat these in daily interactions (Kaiser 1995; Kaiser 2003; McDuffie 2016; Pepper 2004). Coaching might also involve the parent watching the therapist modelling the interaction with the child (Kaiser 2003; Pepper 2004). Young children who have severe speech impairment associated with their motor disorders may be introduced to augmentative and alternative communication (AAC) to supplement or replace their natural forms of communication. AAC includes signing, gesture and use of body movements (unaided AAC), or may introduce equipment such as objects to represent daily activities, photographs, pictures, symbols, speech generating devices and mobile technology communication applications (aided AAC). AAC provides access to a wider range of vocabulary and language but takes children and their parents time to learn. Conversation using AAC often involves the coconstruction of meaning by parents and children as children have access to a restricted vocabulary (Clarke 2017; Solomon-Rice 2011). Training is often provided for parents to teach them how to accommodate the use of the system in

conversation and help their children to produce new vocabulary and language structures via the AAC system (Kent-Walsh 2015). Such training may be incorporated in the generic communication training described above (Pennington 2009), or be provided separately in programmes that focus specifically on AAC (Kent-Walsh 2015).

How the intervention might work

Parent communication training is based on the transactional theory of development, which hypothesises that children and their parents continuously adapt to each other's behaviours (Sameroff 2000). Children with motor disorders communicate using movements and vocalisations/words that may look and sound very different to those of their typically developing peers. Following the transactional hypothesis, helping parents to recognise and interpret their children's idiosyncratic communication behaviours should lead to parents' responding appropriately to children's messages. Such positive responses should, in turn, encourage children to repeat these behaviours and communicate more frequently. Teaching parents about how communication develops should enable them to apply this knowledge to continue to adapt their own communication patterns and provide ongoing support for their children's development from the beginnings of intentional communication to the understanding and use of multi-word utterances (Girolametto 1996). When children are just beginning to communicate intentionally, parents would follow their children's focus of attention and give their children more time to start interactions and produce messages. Children would be encouraged to jointly attend to an object/activity with their parents. Hearing their parents say the names of objects and actions during joint attention would help children to associate the spoken words with their referents to learn language. Over time, parents would introduce more varied play and communication opportunities and hence a need for children to communicate for a greater range of purposes, with a wider range of vocabulary and language structures. For children with limited spoken output, such vocabulary would be supported by the use of AAC; for example by parents pointing to symbols or pictures as well as speaking. In these approaches to therapy, there is an implicit or explicit assumption that changing parents' interaction style will have an effect on children's communication, expressive or receptive language (or both), and interaction (Pickstone 2009). The intervention may also serve to increase parents' confidence in their communication with their children, reduce parental stress as communication breakdowns become less frequent, and help children to interact successfully in a greater number of social activities and with a broader range of people.

Why it is important to do this review

The James Lind Alliance Childhood Disability Research Priority Setting Partnership rated the timing and intensities of interventions, and the effectiveness of communication interventions as the two most important areas for investigation (Morris 2015). Internationally, there has been a drive in research to develop early interventions to maximise the potential skill development associated with brain plasticity in infancy and the early years. Early communication intervention has often focused on training parents to provide parent-mediated intervention (Watson 2015). Previous Cochrane Reviews have considered parent-mediated intervention training programmes for children with autism (Oono 2013), and primary speech and language delay or disorder (Law 2003), and a current review is investigating for parent-mediated intervention children for with Down's syndrome (O'Toole 2016). However, the method of delivery of parent communication training, its contents, dosage and suitability for families of children with motor disorders have not been evaluated. A previous review considered speech and language therapy interventions to improve the communication skills of children with cerebral palsy and included parent-mediated interventions (Pennington 2003); its authors identified one randomised controlled trial (RCT) of a parent-mediated communication intervention. This review updates the section of the previous review that examined training delivered to parents of children with cerebral palsy (Pennington 2003), to identify new empirical data. It also considers interventions provided to parents of preschool children with other non-progressive motor disorders, as their communication development is similarly affected. By including all children with non-progressive motor disorders, we aimed to examine the generic effectiveness of parent-mediated training interventions in the preschool period, extending the utility of the review to service providers and policymakers.

OBJECTIVES

To assess the effectiveness of parent-mediated communication interventions, compared to no intervention, treatment as usual or clinician-mediated interventions, for improving the communication skills of preschool children up to five years of age who have non-progressive motor disorders.

METHODS

Criteria for considering studies for this review

Types of studies

RCTs, including cluster-RCTs, and quasi-RCTs in which participants were allocated to intervention groups by methods that were not strictly random (e.g. date of birth).

Types of participants

Children up to five years of age who had a communication difficulty associated with any non-progressive motor disorder acquired before two years of age. We included children with additional intellectual impairments, including children with Down's syndrome, children whose vision was corrected by spectacles and children whose hearing was amplified by hearing aid(s) if they had identified motor difficulties. We included studies of children with intellectual/developmental disability, where at least one child in each group had additional motor disorders, as other participants would be likely to have delayed motor skills. Subgroup analysis for children with motor disorders was not mandatory. We excluded studies of only children with Down's syndrome, as they are being considered by another review (see O'Toole 2016). We also excluded children whose communication was primarily limited by a sensory impairment, as their communication development differs from children who can see and hear the world around them. We inferred motor disorder from descriptions of children's development and confirmed this with study authors, where possible. Where information about this was provided, children's communication difficulty was diagnosed by speech and language therapists or psychologists.

Parents of the children described above.

Types of interventions

We considered studies of training delivered to parents with the explicit aim of helping them to promote their child's communication development. Training could have been delivered to parents individually or in groups. Training could have been delivered by speech and language therapists, psychologists, early educators or others with specialist knowledge of communication. Training could have taken place in the home or in health, education or community support settings. Training programmes could have varied in dosage (intensity, frequency and duration). Training could have included communication via AAC as one mode of communication in a total communication approach or focus primarily on communication using AAC. We excluded facilitated communication, as it is not a valid form of AAC (Schlosser 2014).

Comparisons of interest were: parent-mediated intervention versus no intervention or waiting-list controls; parent-mediated intervention versus treatment as usual (e.g. multidisciplinary therapy groups providing motor, sensory and language stimulation); parent-mediated intervention versus clinician-mediated intervention.

Types of outcome measures

Primary outcomes

1. Children's ability to communicate effectively in everyday life. Outcomes included children's ability to:

- i) take turns in conversation, initiating conversation and responding to others' conversational gambits;
- ii) use communication for a wide range of purposes such as requesting attention, asking questions, answering questions, making comments and repairing conversation when they have not been understood; and
- iii) use a range of modes of expression by vocalising, speaking, using gesture or using the AAC system.

2. Adverse events, including reductions in the frequency with which children communicate, or increases in negative behaviour. Outcomes could have been measured at the level of activity (i.e. the ability to execute a task), and at the level of participation (i.e. communication in life situations) (WHO 2001).

Measures could have included rating scales (e.g. Focus on the Outcomes of Communication Under Six (Thomas-Stonell 2010) and Therapy Outcome Measures (Enderby 2015)), communication assessments (e.g. Communication and Symbolic Behavior Scales (Wetherby 2002)), and observational coding schemes (e.g. frequency counts of children's initiations and responses in interaction).

Secondary outcomes

1. Child outcomes:

- i) speech and language function, assessed using standardised measures of children's expressive and receptive language skills and speech production (e.g. Pre-school Language Scales (Zimmerman 2002); Communicative Development Inventory (CDI; Fenson 2006); Receptive-Expressive Emergent Language Scale-3 (REEL-3; Bzoch 1970)); non-standardised assessments of gestural ability; or production of messages using AAC on demand, measured using coding schemes developed for individual research studies that include validity and reliability data; and

- ii) children's generic participation, assessed using validated measures (Assessment of Life Habits (Noreau 2007), and Children's Assessment of Participation and Enjoyment (King 2004)).

2. Parent outcomes:

- i) parents' communication and interaction strategies, assessed using non-standardised measures (Responsive Augmentative and Alternative Communication Style (Broberg 2012); coding schemes that measured the frequency of parent communication behaviours (e.g. initiations of conversation; directives) developed for individual research studies that included validity and reliability data);

- ii) family stress and coping (e.g. Questionnaire on Resources and Stress (Friedrich 1983) or Carer Strain Index (Robinson 1983));

- iii) satisfaction of child and family with treatment (e.g. rating scales developed for individual studies, Patient Satisfaction Questionnaire Short Form (PSQ-18; Marshall 1994)); and

- iv) compliance with treatment (e.g. number of sessions missed and reasons for this).

See Table 1 for methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions.

Search methods for identification of studies

Electronic searches

In July 2017, we searched the electronic databases and trials registers listed below, from inception onwards.

1. Cochrane Central Register of Controlled Trials (CENTRAL; 2017, Issue 6) in the Cochrane Library, which includes the Cochrane Developmental, Psychosocial and Learning Problems Specialised Register (searched 26 July 2017).
2. MEDLINE Ovid (1946 to July week 2 2017).
3. MEDLINE In-Process & Other Non-Indexed Citations Ovid (searched 25 July 2017).
4. MEDLINE Epub Ahead of Print Ovid (searched 25 July 2017).
5. Embase Ovid (1974 to 2017 week 30).
6. CINAHL Plus EBSCOhost (Cumulative Index to Nursing and Allied Health Literature; 1937 to 26 July 2017).
7. PsycINFO Ovid (1806 to July week 3 2017).
8. Science Citation Index Web of Science (SCI; 1970 to 26 July 2017).
9. Conference Proceedings Citation Index - Science Web of Science (CPCI-S; 1970 to 26 July 2017).
10. Language and Linguistic Behaviour Abstracts ProQuest (LLBA; 1871 to 3 August 2017).
11. British Education Index EBSCOhost (BEI; 1929 to 26 July 2017).
12. ERIC EBSCOhost (Education Resources Information Center; 1966 to 26 July 2017).
13. *Cochrane Database of Systematic Reviews* (CDSR; 2017, Issue 7) part of the Cochrane Library (searched 26 July 2017).
14. Database of Abstracts of Reviews of Effects (DARE; 2015, Issue 2) part of the Cochrane Library (searched 26 July 2017).
15. LILACS (Latin American and Caribbean Health Science Information database; lilacs.bvsalud.org/en; searched 26 July 2017).
16. Rehabdata National Rehabilitation Information Center (narric.com; searched 27 July 2017).
17. SpeechBITE (speechbite.com; searched 27 July 2017).
18. ClinicalTrials.gov (clinicaltrials.gov; searched 27 July 2017).
19. UK Clinical Trials Gateway (www.ukctg.nihr.ac.uk/clinical-trials; searched 27 July 2017).

20. World Health Organization International Clinical Trials Registry Platform (WHO ICTRP; www.who.int/ictip/en; searched 27 July 2017).

Search strategies for each source are in [Appendix 1](#). We did not limit the search by the country in which the research was undertaken, the language in which the research was reported, year of publication or publication status. See [Table 1](#) for methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions.

Searching other resources

We handsearched the reference lists of relevant papers and reviews for studies not identified by the electronic searches. We approached authors working in the field to locate currently unpublished studies.

Data collection and analysis

We did not use all of the methods stipulated in our published protocol ([Pennington 2017](#)), either because these were not relevant (e.g. we did not encounter cluster-RCTs) or because they could not be deployed (e.g. subgroup analyses). [Table 1](#) summarises the methods section of our protocol, which will be applied in updates of this review if possible. Here, we report only those methods deployed in this review.

Selection of studies

Two review authors (WA and LP) independently screened each title and abstract for eligibility against the inclusion criteria (see [Criteria for considering studies for this review](#)). In the event of a disagreement, resolution was reached in discussion with a third author (JG). When inclusion was uncertain, we obtained the full-text reports. Pairs of review authors (WA, JG, KL or LP) were randomly allocated to each report that appeared from the abstract to fit the inclusion criteria and independently reviewed each one to determine its inclusion. In the event of disagreement regarding inclusion, a third review author (WA, JG, KL or LP) reviewed the report independently and we reached consensus through discussion and by reassessing the inclusion criteria together.

We recorded the selection process in a PRISMA diagram ([Moher 2009](#)).

Data extraction and management

All review authors were involved in data extraction. Two of the four review authors (WA, JG, KL or LP) were randomly assigned to each report and independently extracted data, imputing it into Review Manager 5 ([Review Manager 2014](#)). We developed and piloted a data extraction form for the study to collect data on the following.

1. Country of origin.

2. Type of study: RCT; cluster-RCT; quasi-RCT.
3. Sample size: treatment and control groups; attrition.
4. Study population: parents (age, gender, relationship to child, educational level (high school, further education, higher education)); children (diagnosis of underlying disorder, type of motor disorder (spastic, dyskinetic, ataxic, hypotonic, mixed); age; gender; non-verbal cognitive development (standard scores, percentile rank); receptive language development (standard scores, percentile rank); modes of communication used (vocalisation, speech, gesture, facial expression, body movement, AAC); communicative functions used; number of intelligible words; gross motor function, classified using the Gross Motor Function Classification System (GMFCS; [Palisano 2007](#)); and upper limb function, categorised using the Manual Ability Classification System (MACS; [Eliasson 2006](#)), when possible.
5. Intervention: type of intervention; duration; frequency of sessions; group or individual; content of sessions; inclusion of coaching or didactic teaching only.
6. Comparator intervention: type of intervention; duration; frequency of sessions; group or individual; content of sessions; inclusion of coaching or didactic teaching only.
7. Intervention provider: speech and language therapist (or relevant term in country of origin); psychologist; teacher; other.
8. Fidelity of intervention: how this was assessed and by whom.
9. Outcome measures: parent outcomes; child outcomes; family outcomes.
10. Results: short term (zero to one month following intervention completion), medium term (two to five months after intervention) and long term (six or more months following intervention).
11. Adverse events.
12. Conflicts of interest, including declarations of conflicts of interest.

We resolved disagreements by discussion and by involving a third review author (WA, JG, KL or LP), when necessary.

Assessment of risk of bias in included studies

We rated the risk of bias of each included study using the Cochrane 'Risk of bias' tool ([Higgins 2011a](#)). Pairs of review authors (WA, JG, KL or LP) were randomly allocated to each study to extract data and rate risk of bias. Disagreements were resolved through discussion or by involving a third review author (WA, JG, KL or LP). We rated studies as having low, high or unclear risk of bias in: random sequence generation; allocation concealment; blinding of participants and personnel; blinding of outcome assessment; completeness of data collection; selective reporting; and other sources of bias. We applied the coding schedule in [Appendix 2](#) for each source of bias.

Measures of treatment effect

Continuous data

The included studies reported only continuous data. These were analysed on the assumption that they came from a normally distributed population with no skew. The included studies used different types of measures to evaluate constructs and so we could not combine the data in a meta-analysis. See [Table 1](#) for methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions.

Unit of analysis issues

There were no unit-of-analysis issues as both included studies randomised individual children to one of two study arms. See [Table 1](#) for methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions.

Dealing with missing data

Where necessary, we requested data on participants from study authors, to confirm that at least one child in each group had a motor disorder. If information was not available, we classed the study as 'awaiting classification' and did not include it in this version of the review (e.g. [Fey 2006](#)).

Both included studies reported data on the number of families who dropped out of the study as well as the reasons for dropping out. We reported the number of participants included in the final analysis as a proportion of those participants who began the intervention (see [Characteristics of included studies](#) table). We concluded that the data were missing at random and analysed the remaining data, ignoring the missing data.

We requested missing outcome data from study authors, such pre- and postintervention group mean scores and standard deviations. If data were not available, we included the study in the review and reported, assessed and discussed the extent to which its findings affected the results of the review.

Assessment of heterogeneity

We were unable to assess heterogeneity due to a lack of outcome data available for one of the studies ([Girolametto 1988](#)). See [Table 1](#) for methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions.

Assessment of reporting biases

We were unable to assess the estimate of reporting biases due to the small number of studies included in this review. See [Table 1](#) for methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions.

Data synthesis

Data were unavailable for one study, [Girolametto 1988](#), so we could not conduct a meta-analysis for any of the study outcomes. See [Table 1](#) for methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions.

'Summary of findings' tables

We randomly assigned two review authors (WA, JG, KL or LP) to each outcome. Using the GRADE approach ([GRADEpro 2008](#)), both review authors independently assessed the overall quality of the body of evidence for each outcome as high, moderate, low or very low quality, according to the presence of five criteria (risk of bias, imprecision, inconsistency, indirectness and publication bias). We resolved disagreements by involvement of a third review author (WA, JG, KL or LP). We reported these ratings, along with the number of studies and participants as well as the effect estimate, in [Summary of findings for the main comparison](#), which we constructed using GRADEprofiler ([GRADEpro GDT 2015](#)), having imported data from [Review Manager 2014](#). The table reported the evidence for the following outcomes assessed at postintervention for the comparison 'Parent-mediated communication intervention compared with no intervention': children's ability to communicate effectively in everyday life (taking turns in conversation, initiating conversation and responding to others' conversational gambits); adverse events; child speech and language function; child generic participation: social; parents' communication and interaction strategies: directiveness and responsiveness; family stress and coping; parent stress; satisfaction of child and family with treatment; and compliance with treatment.

Subgroup analysis and investigation of heterogeneity

As we were unable to perform a meta-analysis in this review, we could not undertake subgroup analysis. See [Table 1](#) for details of planned subgroup analyses.

Sensitivity analysis

We were unable to conduct any sensitivity analysis for this version of the review. [Table 1](#) sets out our plans for sensitivity analysis, which we will implement in future updates of this review, if possible.

RESULTS

Description of studies

Results of the search

Our searches identified 8544 records, including six systematic reviews, whose references were handsearched ([Baker 2012](#); [Chorna 2017](#); [Ketelaar 1998](#); [Law 2003](#); [Roberts 2014](#); [Whittingham 2011](#)). After removal of duplicates, two review authors (WA and LP) independently screened the titles and abstracts of 6009 records and subsequently excluded 5943 records on the basis of title and abstract. We retrieved the full-text reports for 63 of the remaining 66 records. We were unable to obtain the full-text reports of three records and classified these as 'awaiting classification' until we

can obtain more information ([Characteristics of studies awaiting classification](#) table). We identified two studies from the 63 full-text reports that met our inclusion criteria ([Included studies](#)), and found two ongoing studies ([Characteristics of ongoing studies](#) table). Fifty-nine of the full-text papers were ineligible for the reasons given in [Figure 1](#), which illustrates the flow of studies through the screening process. From these, we selected 19 excluded papers that at first appeared to meet the eligibility criteria, but on further inspection failed to meet one or more criteria (see [Excluded studies](#)).

Figure 1. Study flow diagram.

Included studies

This review included two RCTs, reported in one paper each (Girolametto 1988; Kim 2005). See [Summary of findings for the main comparison](#).

Participants

One study had 18 child/mother participants, with 10 participants assigned to the intervention group and eight participants to the control group (Kim 2005). The other study had 20 child/mother participants where nine participants assigned to the intervention group, and 11 participants to the control group completed the study (Girolametto 1988). The total number of child participants from the two included studies was 38 (20 boys and 18 girls). Diagnoses included cerebral palsy (10 participants), Down's syndrome (11 participants), developmental delay (11 participants), intellectual disability (three participants), chromosomal abnormalities (two participants) and one participant with an unknown aetiology. Children ranged in age from 15 to 96 months (see [Characteristics of included studies](#) table). Both studies reported cognitive and communication developmental levels substantially lower than expected for their chronological ages for all study participants. Girolametto 1988 reported that all children had hearing within normal limits; Kim 2005 provided no information on children's hearing. All participants attended some special education and lived at home with their parents.

All parents who took part in the evaluation of the interventions were mothers, although fathers did attend the training in Girolametto 1988, but their communication behaviours were not measured. Mothers were similar in age and number of years spent in education across the studies. Mean maternal age was 34 years in both studies. Mean maternal years of education was 13 years in both studies. We retained reference to mothers throughout the review, as results pertain to mothers only.

Participants were recruited from single centres providing special education. There was no information on admission criteria for the services. Therefore, generalisability from the sample to the population of children with communication difficulties and non-progressive motor disorders was unknown. Neither study reported a sample size calculation or statistical power to detect a true difference between groups.

Setting

The two studies took place in Canada (Girolametto 1988), and South Korea (Kim 2005). Girolametto 1988 conducted training at an intervention centre. Kim 2005 did not specify where training was delivered. In both studies, therapists visited families at home

to help parents implement the techniques they had learned in the group sessions.

Intervention content

The intervention provided in both studies was based on the theory that early parent-child interaction that incorporates joint attention, reciprocal turn taking and contingent responding by parents to children's communicative gambits promotes children's engagement in interaction and provide an ideal basis for learning the meaning of gestures and language (Wells 1981). The studies examined different training programmes: Girolametto 1988 provided It Takes Two to Talk (Manolson 1985; Pepper 2004); Kim 2005 delivered Relationship Focussed Intervention, adapted from the Family/Child Curriculum (Mahoney 1999a) and the Principle of Program Learning (Skinner 1958). In both programmes, parents attended group sessions in which they learned about early interaction and communication development, and the importance of following their children's lead and reducing their own directiveness. Sessions also covered how to create opportunities for interaction in play and daily activities. In addition to the group sessions in clinic, intervention also included home visits, in which therapists helped parents to apply principles they had learned in the group sessions in interaction with their children through one-to-one coaching. However, in both studies, details on individual session content and modes of delivery were lacking. It would not be possible to replicate the intervention from the information provided in the two reports. Neither was it possible to determine the extent to which training content overlapped between the two interventions.

Control content

Families allocated to control groups received no specified communication intervention during the studies. Children were either on the waiting list for intervention (Girolametto 1988), or attended the same early special education provision as those allocated to the intervention content, with no input related to responsive interaction (Kim 2005). Parent-child interaction was videorecorded at the same time points as families allocated to the intervention groups, but the studies provided no feedback on the interaction. We found no studies that compared parent-mediated intervention with clinician-mediated intervention in which therapists focused directly on children to teach them new communication behaviours.

Duration of intervention

In one study, the intervention lasted for 11 weeks and consisted of eight group sessions and three individual home visits (Girolametto

1988). The other study had a 12-week intervention period, comprising eight, once-weekly group sessions of 1.5 to two hours followed by four weeks of once-weekly home visits, two of which were to provide instruction and the final two to provide feedback (Kim 2005).

Outcome measures

Although both studies measured parents' and children's interaction behaviours, they used different outcome measures, which precluded meta-analysis. Girolametto 1988 coded videorecorded parent-child interaction on a turn-by-turn basis to show the frequency with which parents and children took obligatory turns in conversation and whether these turns maintained a topic or redirected the conversation to a new topic. Kim 2005 rated parents' and children's communicative behaviours using subjective, four-point rating scales developed for previous research. Studies measured parent-child interaction from 10-minute observations of interaction. Girolametto 1988 also measured children's expressive and receptive communication using the standardised Sequenced Inventory of Communication Development (Hedrick 1975). Kim 2005 measured child behaviour using the Child Behavior Rating Scale (Mahoney 1998) and the Maternal Behavior Rating Scale (Mahoney 1999a), and parental stress using the Korean version (Shin 1997) of the Parent Stress Index (Abidin 1995). This study did not include a measure of the child's comprehension. See [Characteristics of included studies](#) for further details of outcome measures.

Girolametto 1988 reported children's communication behaviours, including comprehension, as group means and ranges, and stated whether statistical comparisons were significant at P less than 0.05 or P less than 0.01 in one-tailed tests. The exact test statistics and P values were not reported, neither were confidence intervals (CIs). Change scores were available for individual children in the intervention group only. Contact with the study author confirmed that unpublished data were no longer available for the study (Girolametto 2018 (personal communication) in Girolametto 1988). Therefore, it was not possible to calculate estimates of effect size for any of the outcomes reported in this study. Kim 2005 reported mean scores and corresponding standard deviations for intervention and control groups at pre- and postintervention for all measures. Published data allowed the calculation of mean differences (MD) for postintervention scores with 95% CIs. All participants who completed the intervention and control content were included in the analyses in both included studies (38 child/parent participants). Studies measured only short-term effects (immediately after intervention). We rated the quality of evidence for all primary and secondary outcomes as very low, using the GRADE

approach for assessing evidence quality (GRADEpro 2008). See [Summary of findings for the main comparison](#).

Funding

One study was funded, in part, by a doctoral fellowship from the Social Sciences and Research Council of Canada (Girolametto 1988). The other study provided no information on funding (Kim 2005).

Excluded studies

Nineteen papers reported 13 studies of parent-mediated training; however, we excluded all on at least one criterion (Adamson 2010; Badr 2006; Barlow 2007; Buschmann 2008; Buschmann 2009; Cologon 2017; Fey 2006; Kaiser 2013; Karaaslan 2013; Kyno 2012; Milgrom 2004; Newnham 2009; Ronski 2010; Seifer 1991; Tannock 1992; Warren 2008; Whitmore 2014; Woyanowski 2014; Yoder 2002). Reasons for exclusion included the following.

1. Children with motor disorders were not included in both arms of the study (16 papers).
2. Children had very mild motor disorders that did not affect their ability to sit unsupported, manipulate objects, produce gesture or spoken language. Their motor disorders were judged by study authors not to affect children's communication (three papers).
3. Interventions did not specifically target communication or include communication outcomes (16 papers).
4. Lack of randomisation (nine papers).
5. Participants were all older than five years of age (three papers).

The reasons for exclusion of individual studies are provided in the [Characteristics of excluded studies](#) table.

Studies awaiting classification

We were unable to obtain the full texts of three reports (Campbell 1977; Kogan 1978; Lee 2012). See [Studies awaiting classification](#) table for further details.

Ongoing studies

We identified two ongoing studies (ACTRN12616000653460; Eliasson 2016). See [Characteristics of ongoing studies](#) for further details.

Risk of bias in included studies

See [Figure 2](#) for a summary of risk of bias.

Figure 2. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

Allocation

Neither study clearly described the allocation process with regard to sequence generation or allocation concealment. One study reported stratified randomisation based on aetiology and sex but provided no information on how this was achieved (Girolametto 1988). We judged the risk of bias for allocation as unclear for both studies (Girolametto 1988; Kim 2005).

Blinding

Due to the nature of the intervention, it was unlikely that parents and interventionists could be blinded to the treatment condition

in either study, and therefore we considered both studies at high risk of performance bias (Girolametto 1988; Kim 2005). One study undertook all outcome assessment blind to group allocation, so we judged this study at low risk of detection bias (Kim 2005). One study blinded outcome assessors to group allocation for speech and language function outcomes, but blinding was unclear for the main outcome measures taken from video-recorded parent-child interaction; hence, we judged this study at unclear risk of detection bias (Girolametto 1988).

Incomplete outcome data

One study was at low risk of attrition bias (Girolametto 1988). Kim 2005 was judged at high risk of attrition bias, as 3/13 children dropped out of intervention group (one family left the treatment centre and one missed three intervention sessions). Inclusion of these children could have impacted the estimates of treatment effectiveness.

Selective reporting

Both studies presented results based on all prespecified outcomes for all participants, so we judged them at low risk of reporting bias (Girolametto 1988; Kim 2005).

Other potential sources of bias

There were no other sources of bias.

Effects of interventions

See: [Summary of findings for the main comparison](#) Parent-mediated communication intervention compared with no intervention for improving the communication skills of preschool children with non-progressive motor disorder

Missing data in one study precluded testing of effect sizes (Girolametto 1988). Therefore, below, we present the outcomes for one study with 18 participants only (Kim 2005).

Primary outcomes

Children's ability to communicate effectively in everyday life

One study showed no evidence of an effect of parent-mediated training on children's initiations of conversation (MD -0.25 points, 95% CI -1.01 to 0.51; $P = 0.54$; [Analysis 1.1](#)), or their engagement in interaction measured through joint attention (MD 0.05 points, 95% CI -0.52 to 0.62; $P = 0.87$; [Analysis 1.2](#)), assessed at postintervention using the Child Behavior Rating Scale (CBRS; therapist rating of child engagement with parent in play in seven domains (interest, attention to activity, persistence, initiation, affect, joint attention, co-operation); items scored one to five on Likert scale; lower scores represent lower levels of coded behaviours) (Mahoney 1999b).

The study did not report children's use of communication (e.g. requesting attention, asking questions) or range of modes of expression (e.g. vocalisation).

Adverse events

The study did not report adverse events.

Secondary outcomes

Child outcomes: speech and language function

One study that measured children's expressive and receptive language development found that missing data precluded analysis of the effect of parent-mediated communication intervention on speech and language function (Girolametto 1988).

Child outcomes: children's generic participation

We found no studies that reported children's generic participation.

Parent outcomes: parents' communication and interaction strategies

Following parent-mediated communication intervention, mothers were more responsive to their children's communication (MD 1.02 points, 95% CI 0.34 to 1.70; $P = 0.01$; [Analysis 2.1](#)), as measured by the Mothers Behaviour rating Scale (MBRS; therapist rating of maternal behaviour on 12 items in four domains (responsiveness, affect, achievement orientation and directiveness); items scored one to five on Likert scale; lower scores represent lower levels of coded behaviours) (Mahoney 1999a). The study showed no evidence of an effect of parent-mediated communication intervention on mothers' directiveness in interaction (MD 0.49 points, 95% CI -0.02 to 1.00; $P = 0.07$; [Analysis 2.2](#)), also assessed at postintervention using the MBRS.

Parent outcomes: family stress and coping

The study showed no evidence of an effect of parent-mediated communication intervention on parental stress in terms of parents' views of their child's temperament (MD -6.50 points, 95% CI -11.64 to -1.36; [Analysis 3.1](#)), parent-child relationship (MD 0.25 points, 95% CI -4.70 to 5.20; $P = 0.92$; [Analysis 3.2](#)), or learning expectation (MD -0.58 points, 95% CI -2.87 to -1.71; [Analysis 3.3](#)), as measured at postintervention using the Korean version of the Parent Stress Index (20 items per scale rated on five-point Likert scale to indicate the extent to which items disturbed the parent in the last week; lower scores indicate lower levels of disturbance) (Abidin 1995; Shin 1997).

Parent outcomes: satisfaction of child and family with treatment

We found no studies that measured satisfaction of child and family with treatment

Parent outcomes: compliance with treatment

We found no studies that measured compliance with treatment.

DISCUSSION

Summary of main results

We identified two small studies that met the inclusion criteria for this review ([Criteria for considering studies for this review](#)), but were only able to calculate effects sizes for [Kim 2005](#) because of missing data in [Girolametto 1988](#). The single study providing data for the review suggested that parent-mediated intervention may have positive effects on mothers' interaction style, helping them to increase their responsiveness to their children. However, we found no evidence of concomitant reductions in mothers' directiveness in conversation. Neither did we find changes in children's interaction patterns, such as increases in initiation of communication or engagement in joint attention for interaction, which would allow them to communicate effectively in everyday life.

We were not able to analyse the effects of parent-mediated communication intervention on children's frequency of communication, frequency of spoken language, speech, or receptive or expressive language development, due to missing data in the one study that measured these outcomes ([Girolametto 1988](#)). Neither study measured wider impacts of parent-mediated communication intervention in children's social participation or considered potential adverse events. One study measured parental stress ([Kim 2005](#)), and while there was no effect of parental training, differences between groups prior to therapy prevented firm conclusions being drawn about the impact of intervention on this outcome.

Studies measured communication in short play-based interactions of 10 minutes' duration immediately after training. We found no evidence to suggest that changes in interaction were maintained over time, generalised to daily interaction or stimulated children's longer-term communication development.

The studies used different tools to measure similar constructs. Examination of the full-text reports considered for the review showed that a wide range of tools has been used to evaluate parent-mediated communication interventions. Consensus is needed on the constructs that should be measured and the tools that should be used, with demonstration of their validity and reliability.

Overall completeness and applicability of evidence

Both studies recruited children with developmental disabilities and included some children who were identified as having motor disorders. All children had some degree of intellectual impairment. We found no studies that examined parent-mediated intervention delivered specifically to parents of children with motor disorders, or studies that included children with motor disorders who did not have intellectual impairment. Neither was it possible to disaggregate the results for children with motor disorders in this review. Because of a lack of detail on participants, it was not possible to

ascertain how children in the studies communicated, and the impact of their motor disorders on the intelligibility of their communication signals. Therefore, we could not ascertain if parent-mediated intervention was effective for children with motor disorders whose communication signals were difficult to interpret because of a lack of consistency and precision in underlying movements. The transactional theory of development suggests that communication patterns may change at any point in life as a result of interaction with others ([Sameroff 2000](#)). We focused on parent-mediated interventions delivered in the preschool years (up to five years of age), as at this point in children's lives their most frequent communication partner is usually their parent. Some of the children in the study by [Kim 2005](#) were older than five years of age and results may suggest that parent-mediated communication intervention could be provided to parents of school-aged children. Although some fathers attended training in the study by [Girolametto 1988](#), we were able to estimate the effects of training mothers only. Therefore, results may not be generalisable to fathers or other primary carers.

All children included in the studies were receiving special education services. Consequently, results may not be applicable to families who are not receiving such services.

Parent-mediated interventions were developed in North America and have been applied further afield, in Europe (e.g. [Badr 2006](#) and [Karaaslan 2013](#)) and Australasia (e.g. [Whittingham 2011](#)). The programmes assume that parents view active communication by children, in the form of conversation initiation, as positive and culturally acceptable. The study by [Kim 2005](#) suggested parent-mediated communication intervention based on the transactional theory of communication development by [Sameroff 2000](#) may be effective in other settings, although the high rate of attrition from parent-mediated communication interventions suggests that their acceptability requires further investigation.

Quality of the evidence

We judged the studies included in this review to be of very low quality using [GRADEpro 2008](#). Sample sizes were small and lacked justification. Studies were at risk of allocation bias; neither provided information on how allocation was achieved or maintained during recruitment. Furthermore, in the study by [Kim 2005](#), parents allocated to the two groups appeared to differ prior to therapy, particularly on parental stress measures. Before intervention, mothers in the intervention group had a higher mean level of stress related to their relationship with their child than mothers in the control group. Postintervention, mean scores reduced to levels similar to the control group. Analysing postintervention scores disregards individual change from baseline. Children with motor disorders who receive intervention to promote their communication vary widely in their communication and language development at the start of therapy (e.g. the frequency with which they initiate interaction and the number of words they produce)

and the rate at which they learn new skills. Similarly, their parents may differ in interaction characteristics and measures of well-being. Therefore, it is important to consider differences both between groups and within groups over time when estimating an intervention's effect. Unfortunately, the data presented in the studies in this review did not allow us to do so, which may have inflated some results (e.g. parents' responsiveness) and negated others (e.g. parent stress).

As with all parent-training interventions, it is not possible to blind participants and personnel providing the intervention to group allocation, so risk of performance bias was also high for both studies. Both studies achieved blinding of outcome assessors, although we rated one study at unclear risk of detection bias as the text did not explicitly state that outcome assessors were blind to group allocation for all outcomes (Girolametto 1988). We judged one study at high risk of attrition bias, as the reasons for loss to follow-up could be related to treatment satisfaction for two out of the three children lost in the parent-mediated intervention group (Kim 2005). All participants who completed the study were included in the analysis of all outcome measures, thereby reducing the risk of reporting biases.

Potential biases in the review process

Terminology to describe developmental disabilities has changed since the 1980s. It is possible that our search did not capture all studies of parent-mediated communication interventions that included children with motor disorders. We excluded studies that involved children with developmental delay and intellectual disability where there was no reference to motor disorder, motor skills or diagnoses that included motor disorders. However, some children in these excluded studies may have had difficulties producing precise and consistent movements in communication as communication was not specified.

Agreements and disagreements with other studies or reviews

A previous version of this review covered parent-mediated intervention and speech and language therapy provided directly to children with cerebral palsy aged up to 19 years and included all study designs (Pennington 2003). That review identified four studies of parent-mediated intervention, involving 34 children aged eight months to 17 years, none of which were RCTs or quasi-RCTs. Those studies suggested similar improvements in children's communication to those observed in the present study with children taking more turns in conversation and becoming more engaged in interaction.

AUTHORS' CONCLUSIONS

Implications for practice

Some speech and language therapists currently provide individual or group communication training for parents, although the content and intensity of such training varies across providers (Watson 2015). Conclusive evidence of the effectiveness of parent-mediated communication interventions for children with motor disorders has not been demonstrated by this review.

Implications for research

Children with motor disorders may have difficulties in producing consistent and precise movements for gestural, vocal and verbal communication, which may render their attempts to communicate unintelligible. This review shows that further research is needed to test the effectiveness of parent-mediated communication interventions for children whose motor disorders impairs their expressive communication. Children with motor disorder may or may not have intellectual impairment. It is important to differentiate between these two groups as rates of change may differ. Studies that focus on children with intellectual/developmental disability should include a subgroup analysis of participants who also have motor disorders because of the complexity of their expressive communication.

To enable comparison across studies, and amalgamation of data in future systematic reviews, there is an urgent need for researchers to agree a core set of descriptor and outcome measures. Difficulties in individual developmental domains should be identified and described in sufficient detail to allow readers to ascertain the potential impact of the domain on the intervention outcome.

To test the theory of transactional communication development, outcomes should consider parents' interaction style, showing how they provide opportunities for children to join and control conversation; children's communication, measuring the frequency with which they take turns in conversation, the intentions they express (communicative functions) and the complexity of their communication (e.g. length of turn, number of different words/referents, phrase structure); acceptability of interventions; and impact on parental stress as both a potential benefit but also a potential adverse event.

Outcomes should be measured in the medium term (six months) to capture transactional changes in parents' and children's conversation behaviours, and the longer (e.g. one year) to evaluate if training facilitates ongoing, positive change to children's communication development trajectories.

Studies should recruit children who receive a variety of different services via a range of different sources to ensure generalisation of findings.

The studies included in this review involved small samples, limiting their ability to detect true differences and generalisability. Future research must provide sample size calculations to show that

it is capable of detecting differences between groups. Research reporting must also be improved, particularly in the area of participant identification, recruitment, and selection and allocation to demonstrate risk of selection bias. Application of [CONSORT 2010](#) may help in this regard. This review was hampered by poor descriptions of the interventions provided. Future research must ensure adequate descriptions of interventions through adoption of the template for intervention description and replication (TIDieR) guidelines ([Hoffmann 2014](#)), to allow testing of adherence and development of a core set of active ingredients.

As stated above ([Overall completeness and applicability of evidence](#)), parent-mediated interventions have been adopted in cultures that expect children to be equal, independent, communication partners in conversations with parents. Future research should explore their acceptability and effectiveness in other cultures.

The group of communication partners who receive training should also be considered. The studies included in this review are not un-

usual in involving only mothers in their evaluations of treatment effect, as they are often children's primary carers. However, fathers, other family members and paid carers may also spend considerable time with children and influence their development. Future research should also consider the effects of training on their interaction and influence.

Finally, the economic impact of communication training provision and its outcomes should be evaluated in any future trial.

ACKNOWLEDGEMENTS

We thank Dr Joanne Duffield, Professor Geraldine Macdonald, and Dr Adrienne Harvey of Cochrane Developmental, Psychosocial and Learning Problems (CDPLP), for feedback; Margaret Anderson, CDPLP, for her help in developing the search strategies; and the anonymous reviewers for their helpful feedback on previous drafts of the review.

REFERENCES

References to studies included in this review

Girolametto 1988 *{published data only}*

Girolametto L. Participant characteristics and trial results [personal communication]. Email to: L Pennington 8 January 2018.

* Girolametto LE. Improving the social-conversational skills of developmentally delayed children: an intervention study. *Journal of Speech and Hearing Disorders* 1988;**53**(2):156–67. DOI: 10.1044/jshd.5302.156; PUBMED: 2452300

Kim 2005 *{published data only}*

Kim JM, Mahoney G. The effects of relationship focused intervention on Korean parents and their young children with disabilities. *Research in Developmental Disabilities* 2005;**26**(2):117–30. DOI: 10.1016/j.ridd.2004.08.001; PUBMED: 15590243

References to studies excluded from this review

Adamson 2010 *{published data only (unpublished sought but not used)}*

Adamson LB, Ronski MA, Bakeman R, Sevcik RA. Augmented language intervention and the emergence of symbol-infused joint engagement. *Journal of Speech, Language, and Hearing Research* 2010;**53**(6):1769–73. DOI: 10.1044/1092-4388(2010/09-0208); PMC4428337; PUBMED: 20705741

Badr 2006 *{published data only}*

Badr LK, Garg M, Kamath M. Intervention for infants with brain injury: results of a randomized controlled study. *Infant Behavior & Development* 2006;**29**(1):80–90. DOI: 10.1016/j.infbeh.2005.08.003; 17138264; PMC2700252

Barlow 2007 *{published data only}*

Barlow J, Powell L, Cheshire A. The Training and Support Programme (involving basic massage) for parents of children with cerebral palsy: an implementation study. *Journal of Bodywork and Movement Therapies* 2007;**11**(1):44–53. DOI: 10.1016/j.jbmt.2006.03.002

Buschmann 2008 *{published data only}*

Buschmann A, Jooss B, Rupp A, Dockter S, Blaschikowitz H, Heggen I, et al. Children with developmental language delay at 24 months of age: results of a diagnostic work-up. *Developmental Medicine & Child Neurology* 2008;**50**(3):223–9. DOI: 10.1111/j.1469-8749.2008.02034.x; PUBMED: 18266869

Buschmann 2009 *{published data only}*

Buschmann A, Jooss B, Rupp A, Feldhusen F, Pietz J, Philippi H. Parent based language intervention for 2-year-old children with specific expressive language delay: a randomised controlled trial. *Archives of Disease in Childhood* 2009;**94**(2):110–6. DOI: 10.1136/adc.2008.141572; PMC2614563; PUBMED: 18703544

Cologon 2017 *{published data only}*

Cologon K, Wicks L, Salvador A. Supporting caregivers in developing responsive communication partnerships with their children: extending a caregiver-led interactive language program. *Child Language Teaching and Therapy* 2017;**33**(2):157–69. DOI: 10.1177/0265659016650978

Fey 2006 *{published data only}*

Fey ME, Warren SF, Brady N, Finestack LH, Bredin-Oja SL, Fairchild M, et al. Early effects of responsivity education/prelinguistic milieu teaching for children with developmental delays and their parents. *Journal of Speech,*

- Language, and Hearing Research* 2006;**49**(3):526–47. DOI: 10.1044/1092-4388(2006/039); PUBMED: 16787894
- Kaiser 2013** *{published data only}*
Kaiser AP, Roberts MY. Parent-implemented enhanced milieu teaching with preschool children who have intellectual disabilities. *Journal of Speech, Language, and Hearing Research* 2013;**56**(1):295–309. DOI: 1092-4388 (2012/11-0231); PMC3740334; PUBMED: 22744141
- Karaaslan 2013** *{published data only}*
Karaaslan O, Diken IH, Mahoney GH. A randomized control study of responsive teaching with young Turkish children and their mothers. *Topics in Early Childhood Special Education* 2013;**33**(1):18–27. DOI: 10.1177/0271121411429749
- Kyno 2012** *{published data only}*
Kyno NM, Ravn IH, Lindemann R, Fagerland MW, Smeby NA, Torgersen AM. Effect of an early intervention programme on development of moderate and late preterm infants at 36 months: a randomized controlled study. *Infant Behavior & Development* 2012;**35**(4):916–26. DOI: 10.1016/j.infbeh.2012.09.004; PUBMED: 23063851
- Milgrom 2004** *{published data only}*
Milgrom J, Westley DT, Gemmill AW. The mediating role of maternal responsiveness in some longer term effects of postnatal depression on infant development. *Infant Behavior & Development* 2004;**27**(4):443–54. DOI: 10.1016/j.infbeh.2004.03.003
- Newnham 2009** *{published data only}*
Newnham CA, Milgrom J, Skouteris H. Effectiveness of a modified mother-infant transaction program on outcomes for preterm infants from 3 to 24 months of age. *Infant Behavior & Development* 2009;**32**(1):17–26. DOI: 10.1016/j.infbeh.2008.09.004; PUBMED: 19026450
- Romski 2010** *{published data only (unpublished sought but not used)}*
* Romski M, Sevcik RA, Adamson LB, Cheslock M, Smith A, Barker RM, et al. Randomized comparison of augmented and non-augmented language interventions for toddlers with developmental delays and their parents. *Journal of Speech, Language, and Hearing Research* 2010;**53**(2):350–64. DOI: 10.1044/1092-4388(2009/08-0156); PUBMED: 20360461
- Seifer 1991** *{published data only}*
Seifer R, Clark GN, Sameroff AJ. Positive effects of interaction coaching on infants with developmental disabilities and their mothers. *American Journal of Mental Retardation* 1991;**96**(1):1–11.
- Tannock 1992** *{published data only}*
Tannock R, Girolametto L, Siegel LS. Language intervention with children who have developmental delays: effects of an interactive approach. *American Journal on Mental Retardation* 1992;**97**(2):145–60. SSN–0895–8017; PUBMED: 1384566]
- Warren 2008** *{published data only}*
Warren SF, Fey ME, Finestack LH, Brady NC, Bredin-Oja SL, Fleming KK. A randomized trial of longitudinal effects of low-intensity responsivity education/prelinguistic milieu teaching. *Journal of Speech, Language, and Hearing Research* 2008;**51**(2):451–70. DOI: 10.1044/1092-4388(2008/033); PUBMED: 18367689
- Whitmore 2014** *{published data only}*
Whitmore AS, Romski MA, Sevcik RA. Early augmented language intervention for children with developmental delays: potential secondary motor outcomes. *Augmentative and Alternative Communication* 2014;**30**(3):200–12. DOI: 10.3109/07434618.2014.940466; PUBMED: 25109299
- Woynaroski 2014** *{published data only}*
Woynaroski T, Yoder PJ, Fey ME, Warren SF. A transactional model of spoken vocabulary variation in toddlers with intellectual disabilities. *Journal of Speech, Language, and Hearing Research* 2014;**57**(5):1754–63. DOI: 10.1044/2014.JSLHR-L-13-0252; PMC4192117; PUBMED: 24802090
- Yoder 2002** *{published data only}*
Yoder PJ, Warren SF. Effects of prelinguistic milieu teaching and parent responsivity education on dyads involving children with intellectual disabilities. *Journal of Speech, Language, and Hearing Research* 2002;**45**(6):1158–74. DOI: 10.1044/1092-4388(2002/094); PUBMED: 12546485

References to studies awaiting assessment

- Campbell 1977** *{published data only}*
Campbell AA. *A Comparison of the Effect of Two Types of Treatment Involving Professionals and Parents on the Developmental Progress of Preschool Developmentally Delayed Children [Thesis]*. Memphis (TN): Memphis State University, 1977.
- Kogan 1978** *{published data only}*
Kogan KL, Tyler NB. *Comparing Ways of Altering Parent-Child Interaction*. Washington (DC): Washington University, 1978.
- Lee 2012** *{published data only}*
Lee H-Y, Chen C-L, Cheng H-YK, Hwang A-W, Chen L-J, Liu S-W, et al. Effects of home program intervention for preschool children with developmental delay: a preliminary study. *Journal of Rehabilitation Medicine* 2012;**44**(Suppl 52):103. 0522PP049; htresearch.cgu.edu.tw/ezfiles/14/1014/img/651/101-55-B.pdf]

References to ongoing studies

- ACTRN12616000653460** *{published data only}*
ACTRN12616000653460. LEAP-CP: Learning through Everyday Activities with Parents for infants at high risk of Cerebral Palsy in a low-income country [Community-based parent delivered early detection and intervention program for infants at high risk of cerebral palsy in a low-resource setting: a randomised controlled trial]. www.anzctr.org.au/Trial/Registration/TrialReview.aspx?id=370512 (date first received 19 May 2016).
- Eliasson 2016** *{published data only}*
Eliasson AC, Holmström L, Aarne P, Nakeva von Mentzer C, Weiland AL, Sjöstrand L, et al. Efficacy of the small

step program in a randomised controlled trial for infants below age 12 months with clinical signs of CP: a study protocol. *BMC Pediatrics* 2016;**16**(1):175. DOI: 10.1186/s12887-016-0711-x; NCT02166801 ; PMC5093986; PUBMED: 27809886

Additional references

Abidin 1995

Abidin RR. *Parenting Stress Index*. 3rd Edition. Odessa (FL): Psychological Assessment Resources, 1995.

ACPR 2016

Australian Cerebral Palsy Register Group (ACPR). *Report of the Australian Cerebral Palsy Register 2016. Birth Years 1993-2009*. Sydney (AU): Cerebral Palsy Alliance; CP Register, 2016.

APA 2000

American Psychological Association. *Diagnostic and statistical manual of mental disorders*. 4th Edition. Washington (DC): American Psychological Association, 2000.

Baker 2012

Baker T, Haines S, Yost J, DiClaudio S, Braun C, Holt S. The role of family-centered therapy when used with physical or occupational therapy in children with congenital or acquired disorders. *Physical Therapy Reviews* 2012;**17**(1): 29–36. DOI: 10.1179/1743288X11Y.0000000049

Bax 2006

Bax M, Tydeman C, Flodmark O. Clinical and MRI correlates of cerebral palsy: the European Cerebral Palsy Study. *Journal of the American Medical Association* 2006;**296**(13):1602–8. DOI: 10.1001/jama.296.13.1602; PUBMED: 17018805

Blackburn 2012

Blackburn JS, Mink JW, Augustine EF. Pediatric movement disorders: five new things. *Neurology. Clinical Practice* 2012;**2**(4):311–8. DOI: 10.1212/CPJ.0b013e318278bf06; PMC3613211; PUBMED: 23634375

Boyle 2011

Boyle CA, Boulet S, Schieve LA, Cohen RA, Blumberg SJ, Yeargin-Allsopp M, et al. Trends in the prevalence of developmental disabilities in US children, 1997-2008. *Pediatrics* 2011;**127**(6):1034–42. DOI: 10.1542/peds.2010-2989; PUBMED: 21606152

Broberg 2012

Broberg M, Ferm U, Thunberg G. Measuring responsive style in parents who use AAC with their children: development and evaluation of a new instrument. *Augmentative and Alternative Communication* 2012;**28**(4):243–53. DOI: 10.3109/07434618.2012.740686; PUBMED: 23256856

Bruno 1998

Bruno J, Dribbon M. Outcomes in AAC: evaluating the effectiveness of a parent training program. *Augmentative and Alternative Communication* 1998;**14**(2):59–70. DOI: 10.1080/07434619812331278216

Bzoch 1970

Bzoch KR, League R. *Receptive Expressive Emergent Language Scale*. Gainesville (FL): Tree of Life Press, 1970.

Cans 2008

Cans C, De-la-Cruz J, Mermet M. Epidemiology of cerebral palsy. *Paediatrics and Child Health* 2008;**18**(9):393–8. DOI: 10.1016/j.paed.2008.05.015

Chorna 2017

Chorna O, Hamm E, Cummings C, Fethers A, Maitre NL. Speech and language interventions for infants aged 0 to 2 years at high risk for cerebral palsy: a systematic review. *Developmental Medicine & Child Neurology* 2017;**59**(4): 355–60. DOI: 10.1111/dmcn.13342; PMC5395422; PUBMED: 27897320

Clarke 2017

Clarke MT, Soto G, Nelson K. Language learning, recasts, and interaction involving AAC: background and potential for intervention. *Augmentative and Alternative Communication* 2017;**33**(1):42–50. DOI: 10.1080/07434618.2016.1278130; PUBMED: 28166652

CONSORT 2010

Schulz KF, Altman DG, Moher D, CONSORT Group. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMJ* 2010; Vol. 340:c332. DOI: 10.1136/bmj.c332; PMC2844940; PUBMED: 20332509

De Kloet 2013

De Kloet AJ, Hilberink SR, Roebroek ME, Catsman-Berervoets CE, Peeters E, Lambregts SA, et al. Youth with acquired brain injury in the Netherlands: a multi-centre study. *Brain Injury* 2013;**27**(7-8):843–9. DOI: 10.3109/02699052.2013.775496; PUBMED: 23758314

Dickinson 2007

Dickinson HO, Parkinson KN, Ravens-Sieberger U, Schirripa G, Thyen U, Arnaud C, et al. Self-reported quality of life of 8-12-year-old children with cerebral palsy: a cross-sectional European study. *Lancet* 2007;**369**(9580):2171–8. DOI: 10.1016/S0140-6736(07)61013-7; PUBMED: 17604799

Dunst 1985

Dunst CJ. Communicative competence and deficits: effects of early social interactions. In: McDonald ET, Gallagher DL editor(s). *Facilitating Social-Emotional Development in Multiply Handicapped Children*. Philadelphia (PA): Home of the Merciful Savior for Crippled Children, 1985:93–140.

Egger 1997

Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**(7109):629–34. DOI: 10.1136/bmj.315.7109.629

Eliasson 2006

Eliasson AC, Krumlinde-Sundholm L, Rösblad B, Beckung E, Arner M, Öhrvall AM, et al. The Manual Ability Classification System (MACS) for children with cerebral palsy: scale development and evidence of validity and reliability. *Developmental Medicine & Child Neurology*

- 2006;**48**(7):549–54. DOI: 10.1017/S0012162206001162; PUBMED: 16780622
- Enderby 2015**
Enderby P, John A. *Therapy Outcome Measures for Rehabilitation Professionals*. Guildford (UK): J&R Press, 2015.
- Fauconnier 2009**
Fauconnier J, Dickinson HO, Beckung E, Marcelli M, McManus V, Michelsen SI, et al. Participation in life situations of 8-12 year old children with cerebral palsy: cross sectional European study. *BMJ* 2009;**338**:b1458–71. DOI: 10.1136/bmj.b1458; PMC2673343; PUBMED: 19395424
- Fenson 2006**
Fenson L, Marchman VA, Thal DJ, Dale PS, Reznick JS, Bates E. *The MacArthur-Bates Communicative Development Inventories User's Guide and Technical Manual*. 2nd Edition. Baltimore (MD): Brookes, 2006.
- Folio 2000**
Folio MR, Fewell RR. *Peabody Developmental Motor Scales, PDMS-2*. 2nd Edition. Austin (TX): Pro ED, 2000.
- Friedrich 1983**
Friedrich WN, Greenberg MT, Crnic K. A short-form of the Questionnaire on Resources and Stress. *American Journal of Mental Deficiency* 1983;**88**(1):41–8. [PUBMED: 6225338]
- Gibbard 2004**
Gibbard D, Cogan L, MacDonald J. Cost-effectiveness analysis of current practice and parent intervention for children under 3 years presenting with expressive language delay. *International Journal of Language & Communication Disorders* 2004;**39**(2):229–44. DOI: 10.1080/13682820310001618839; PUBMED: 15204453
- Girolametto 1986**
Girolametto L, Greenberg J, Manolson HA. Developing dialogue skills: the Hanen early language parent program. *Seminars in Speech and Language* 1986;**7**:367–82.
- Girolametto 1996**
Girolametto L, Pearce PS, Weitzman E. Interactive focused stimulation for toddlers with expressive vocabulary delays. *Journal of Speech and Hearing Research* 1996;**39**(6):1274–83. DOI: 10.1044/jshr.3906.1274; PUBMED: 8959612
- GRADEpro 2008 [Computer program]**
Brozek J, Oxman A, Schünemann H. GRADEpro 3.2 for Windows. Hamilton (ON): GRADE Working Group, McMaster University, 2008.
- GRADEpro GDT 2015 [Computer program]**
GRADE Working Group, McMaster University. GRADEpro GDT: GRADEpro Guideline Development Tool. Version accessed prior to 28 November 2016. Hamilton (ON): GRADE Working Group, McMaster University, 2015.
- Griffiths 1970**
Griffiths R. *The Abilities of Young Children: A Comprehensive System of Mental Measurement for the First Eight Years of Life*. London (UK): Child Development Research Centre, 1970.
- Hanzlik 1990**
Hanzlik JR. Nonverbal interaction patterns of mothers and their infants with cerebral palsy. *Education and Training in Mental Retardation* 1990;**25**(4):333–43.
- Hedrick 1975**
Hedrick DL, Prather EM. *Sequenced Inventory of Communication Development*. Torrance (CA): WPS, 1975.
- Hemmeter 1994**
Hemmeter ML, Kaiser AP. Enhanced milieu teaching: effects of parent-implemented language intervention. *Journal of Early Intervention* 1994;**18**(3):269–89. DOI: 10.1177/105381519401800303
- Higgins 2002**
Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* 2002;**21**(11):1539–58. DOI: 10.1002/sim.1186; PUBMED: 12111919
- Higgins 2011a**
Higgins JP, Altman DG, Sterne JA. Chapter 8: Assessing risk of bias in included studies. In: Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.
- Higgins 2011b**
Higgins JP, Deeks JJ, Altman DD. Chapter 16: Special topics in statistics. In Higgins JP, Green S, editor (s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.
- Hoffmann 2014**
Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;**348**:g1687. DOI: 10.1136/bmj.g1687; PUBMED: 24609605
- Johnson-Martin 2004**
Johnson-Martin MM, Attermeier SM, Hacker BJ. *The Carolina Curriculum for Infants and Toddlers with Special Needs*. Baltimore (MD): Paul H Brookes, 2004.
- Kaiser 1987**
Kaiser A, Alpert CL, Hemmeter ML, Ostrosky MM. *Milieu Teaching Manual*. Nashville (TN): Vanderbilt University, 1987.
- Kaiser 1995**
Kaiser AP, Hemmeter ML, Ostrosky MM, Alpert CL, Hancock TB. The effects of group training and individual feedback on parent use of milieu teaching. *Journal of Childhood Communication Disorders* 1995;**16**(2):39–48. DOI: 10.1177/152574019501600206
- Kaiser 2001**
Kaiser AP, Hester PP, McDuffie AS. Supporting communication in young children with developmental disabilities. *Mental Retardation and Developmental Disabilities Research Reviews* 2001;**7**(2):143–50. DOI: 10.1002/mrdd.1020

Kaiser 2003

Kaiser AP, Hancock TB. Teaching parents new skills to support their young children's development. *Infants and Young Children* 2003;**16**(1):9–21.

Kent-Walsh 2015

Kent-Walsh J, Murza KA, Malani MD, Binger C. Effects of communication partner instruction on the communication of individuals using AAC: a meta-analysis. *Augmentative and Alternative Communication* 2015;**31**(4):271–84. DOI: 10.3109/07434618.2015.1052153; PUBMED: 26059542

Ketelaar 1998

Ketelaar M, Vermeer A, Helders PJM, Hart HT. Parental participation in intervention programs for children with cerebral palsy: a review of research. *Topics in Early Childhood Special Education* 1998;**18**(2):107–17. DOI: 10.1177/027112149801800206

King 2004

King G, Law L, King S, Hurley P, Rosenbaum P, Hanna S, et al. *Children's Assessment of Participation and Enjoyment (CAPE) and Preferences for Activities of Children (PAC)*. San Antonio (TX): Pearson, 2004.

Kirby 2011

Kirby RS, Wingate MS, Van Naarden Braun K, Doernberg NS, Arneson CL, Benedict RE, et al. Prevalence and functioning of children with cerebral palsy in four areas of the United States in 2006: a report from the Autism and Developmental Disabilities Monitoring Network. *Research in Developmental Disabilities* 2011;**32**(2):462–9. DOI: 10.1016/j.ridd.2010.12.042; PUBMED: 21273041

Law 2003

Law J, Garrett Z, Nye C. Speech and language therapy interventions for children with primary speech and language delay or disorder. *Cochrane Database of Systematic Reviews* 2003, Issue 3. DOI: 10.1002/14651858.CD004110

Light 1985

Light J, Collier B, Parnes P. Communicative interaction between young nonspeaking physically disabled children and their primary caregivers: part II - communicative function. *AAC: Augmentative and Alternative Communication* 1985;**1**(3):98–107. DOI: 10.1080/07434618512331273591

Mahoney 1988

Mahoney G, Powell A. Modifying parent-child interaction: enhancing the development of handicapped children. *Journal of Special Education* 1988;**22**(1):82–6. DOI: 10.1177/002246698802200110

Mahoney 1998

Mahony G, Bella JM. Examination is centred early intervention on child and family outcomes. *Topics in Early Childhood Special Education* 1998;**18**(2):83–94. DOI: 10.1177/027112149801800204

Mahoney 1999a

Mahoney G. *Family/Child Curriculum: a Relationship Focused Approach to Parent Education/Early Intervention*. Tallmadge (OH): Family Child Learning Center, 1999.

Mahoney 1999b

Mahoney G, Wheeden CA. The effect of the teacher style on interactive engagement of preschool aged children with special learning needs. *Early Childhood Research Quarterly* 1999;**14**(1):51–68.

Manolson 1985

Manolson A. *It Takes Two to Talk: the Hanen Program*. Toronto (CA): Hanen Early Language Resource Centre, 1985.

Marshall 1994

Marshall GN, Hays RD. *The Patient Satisfaction Questionnaire Short Form (PSQ-18)*. Santa Monica (CA): RAND Corporation, 1994.

McDuffie 2016

McDuffie A, Oakes A, Machalicek W, Ma M, Bullard L, Nelson S, et al. Early language intervention using distance video-teleconferencing: a pilot study of young boys with Fragile X Syndrome and their mothers. *American Journal of Speech-Language Pathology* 2016;**25**:46–66. DOI: 10.1044/2015_AJSLP-14-0137

Moher 2009

Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine* 2009;**6**(7):e1000097. DOI: 10.1371/journal.pmed.1000097; PMC2707599; PUBMED: 19621072

Morris 2015

Morris C, Simkiss D, Busk M, Morris M, Allard A, Denness J, et al. Setting research priorities to improve the health of children and young people with neurodisability: a British Academy of Childhood Disability-James Lind Alliance Research Priority Setting Partnership. *BMJ Open* 2015;**5**(1):e006233. DOI: 10.1136/bmjopen-2014-006233

Nordberg 2013

Nordberg A, Miniscalco C, Lohmander A, Himmelmann K. Speech problems affect more than one in two children with cerebral palsy: Swedish population-based study. *Acta Paediatrica* 2013;**102**(2):161–6. DOI: 10.1111/apa.12076; PUBMED: 23186066

Noreau 2007

Noreau L, Lepage C, Boissiere L, Picard R, Fougereyrollas P, Mathieu J, et al. Measuring participation in children with disabilities using the Assessment of Life Habits. *Developmental Medicine & Child Neurology* 2007;**49**(9):666–71. DOI: 10.1111/j.1469-8749.2007.00666.x; PUBMED: 17718822

Novak 2012

Novak I, Hines M, Goldsmith S, Barclay R. Clinical prognostic messages from a systematic review on cerebral palsy. *Pediatrics* 2012;**130**(5):e1285–312. DOI: 10.1542/peds.2012-0924; PUBMED: 23045562

O'Toole 2016

O'Toole C, Lee ASY, Gibbon FE, van Bysterveldt AK, Conway P, Hart NJ. Parent-mediated interventions to promote communication and language development in

- children with Down syndrome aged between birth and six years. *Cochrane Database of Systematic Reviews* 2016, Issue 2. DOI: 10.1002/14651858.CD012089
- Oono 2013**
Oono IP, Honey EJ, McConachie H. Parent-mediated early intervention for young children with autism spectrum disorders (ASD). *Cochrane Database of Systematic Reviews* 2013, Issue 4. DOI: 10.1002/14651858.CD009774.pub2
- Palisano 2007**
Palisano R, Rosenbaum P, Bartlett D, Livingston M. *Gross Motor Function Classification System Expanded and Revised*. Toronto (ON): CanChild Centre for Childhood Disability Research, McMaster University, 2007.
- Parker 2010**
Parker SE, Mai CT, Canfield MA, Rickard R, Wang Y, Meyer RE, et al. Updated National Birth Prevalence estimates for selected birth defects in the United States, 2004-2006. *Birth Defects Research. Part A, Clinical and Molecular Teratology* 2010;**88**(12):1008–16. DOI: 10.1002/bdra.20735; PUBMED: 20878909
- Parkes 2010**
Parkes J, Hill N, Platt MJ, Donnelly C. Oromotor dysfunction and communication impairments in children with cerebral palsy: a register study. *Developmental Medicine & Child Neurology* 2010;**52**(12):1113–9. DOI: 10.1111/j.1469-8749.2010.03765.x; PUBMED: 20813020
- Parkes 2011**
Parkes J, Caravale B, Marcelli M, Franco F, Colver A. Parenting stress and children with cerebral palsy: a European cross-sectional survey. *Developmental Medicine & Child Neurology* 2011;**53**(9):815–21. DOI: 10.1111/j.1469-8749.2011.04014.x; PUBMED: 21707599
- Pennington 2001**
Pennington L, McConachie H. Interaction between children with cerebral palsy and their mothers: the effects of speech intelligibility. *Journal of Language and Communication Disorders* 2001;**36**(3):371–93. [PUBMED: 11491485]
- Pennington 2003**
Pennington L, Goldbart J, Marshall J. Speech and language therapy to improve the communication skills of children with cerebral palsy. *Cochrane Database of Systematic Reviews* 2003, Issue 3. DOI: 10.1002/14651858.CD003466.pub2
- Pennington 2009**
Pennington L, Thomson K, James P, Martin L, McNally R. Effects of It Takes Two to Talk - the Hanen Program for parents of preschool children with cerebral palsy: findings from an exploratory study. *Journal of Speech, Language, and Hearing Research* 2009;**52**(5):1121–38. DOI: 10.1044/1092-4388(2009/07-0187); PUBMED: 19635943
- Pepper 2004**
Pepper J, Weitzman E, McDade A. *Making Hanen happen: It Takes Two to Talk - the Hanen Program for Parents. Leader's Guide for Certified Speech-Language Pathologists*. Toronto (ON): Hanen Centre, 2004.
- Pickstone 2009**
Pickstone C, Goldbart J, Marshall J, Rees A, Roulstone S. A systematic review of environmental interventions to improve child language outcomes for children with or at risk of primary language impairment. *Journal of Research in Special Educational Needs* 2009;**9**(2):66–79. DOI: 10.1111/j.1471-3802.2009.01119.x
- Pirila 2007**
Pirila S, van der Meere J, Pentikainen T, Ruusu-Niemi P, Korpela R, Kilpinen J, et al. Language and motor speech skills in children with cerebral palsy. *Journal of Communication Disorders* 2007;**40**(2):116–28. DOI: 10.1016/j.jcomdis.2006.06.002; PUBMED: 16860820
- Pousada 2013**
Pousada M, Guillamón N, Hernández-Encuentra E, Muñoz E, Redolar D, Boixadós M, et al. Impact of caring for a child with cerebral palsy on the quality of life of parents: a systematic review of the literature. *Journal of Developmental and Physical Disabilities* 2013;**25**(5):545–77. DOI: 10.1007/s10882-013-9332-6
- Reid 2011**
Reid SM, Carlin JB, Reddihough DS. Rates of cerebral palsy in Victoria, Australia, 1970 to 2004: has there been a change?. *Developmental Medicine & Child Neurology* 2011;**53**(10):907–12. DOI: 10.1111/j.1469-8749.2011.04039.x; PUBMED: 21752018
- Review Manager 2014 [Computer program]**
Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager 5 (RevMan 5). Version 5.3. Copenhagen: Nordic Cochrane Centre, The Cochrane Collaboration, 2014.
- Roberts 2014**
Roberts MY. Parenting interventions for children with cerebral palsy: insufficient quantity and quality of evidence. *Evidence Based Communication Assessment and Intervention* 2014;**8**(1):1–4. DOI: 10.1080/17489539.2014.925270
- Robinson 1983**
Robinson BC. Validation of a Caregiver Strain Index. *Journal of Gerontology* 1983;**38**(3):344–8. [PUBMED: 6841931]
- Rosenbaum 2007**
Rosenbaum P, Paneth N, Leviton A, Goldstein M, Bax M, Damiano D, et al. A report: the definition and classification of cerebral palsy April 2006. *Developmental Medicine & Child Neurology* 2007;**49**(109):8–14. [PUBMED: 17370477]
- Sameroff 2000**
Sameroff AJ, Feise BH. Models of development and developmental risk. In: Zeanah CH Jr editor(s). *Handbook of Infant Mental Health*. New York (NY): Guilford Press, 2000:3–19.
- Schlosser 2014**
Schlosser RW, Balandin S, Helmsley B, Iacono T, Probst T, von Tetzchner S. Facilitated communication and authorship: a systematic review. *Augmentative and*

- Alternative Communication* 2014;**30**(4):359–68. DOI: 10.3109/07434618.2014.971490; PUBMED: 25384895
- Shin 1997**
Shin SJ. *Effects of Parenting Stress, Social Support and Parenting Efficacy on Mothers' Parenting Behavior [PhD thesis]*. Seoul (KR): Yonsei University, 1997.
- Skinner 1958**
Skinner BF. Teaching machine; from the experimental study of learning come devices which arrange optimal conditions for self instruction. *Science* 1958;**128**(3330):969–77. [PUBMED: 13592277]
- Solomon-Rice 2011**
Solomon-Rice P, Soto G. Co-construction as a facilitative factor in supporting the personal narratives of children who use augmentative and alternative communication. *Communication Disorders Quarterly* 2011;**32**(2):70–82. DOI: 10.1177/1525740109354776
- Sparling 2008**
Sparling J, Lewis I. *The Creative Curriculum Learning Games*. Washington (DC): Teaching Strategies LLC, 2008.
- Thomas-Stonell 2010**
Thomas-Stonell NL, Oddson B, Robertson B, Rosenbaum PL. Development of the FOCUS (Focus on the Outcomes of Communication Under Six), a communication outcome measure for preschool children. *Developmental Medicine & Child Neurology* 2010;**52**(1):47–53. DOI: 10.1111/j.1469-8749.2009.03410.x; PUBMED: 19709136
- Watson 2015**
Watson RM, Pennington L. Assessment and management of the communication difficulties of children with cerebral palsy: a UK survey of SLT practice. *International Journal of Language and Communication Disorders* 2015;**50**(2):241–59. DOI: 10.1111/1460-6984.12138; PMC4371637
- Wells 1981**
Wells G. *Language Through Interaction: the Study of Language Development*. New York (NY): Cambridge University Press, 1981.
- Wetherby 2002**
Wetherby A, Prizant BM. *Communication and Symbolic Behavior Scales Developmental Profile™ (CSBS DP™)*. Baltimore (MD): Paul H Brookes Publishing Co, 2002.
- Whittingham 2011**
Whittingham K, Wee D, Boyd R. Systematic review of the efficacy of parenting interventions for children with cerebral palsy. *Child: Care, Health & Development* 2011; **37**(4):475–83. DOI: 10.1111/j.1365-2214.2011.01212.x; PUBMED: 21392055
- WHO 2001**
World Health Organization. *International Classification of Functioning, Disability and Health: ICF*. Geneva: WHO, 2001.
- Zimmerman 2002**
Zimmerman I, Steiner V, Pond R. *Preschool Language Scale - 4*. San Antonio (TX): Pearson, 2002.
- References to other published versions of this review**
- Pennington 2017**
Pennington L, Laws K, Goldbart J. Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders. *Cochrane Database of Systematic Reviews* 2017, Issue 1. DOI: 10.1002/14651858.CD012507
- * Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Girolametto 1988

Methods	Parallel-group randomised controlled trial
Participants	<p>Setting: Toronto, Canada</p> <p>Children</p> <p>Sample size: 20. Intervention = 9 (5 boys, 4 girls); control = 11 (5 boys, 6 girls)</p> <p>Completers: 20/22 children completed the study (intervention = 9/11; control = 11/11)</p> <p>Mean age: intervention = 3.3 (SD 1.2) years; control = 3.1 (SD 1.2) years</p> <p>Diagnoses: 11 = Down's syndrome; 4 = cerebral palsy; 2 = chromosomal abnormalities; 3 = unknown aetiology</p> <p>Developmental measures</p> <ol style="list-style-type: none"> 1. Global developmental age (Griffiths Mental Development Scales, Griffiths 1970): intervention = 16.4 (SD 5.8) months; control = 18.5 (SD 6.8) months 2. Receptive language age (SICD): intervention = 18.4 (SD 7.2) months; control = 19.6 (SD 6.8) months 3. Expressive language age (SICD): intervention = 15.1 (SD 7.1); control = 14.4 (SD 4.5) months 4. All children had good head control and use of hands <p>Parents</p> <p>Sample size: 20 mothers; intervention = 9; control = 11</p> <p>Mean age: intervention = 35.6 (SD 4.6) years; control = 35.1 (SD 6.3) years</p> <p>Mean years of education: intervention = 13 (SD 2.4) years; control = 13.9 (SD 2.7) years</p> <p>Employment: not reported</p>
Interventions	<p>Intervention: It Takes Two to Talk, the Hanen Early Language Parent Programme (Manolson 1985)</p> <p>Method: group teaching, video-coaching, feedback</p> <p>Contents: observing child, following child's lead, contingent responding, creating opportunities for communication. Training provided by certified speech-language pathologist and 2 parent assistants</p> <p>Dosage: 1 full-day workshop and 1 session per week for 11 weeks consisting of 8 × 3-hour group sessions and 3 individual home visits</p> <p>Control: waiting list</p>
Outcomes	<p>Timing of outcome measurement: pretest and post-test, exact timing not specified</p> <p>Child outcomes</p> <ol style="list-style-type: none"> 1. Discourse analysis of parent-child interaction: turn taking, contingent responsiveness, topic control, topic maintenance, responsiveness 2. Communication development: SICD (Hedrick 1975) <p>Parent outcomes</p> <ol style="list-style-type: none"> 1. Discourse analysis of parent-child interaction: turn taking, contingent responsiveness, topic control, topic maintenance, responsiveness <p>All primary outcomes</p>

Notes	Study start and end dates: not known Conflicts of interest: not stated	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not described
Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias) All outcomes	High risk	Parents and interventionists could not be blinded to treatment
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Communication development assessment measure completed by assessor blind to treatment and control group. Blinding of coders in discourse analysis measures not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	2 participants lost in intervention group; 1 due to difficulty attending training, no reason stated for the other
Selective reporting (reporting bias)	Low risk	All expected outcomes reported
Other bias	Unclear risk	None identified

Kim 2005

Methods	Parallel-group randomised controlled trial
Participants	<p>Setting: South Korea</p> <p>Children</p> <p>Sample size: 18 (10 boys, 8 girls); intervention = 10; control = 8</p> <p>Completers: 18/23 children completed the study (intervention = 10/13; control = 8/10)</p> <p>Mean age: intervention = 6.2 (SD 1.7) years; control = 6.1 (1.8) years</p> <p>Diagnoses: mental retardation or developmental disorders by psychiatrist or psychologist. Met DSM-IV criteria for mental retardation, motor skills disorder or pervasive developmental disorders</p> <p>Developmental measures</p> <ol style="list-style-type: none"> 1. Vineland Social Maturity age: intervention = 3.9 (SD 2.5) years; control = 3.9 (SD 2.3) years 2. Cognitive age (Carolina Curriculum, Johnson-Martin 2004): intervention = 31.8 (SD 17.8) months; control = 32.9 (SD 19.5) months

	<p>3. Communication age (Carolina Curriculum, Johnson-Martin 2004): intervention = 25.1 (SD 14.8) months; control = 28.5 (SD 18.4) months</p> <p>4. Social adaptation age (Carolina Curriculum, Johnson-Martin 2004): intervention = 36.4 (SD 19.7) months; control 37.9 (SD 22.4) months</p> <p>5. Fine motor age in months (Carolina Curriculum, Johnson-Martin 2004): intervention = 39.9 (SD 22.3) months; control 37.9 (SD 24.0) months</p> <p>6. Gross motor age in months (Carolina Curriculum, Johnson-Martin 2004): intervention = 37.8 (SD 23.3) months; control 38.4 (SD 24.0) months</p> <p>Parents</p> <p>Sample size: 18 mothers; intervention = 10; control = 8</p> <p>Mean age: intervention = 34.0 (SD 2.9) years; control = 34.0 (SD 5.9) years</p> <p>Mean years of education: intervention = 13.6 (SD 2.1) years; control = 12.6 (2.3) years</p> <p>Employment: 3 mothers employed (intervention = 1; control = 2)</p>	
Interventions	<p>Intervention: Relationship Focussed Intervention, adapted from the Family/Child Curriculum (Mahoney 1999a) and the Principle of Program Learning (Skinner 1958)</p> <p>Method: classroom-based instruction; home-based instruction; feedback and evaluation</p> <p>Content: responsive interaction; level of developmental functioning; interacting with children in play; turn taking; following the child's lead; increasing responsiveness; decreasing directiveness; child development; feedback about parent-child interaction in their daily routine; implementing responsive interaction strategies in the daily routine</p> <p>Dosage: 1.5-2 hours per week, every week for 3 months</p> <p>Control: no parent intervention. Children attended special education, as per intervention group</p>	
Outcomes	<p>Timing of outcome measurement: pretest and post-test, exact timing not specified</p> <p>Child outcomes</p> <p>1. Child Behavior Rating Scale (Mahoney 1999b), rating attention to activity, persistence, initiation, joint attention and co-operation</p> <p>Parent outcomes</p> <p>1. Maternal Behavior Rating Scale (Mahoney 1999a) rating responsiveness, affect, achievement orientation and directiveness</p> <p>2. Korean version of the Parent Stress Index (Abidin 1995)</p> <p>All primary outcomes</p>	
Notes	<p>Study start and end dates: not known</p> <p>Conflicts of interest: none identified</p>	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not stated
Allocation concealment (selection bias)	Unclear risk	Not stated

Kim 2005 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	Parents and interventionists could not be blinded to the intervention
Blinding of outcome assessment (detection bias) All outcomes	Low risk	2 raters, blinded to the intervention condition, coded videotaped observations
Incomplete outcome data (attrition bias) All outcomes	High risk	18/23 children completed the study; 3/13 children dropped out of intervention group (1 hospitalised, 1 left centre, 1 missed 3 intervention sessions), 2/10 children in control group did not complete postintervention measures (1 family refused to be videorecorded, 1 hospitalised)
Selective reporting (reporting bias)	Low risk	All expected outcomes reported
Other bias	Unclear risk	No other sources of bias identified

DSM-IV: Diagnostic and Statistical Manual of Mental Disorders (APA 2000); SD: standard deviation; SICD: Sequenced Inventory of Communication Development.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Adamson 2010	No reference to motor impairment. Cerebral palsy was listed among the aetiologies, but no other reference to motor impairment or physical disability. Part of the same study as Ronski 2010 .
Badr 2006	Participants not reported as having communication difficulties
Barlow 2007	Non-experimental design. Did not target communication development
Buschmann 2008	Focused on evaluation of a diagnostic workup, not interventions to improve language and communication in children
Buschmann 2009	Children did not have motor disorders
Cologon 2017	Children did not have motor disorders; both arms of the study received parent-mediated training, hence no non-parent-mediated comparator
Fey 2006	Children had very mild motor disorders that did not affect their ability to sit unsupported, to manipulate objects, to produce gesture or spoken language

(Continued)

Kaiser 2013	Children did not have motor disorders
Karaaslan 2013	Children did not have motor disorders
Kynø 2012	Intervention not aimed at improving communication
Milgrom 2004	Did not examine parent-mediated intervention to improve communication. Not an RCT
Newnham 2009	Intervention not aimed at improving communication
Romski 2010	Inadequate information on presence or absence of motor disorder among participants. Compared 3 parent-mediated interventions and had no control without parent-mediated intervention
Seifer 1991	Intervention not specifically for children with communication difficulties. Study had a young age group, for which specific communication difficulties may not be identified
Tannock 1992	Only 1 child in 1 arm of study had a motor disorder
Warren 2008	From same study as Fey 2006 . Children had very mild motor disorders that did not affect their ability to sit unsupported, to manipulate objects, to produce gesture or spoken language
Whitmore 2014	From the same study as Romski 2010 ; looked at secondary motor outcomes
Woynaroski 2014	From same study as Fey 2006 . Children had very mild motor disorders that did not affect their ability to sit unsupported, to manipulate objects, to produce gesture or spoken language
Yoder 2002	Children with intellectual disability, 1 participant had 'mild cerebral palsy'

RCT: randomised controlled trial.

Characteristics of studies awaiting assessment *[ordered by study ID]*

[Campbell 1977](#)

Methods	No information available
Participants	No information available
Interventions	No information available
Outcomes	No information available
Notes	Title only, full text yet to be made available

Kogan 1978

Methods	Study design unknown
Participants	Developmentally delayed children aged 3-5 years
Interventions	Parent instruction programme
Outcomes	Changes in parent-child interactions, decrease areas of conflict, and increase self-esteem of both parent and child
Notes	Abstract only, full text yet to be made available

Lee 2012

Methods	Study design unknown
Participants	Children aged 1.5-5 years with developmental delay
Interventions	A home programme intervention
Outcomes	Comprehensive Developmental Inventory for Infants and Toddlers (CDIIT) assessments, including cognition, receptive language, expressive language, gross motor, fine motor, social, and self-care domains, at baseline and after 8 weeks (postintervention)
Notes	Abstract only, yet to receive any response from contacted authors

Characteristics of ongoing studies *[ordered by study ID]*

ACTRN12616000653460

Trial name or title	Community-based parent delivered early detection and intervention program for infants at high risk of cerebral palsy in a low-resource setting: a randomised controlled trial
Methods	Randomised controlled trial
Participants	Inclusion criteria: children with cerebral palsy or at risk of cerebral palsy, scored by a certified General Movements Assessors. Aged 12-40 weeks' corrected age Exclusion criteria: no developmental concerns at 12-40 weeks' chronological age Sample size: 142 Recruitment: through developmental healthcare professionals and community workers
Interventions	Intervention: multidisciplinary, family-centred intervention delivered peer-to-peer for 15 fortnightly visits and then delivered by carer to infant. The intervention is based on games and strategies based on the Creative Curriculum Learning Games (Sparling 2008) Control: health advice during 15 fortnightly visits
Outcomes	Primary outcomes 1. Infant functional abilities measured using the mobility domain of Pediatric Evaluation of Disability

	<p>Inventory - Computer Adaptive Test (PEDI-CAT)</p> <p>2. Carers' mental health assessed using Depression, Anxiety, Stress Scale - Short Form (DASS)</p> <p>Secondary outcomes</p> <ol style="list-style-type: none"> 1. Infant cognitive development (Bayley Scales of Infant Development III) 2. Infant nutritional status (weight, body mass index, height/length, head circumference, mean upper arm circumference) 3. Health resource use 4. Infant motor skills (Peabody Developmental Motor Scales) 5. Infant neurological status (Hammersmith Infant Neurological Examination) 6. Quality and extent of stimulation at home (HOME Inventory) 7. Near vision detection scale and Canadian occupational performance measure <p>Timing of outcome measurement: baseline and postintervention (18 months' chronological age)</p>
Starting date	<p>Date registered: 19 May 2016</p> <p>Actual start of recruitment: 13 March 2017</p> <p>Status: recruiting</p>
Contact information	Dr Katherine Benfer. Email: k.benfer@uq.edu.au
Notes	<p>Public title: LEAP-CP: Learning through everyday activities with parents for infants at high risk of cerebral palsy in a low-income country</p> <p>ANZCTR identifier: ACTRN12616000653460</p> <p>Funding source: Endeavour Queen Elizabeth II Diamond Jubilee Scholarship</p> <p>Conflict of interest: none identified</p>

Eliasson 2016

Trial name or title	Efficacy of the small step program in a randomised controlled trial for infants below age 12 months with clinical signs of cerebral palsy
Methods	Parallel-group randomised controlled trial
Participants	<p>Inclusion criteria: 3- to 8-month old children at risk for cerebral palsy due to early neonatal events affecting the brain</p> <p>Exclusion criteria: participants with unstable health, uncontrolled epilepsy, progressive disorders or diagnosis with a specific syndrome</p> <p>Sample size: 30 (15 in each arm)</p> <p>Recruitment: hospital-based recruitment. Participants will be recruited based on neurological signs of cerebral palsy, with a formal diagnosis made at 2 years</p>
Interventions	<p>Intervention: small step programme, an intervention targeting hand use, mobility and communication, delivered by parents who will be coached weekly by therapists. Training and intervention goals will be individualised to the child's need</p> <p>Control: usual care, the hospital follow-up programme, not standardised, but includes instructions to parents regarding home training</p>
Outcomes	<p>Primary outcome</p> <ol style="list-style-type: none"> 1. Peabody Developmental Motor Scale (second edition, Folio 2000)

	<p>Timing of outcome measurement: baseline, 14 weeks' and 35 weeks' postintervention, plus additional measures at 7 and 21 weeks' postintervention and at 2 years' adjusted age</p> <p>Secondary outcomes</p> <ol style="list-style-type: none"> 1. Child development 2. Specific brain pathology 3. Parents' perspective of the programme <p>Timing of outcome measurement: baseline, 14 weeks', 35 weeks' and 2 years' postintervention</p>
Starting date	<p>Study start date: January 2014</p> <p>Status: recruiting</p>
Contact information	<p>Linda Holmström, PhD. Email: Linda.Holmstrom@ki.se</p> <p>Ann-Christin Eliasson, PhD. Email: ann-christin.eliaasson@ki.se</p>
Notes	<p>ClinicalTrials.gov identifier: NCT02166801</p> <p>Funding source: Swedish Research Council (grant number 521-2013-3096), Special grants supporting Stockholm City Council (LS 1411-1372), Stiftelsen Frimurare-Barnhuset in Stockholm, Foundation Sunnerdahls Handikappfond, Olle Engkvist Byggmästare, Promobilia, Norrbacka-Eugenia Stiftelsen, Stiftelsen and Sven Jerrings Fond</p> <p>Official title: A small-step program for development of new treatment principles for children with cerebral palsy and other neurodevelopmental disorders</p>

DATA AND ANALYSES

Comparison 1. Parent-mediated intervention versus no intervention: child communicative participation outcomes

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Initiating conversation	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
2 Joint attention	1		Mean Difference (IV, Random, 95% CI)	Totals not selected

Comparison 2. Parent-mediated intervention versus no intervention: parents' communication and interaction strategies

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Maternal Behavior Rating Scale (MBRS) Responsiveness	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
2 MBRS Directiveness	1		Mean Difference (IV, Random, 95% CI)	Totals not selected

Comparison 3. Parent-mediated intervention versus no intervention: family stress and coping

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Parent Stress Index: child temperament	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
2 Parent Stress Index: parent-child relationship	1		Mean Difference (IV, Random, 95% CI)	Totals not selected
3 Parent Stress Index: learning expectation	1		Mean Difference (IV, Random, 95% CI)	Totals not selected

Analysis 1.1. Comparison 1 Parent-mediated intervention versus no intervention: child communicative participation outcomes, Outcome 1 Initiating conversation.

Review: Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders

Comparison: 1 Parent-mediated intervention versus no intervention: child communicative participation outcomes

Outcome: 1 Initiating conversation

Analysis 1.2. Comparison 1 Parent-mediated intervention versus no intervention: child communicative participation outcomes, Outcome 2 Joint attention.

Review: Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders

Comparison: 1 Parent-mediated intervention versus no intervention: child communicative participation outcomes

Outcome: 2 Joint attention

Analysis 2.1. Comparison 2 Parent-mediated intervention versus no intervention: parents' communication and interaction strategies, Outcome 1 Maternal Behavior Rating Scale (MBRS) Responsiveness.

Review: Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders

Comparison: 2 Parent-mediated intervention versus no intervention: parents' communication and interaction strategies

Outcome: 1 Maternal Behavior Rating Scale (MBRS) Responsiveness

Analysis 2.2. Comparison 2 Parent-mediated intervention versus no intervention: parents' communication and interaction strategies, Outcome 2 MBRS Directiveness.

Review: Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders

Comparison: 2 Parent-mediated intervention versus no intervention: parents' communication and interaction strategies

Outcome: 2 MBRS Directiveness

Analysis 3.1. Comparison 3 Parent-mediated intervention versus no intervention: family stress and coping, Outcome 1 Parent Stress Index: child temperament.

Review: Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders

Comparison: 3 Parent-mediated intervention versus no intervention: family stress and coping

Outcome: 1 Parent Stress Index: child temperament

Analysis 3.2. Comparison 3 Parent-mediated intervention versus no intervention: family stress and coping, Outcome 2 Parent Stress Index: parent-child relationship.

Review: Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders

Comparison: 3 Parent-mediated intervention versus no intervention: family stress and coping

Outcome: 2 Parent Stress Index: parent- child relationship

Analysis 3.3. Comparison 3 Parent-mediated intervention versus no intervention: family stress and coping, Outcome 3 Parent Stress Index: learning expectation.

Review: Parent-mediated communication interventions for improving the communication skills of preschool children with non-progressive motor disorders

Comparison: 3 Parent-mediated intervention versus no intervention: family stress and coping

Outcome: 3 Parent Stress Index: learning expectation

ADDITIONAL TABLES

Table 1. Methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions

Types of outcome measures	<p>We will compare baseline measures with outcomes grouped into the following time points: short term (0-1 month following intervention completion), medium term (2-5 months after intervention) and long term (≥ 6 months following intervention)</p> <p>We will combine results from studies where tools measure the same outcome using the same type of data (e.g. frequency of child communication behaviours; standard scores on child language measures)</p>
Electronic searches	We will seek translations of papers published in languages other than English, when necessary
Measures of treatment effect	<p>Binary data For binary data, we will calculate an OR with a 95% CI.</p> <p>Continuous We will report the effect size as an MD with 95% CI if studies have use the same continuous outcome measure. For studies that evaluate the same construct using different continuous outcome measures that share the same method of administration (e.g. questionnaires; frequency counts of behaviours measured in direct observation), we will summarise results using the SMD with 95% CI</p>
Unit of analysis issues	<p>Cluster-randomised trials We will use the <i>Cochrane Handbook of Systematic Reviews of Interventions</i></p>

Table 1. Methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions (Continued)

	<p>for advice on analysis of cluster-randomised trials (Higgins 2011b).</p> <hr/> <p>Cross-over trials In cross-over trials, we will include data from the first period only, so as not to include data from the same participant twice</p> <hr/> <p>Studies with multiple treatment groups If a study investigates multiple treatment groups, we will make single pair-wise comparisons by combining data from all eligible parent-mediated intervention groups and comparing these with data combined from all eligible control groups</p>
Assessment of heterogeneity	We will assess heterogeneity in clinical characteristics of study samples (e.g. parents' years in education; ratio of mothers to fathers in group composition; children's age; level of communication development; use of AAC) using the Chi^2 test to assess if statistical heterogeneity is likely to be due to chance alone. We will use the I^2 test and Tau^2 to describe the variation in effect estimates that is due to heterogeneity rather than sampling error (Higgins 2002).
Assessment of reporting bias	If we identify > 10 studies that fit the inclusion criteria, we will use funnel plots of effect estimates to assess the possibility of publication bias on primary outcomes, and we will use Egger's test to test for funnel plot asymmetry (Egger 1997).
Data synthesis	We will undertake meta-analysis using RevMan 5 applying a random-effects model, if two or more studies report interventions that are similar in terms of topic, delivery methods and dosage (duration, frequency and intensity of sessions), and include similar participants (parents and children) and use similar outcome measures (Review Manager 2014).
Subgroup analysis and investigation of heterogeneity	<p>If we identify multiple studies with sufficiently similar participants, interventions and outcome measures, we will conduct the following subgroup analyses to explore possible sources of heterogeneity</p> <ol style="list-style-type: none"> 1. In the presence of severe or profound intellectual or receptive language impairment (impairment in either function ≥ -1.9 standard deviations versus non-verbal or receptive language score < -2 standard deviations) 2. In parental education (high school versus further or higher education) 3. In dosage of intervention (frequency and duration of sessions) and 4. Between specific 'named' interventions (e.g. Hanen programmes (see, for example, Pepper 2004) or Enhanced Milieu Teaching (see, for example, Hemmeter 1994)).
Sensitivity analysis	We will use the 'Risk of bias' assessment to inform sensitivity analyses. As it is difficult to blind parents and training providers to the type of intervention, sensitivity analyses will use data from risk of bias arising from

Table 1. Methods stipulated in the protocol that were not required in this version of the review, but which will be applied in future versions (Continued)

	random allocation generation, allocation concealment, loss to follow-up and incomplete reporting of outcomes. We will remove studies judged to have a high risk of bias in these areas to determine their effect on the pooled estimate. We will undertake a sensitivity analysis of binary outcomes if data are considered missing at random, adopting both a best-case and worst-case scenario in which, for example, children in the experimental group are imputed to have a good outcome (best case) and poor outcome (worst outcome)
--	--

ACC: augmentative and alternative communication; CI: confidence interval; MD: mean difference; OR: odds ratio; SMD: standardised mean difference.

APPENDICES

Appendix I. Search strategies

Cochrane Central Register of Controlled Trials (CENTRAL) in the Cochrane Library, and which includes the Cochrane Developmental, Psychosocial and Learning Problems Specialised Register

```
#1[mh ^"Cerebral Palsy"]
#2cerebral pals*
#3(ataxic or ataxia*)
#4spastic*
#5dyskinetic*
#6(non-progressive near/3 (disabilit* or disorder* or impair*))
#7[mh ^"Movement Disorders"]
#8[mh "motor disorders"]
#9[mh ^"motor skills disorder"]
#10[mh "Motor Skills"]
#11(movement near/3 (disorder* or disabilit* or impair*))
#12(motor near/3 (dysfunction* or dysfunction* or disorder* or disabilit* or impair*))
#13[mh ^"Brain Injuries"]
#14[mh "Brain Damage, Chronic"]
#15[mh "Brain injury, chronic"]
#16("acquired brain injur*" or ABI)
#17[mh ^"Developmental disabilities"]
#18(developmental* near/3 (delay* or disab*))
#19[mh ^"Down Syndrome"]
#20down* next syndrome*
#21[mh dyskinesias]
#22[mh Chorea]
#23(dyskinesia* or chorea*)
#24[mh "Angelman Syndrome"]
#25((Angelman* or "Happy Puppet") near/1 syndrome*)
```

#26(GLUT1 or GLUT-1)
 #27(PRRT2 or PRRT-2)
 #28{or #1-#27}
 #29[mh Infant]
 #30[mh child]
 #31(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten* or kindergarten*)
 #32{or #29-#31}
 #33#28 and #32
 #34[mh "Parent-Child Relations"]
 #35[mh Parents]
 #36[mh Parenting]
 #37[mh Family]
 #38[mh Caregivers]
 #39("at home" or (in near/3 home) or "home based")
 #40(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or famil* or adult* or teacher* or therapist*)
 #41{or #34-#40}
 #42[mh education]
 #43[mh teaching]
 #44[mh "Early intervention (Education)"]
 #45"early intervent*"

#46[mh "Education of Intellectually Disabled"]
 #47[mh "education, special"]
 #48[mh "speech therapy"]
 #49[mh "language therapy"]
 #50[mh "speech-language pathology"]
 #51[mh "speech disorders"]
 #52[mh "language disorders"]
 #53[mh "Sign language"]
 #54[mh "Manual Communication"]
 #55[mh "Nonverbal communication"]
 #56[mh "Communication Aids for Disabled"]
 #57[mh "Self-Help Devices"]
 #58[mh ^communication]
 #59[mh "communication disorders"]
 #60(speech* or languag* or communicat* or sign* or nonverbal* or non-verbal* or nonvocal* or non-vocal* or cue* or symbol* or picture* or pictorial* or icon* or augment*)
 #61{or #42-#60}
 #62#61 and #41
 #63((parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*) near/3 (coach* or educat* or intervention* or learn* or program* or teach* or train*))
 #64((parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*) near/3 (interact* or inter-act* or involv* or mediat* or respon*))
 #65(naturalistic near/2 (teach* or approach* or setting* or environment*))
 #66"focus*ed stimulation"
 #67(milieu near/2 teaching)
 #68(responsiv* near/2 education)
 #69(responsiv* near/2 teaching)
 #70Hanen*
 #71{or #62-#70}
 #72#33 and #71 in Trials

MEDLINE Ovid

1 Cerebral Palsy/
2 cerebral pals\$.tw,kf.
3 (ataxic or ataxia\$).tw,kf.
4 spastic\$.tw,kf.
5 dyskinetic\$.tw,kf.
6 (non-progressive adj3 (disabilit\$ or disorder\$ or impair\$)).tw,kf.
7 Movement Disorders/
8 motor disorders/
9 motor skills disorders/
10 Motor Skills/
11 (movement adj3 (disorder\$ or disabilit\$ or impair\$)).tw,kf.
12 (motor adj3 (disfunction\$ or dysfunction\$ or disorder\$ or disabilit\$ or impair\$)).tw,kf.
13 Brain Injuries/
14 Brain Damage, Chronic/
15 brain injury, chronic/
16 (acquired brain injur\$ or ABI).tw,kf.
17 Developmental disabilities/
18 (developmental\$ adj3 (delay\$ or disab\$)).tw,kf.
19 Down Syndrome/
20 down\$ syndrome\$.tw,kf.
21 dyskinesias/
22 Chorea/
23 (dyskinesia\$ or chorea\$).tw,kf.
24 Angelman Syndrome/
25 Angelman\$ syndrome\$.tw,kf.
26 (GLUT1 or GLUT-1).tw,kf.
27 (PRRT2 or PRRT-2).tw,kf.
28 or/1-27
29 Infant/
30 exp child/
31 (child\$ or infant\$ or babies or baby or toddler\$ or girl\$ or boy\$ or pre-school\$ or preschool\$ or nurser\$ or kindergarten\$ or kindergarten\$).tw,kf.
32 or/29-31
33 28 and 32
34 exp Parent-Child Relations/
35 exp Parents/
36 Parenting/
37 Family/
38 Caregivers/
39 (at home or (in adj3 home) or home-based).tw,kf.
40 (parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or famil\$ or adult\$ or teacher\$ or therapist\$).tw,kf.
41 or/34-40
42 education/
43 teaching/
44 "Early intervention (Education)"/
45 early intervent\$.tw,kf.
46 Education of Intellectually Disabled/
47 education, special/
48 speech therapy/
49 language therapy/

50 speech-language pathology/
 51 speech disorders/
 52 language disorders/
 53 Sign language/
 54 Manual Communication/
 55 Nonverbal communication/
 56 Communication Aids for Disabled/
 57 Self-Help Devices/
 58 communication/
 59 communication disorders/
 60 early intervent\$.tw,kf.
 61 (speech\$ or languag\$ or communicat\$ or sign\$1 or signing or nonverbal\$ or non-verbal\$ or nonvocal\$ or non-vocal\$ or cue\$ or symbol\$ or picture\$ or pictorial\$ or icon\$ or augment\$).tw,kf.
 62 or/42-61
 63 exp Parents/ed
 64 Caregivers/ed
 65 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$) adj3 (coach\$ or educat\$ or intervention\$ or learn\$ or program\$ or teach\$ or train\$)).tw,kf.
 66 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$) adj3 (interact\$ or inter-act\$ or involv\$ or mediat\$ or respon\$)).tw,kf.
 67 or/63-66
 68 (naturalistic adj2 (teach\$ or approach\$ or setting\$ or environment\$)).tw,kf.
 69 focus?ed stimulation.tw,kf.
 70 (milieu adj2 teaching).tw,kf.
 71 (responsiv\$ adj2 education).tw,kf.
 72 (responsiv\$ adj2 teaching).tw,kf.
 73 Hanen\$.tw,kf.
 74 or/68-73
 75 randomized controlled trial.pt.
 76 controlled clinical trial.pt.
 77 randomi#ed.ab.
 78 placebo\$.ab.
 79 drug therapy.fs.
 80 randomly.ab.
 81 trial.ab.
 82 groups.ab.
 83 or/75-82
 84 exp animals/ not humans.sh.
 85 83 not 84
 86 33 and 67 and 85

MEDLINE In-Process & Other Non-Indexed Citations Ovid

1 cerebral pals\$.tw,kf.
 2 (ataxic or ataxia\$).tw,kf.
 3 spastic\$.tw,kf.
 4 dyskinetic\$.tw,kf.
 5 (non-progressive adj3 (disabilit\$ or disorder\$ or impair\$)).tw,kf.
 6 (movement adj3 (disorder\$ or disabilit\$ or impair\$)).tw,kf.
 7 (motor adj3 (dysfunction\$ or dysfunction\$ or disorder\$ or disabilit\$ or impair\$ or skills\$)).tw,kf.
 8 (acquired brain injur\$ or ABI).tw,kf.
 9 (developmental\$ adj3 (delay\$ or disab\$)).tw,kf.
 10 down\$ syndrome\$.tw,kf.

11 ((Angelman\$ or Happy Puppet) adj1 syndrome\$.tw,kf.
 12 (GLUT1 or GLUT-1).tw,kf.
 13 (PRRT2 or PRRT-2).tw,kf.
 14 or/1-13
 15 (child\$ or infant\$ or babies or baby or toddler\$ or girl\$ or boy\$ or pre-school\$ or preschool\$ or nurser\$ or kindergarten\$ or kindergarten\$.tw,kf.
 16 14 and 15
 17 (parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or famil\$ or adult\$ or teacher\$ or therapist\$.tw,kf.
 18 (at home or (in adj3 home) or home-based).tw,kf.
 19 17 or 18
 20 (speech\$ or languag\$ or communicat\$ or sign\$ or symbol\$ or nonverbal\$ or non-verbal\$ or nonvocal\$ or non-vocal\$ or cue\$ or picture\$ or pictorial\$ or augment\$.tw,kf.
 21 19 and 20
 22 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$) adj3 (coach\$ or educat\$ orv intervention\$ or learn\$ or program\$ or teach\$ or train\$)).tw,kf.
 23 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$) adj3 (interact\$ or inter-act\$ or involv\$ or mediat\$ or respon\$)).tw,kf.
 24 (naturalistic adj2 (teach\$ or approach\$ or setting\$ or environment\$)).tw,kf.
 25 focus?ed stimulation.tw,kf.
 26 (milieu adj2 teaching).tw,kf.
 27 (responsiv\$ adj2 education).tw,kf.
 28 (responsiv\$ adj2 teaching).tw,kf.
 29 Hanen\$.tw,kf.
 30 or/22-29
 31 21 or 30
 32 16 and 31
 33 (random\$ or control\$ or group\$ or cluster\$ or placebo\$ or trial\$ or assign\$ or prospectiv\$ or meta-analysis or systematic review or longitudinal\$.tw,kf.
 34 32 and 33

MEDLINE Epub Ahead of Print Ovid

1 cerebral pals\$.tw,kf.
 2 (ataxic or ataxia\$.tw,kf.
 3 spastic\$.tw,kf.
 4 dyskinetic\$.tw,kf.
 5 (non-progressive adj3 (disabilit\$ or disorder\$ or impair\$)).tw,kf.
 6 (movement adj3 (disorder\$ or disabilit\$ or impair\$)).tw,kf.
 7 (motor adj3 (dysfunction\$ or dysfunction\$ or disorder\$ or disabilit\$ or impair\$ or skills\$)).tw,kf.
 8 (acquired brain injur\$ or ABI).tw,kf.
 9 (developmental\$ adj3 (delay\$ or disab\$)).tw,kf.
 10 down\$ syndrome\$.tw,kf.
 11 ((Angelman\$ or Happy Puppet) adj1 syndrome\$.tw,kf.
 12 (GLUT1 or GLUT-1).tw,kf.
 13 (PRRT2 or PRRT-2).tw,kf.
 14 or/1-13
 15 (child\$ or infant\$ or babies or baby or toddler\$ or girl\$ or boy\$ or pre-school\$ or preschool\$ or nurser\$ or kindergarten\$ or kindergarten\$.tw,kf.
 16 14 and 15
 17 (parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or famil\$ or adult\$ or teacher\$ or therapist\$.tw,kf.
 18 (at home or (in adj3 home) or home-based).tw,kf.

19 17 or 18
 20 (speech\$ or languag\$ or communicat\$ or sign\$ or symbol\$ or nonverbal\$ or non-verbal\$ or nonvocal\$ or non-vocal\$ or cue\$ or picture\$ or pictorial\$ or augment\$).tw,kf.
 21 19 and 20
 22 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$) adj3 (coach\$ or educat\$ or intervention\$ or learn\$ or program\$ or teach\$ or train\$)).tw,kf.
 23 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$) adj3 (interact\$ or inter-act\$ or involv\$ or mediat\$ or respon\$)).tw,kf.
 24 (naturalistic adj2 (teach\$ or approach\$ or setting\$ or environment\$)).tw,kf.
 25 focus?ed stimulation.tw,kf.
 26 (milieu adj2 teaching).tw,kf.
 27 (responsiv\$ adj2 education).tw,kf.
 28 (responsiv\$ adj2 teaching).tw,kf.
 29 Hanen\$.tw,kf.
 30 or/22-29
 31 21 or 30
 32 16 and 31
 33 (random\$ or control\$ or group\$ or cluster\$ or placebo\$ or trial\$ or assign\$ or prospectiv\$ or meta-analysis or systematic review or longitudinal\$).tw,kf.
 34 32 and 33

Embase Ovid

1 cerebral palsy/
 2 cerebral pals\$.tw,kw.
 3 (ataxic or ataxia\$).tw,kw.
 4 spastic\$.tw,kw.
 5 dyskinetic\$.tw,kw.
 6 (non-progressive adj3 (disabilit\$ or disorder\$ or impair\$)).tw,kw.
 7 motor dysfunction/
 8 motor performance/
 9 psychomotor disorder/
 10 (movement adj3 (disorder\$ or disabilit\$ or impair\$)).tw,kw.
 11 (motor adj3 (disorder\$ or disabilit\$ or disfunction\$ or dysfunction\$ or impair\$)).tw,kw.
 12 brain injury/
 13 brain damage/
 14 (acquired brain injur\$ or ABI).tw,kw.
 15 developmental disorder/
 16 (developmental\$ adj3 (delay\$ or disab\$)).tw,kw.
 17 Down Syndrome/)
 18 down\$ syndrome\$.tw,kw.
 19 dyskinesia/
 20 chorea/
 21 (dyskinesia\$ or chorea\$).tw,kw.
 22 happy puppet syndrome/
 23 ((Angelman or happy puppet) adj1 syndrome).tw,kw.
 24 (GLUT1 or GLUT-1).tw,kw.
 25 (PRRT2 or PRRT-2).tw,kw.
 26 or/1-25
 27 exp child/
 28 (child\$ or infant\$ or babies or baby or toddler\$ or girl\$ or boy\$ or pre-school\$ or preschool\$ or nurser\$ or kindergarten\$ or kindergarten\$).tw,kw.
 29 27 or 28

30 26 and 29
 31 exp child parent relation/
 32 exp parent/
 33 family/
 34 caregiver/
 35 or/31-34
 36 education/
 37 teaching/
 38 early childhood intervention/
 39 "education of intellectually disabled"/
 40 special education/
 41 speech therapy/
 42 language therapy/
 43 exp communication disorder/
 44 speech disorder/
 45 sign language/
 46 exp nonverbal communication/
 47 communication aid/
 48 or/36-47
 49 35 and 48
 50 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$)
 adj3 (coach\$ or educat\$ or intervention\$ or learn\$ or program\$ or teach\$ or train\$)).tw,kw.
 51 ((at home or (in adj3 home) or home-based) adj3 (coach\$ or educat\$ or intervention\$ or learn\$ or program\$ or teach\$ or
 train\$)).tw,kw.
 52 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$)
 adj3 (interact\$ or inter-act\$ or involv\$ or mediat\$ or respon\$)).tw,kw.
 53 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$)
 adj5 (augment\$ or communicat\$ or cue\$ or icon\$ or languag\$ or nonverbal\$ or non-verbal\$ or nonvocal\$ or non-vocal\$ or picture\$
 or pictorial\$ or sign\$1 or signing or speech\$)).tw,kw.
 54 (naturalistic adj2 (teach\$ or approach\$ or setting\$ or environment\$)).tw,kw.
 55 focus?ed stimulation.tw,kw.
 56 (milieu adj2 teaching).tw,kw.
 57 (responsiv\$ adj2 education).tw,kw.
 58 (responsiv\$ adj2 teaching).tw,kw.
 59 Hanen\$.tw,kw.
 60 or/49-59
 61 Randomized controlled trial/
 62 controlled clinical trial/
 63 Single blind procedure/
 64 Double blind procedure/
 65 triple blind procedure/
 66 Crossover procedure/
 67 (crossover or cross-over).tw.
 68 ((singl\$ or doubl\$ or tripl\$ or trebl\$) adj1 (blind\$ or mask\$)).tw.
 69 Placebo/
 70 placebo.tw.
 71 prospective.tw.
 72 factorial\$.tw.
 73 random\$.tw.
 74 assign\$.ab.
 75 allocat\$.tw.
 76 volunteer\$.ab.
 77 or/61-76

CINAHL Plus EBSCOhost (Cumulative Index to Nursing and Allied Health Literature)

S84S64 AND S83
S83S65 OR S66 OR S67 OR S68 OR S69 OR S70 OR S71 OR S72 OR S73 OR S74 OR S75 OR S76 OR S77 OR S78 OR S79
OR S80 OR S81 OR S82
S82TI(random* OR RCT or allocat* OR assign* OR group* or placebo*) OR AB(random* OR RCT or allocat* OR assign* OR
group* or placebo*)
S81TI (evaluat* N2 study or evaluat* N2 research) OR AB (evaluat* N2 study or evaluat* N2 research)
S80(MH "Evaluation Research")
S79TI ((tripl* N2 mask*) or (tripl* N2 blind*)) or ((trebl* N2 mask*) or (trebl* N2 blind*)) OR AB((tripl* N2 mask*) or (tripl* N2
blind*)) or ((trebl* N2 mask*) or (trebl* N2 blind*))
S78TI((doubl* N2 mask*) or (doubl* N2 blind*)) OR AB((doubl* N2 mask*) or (doubl* N2 blind*))
S77TI(single N2 mask* or single N2 blind*) OR AB(single N2 mask* or single N2 blind*)
S76(MH "Treatment Outcomes")
S75(MH "Program Evaluation")
S74(prospectiv* study or prospectiv* research)
S73("follow-up study" or "follow-up research")
S72(clinical trial*) or (control* N2 trial*)
S71PT Clinical trial
S70PT randomized controlled trial
S69(MH "Quantitative Studies")
S68(MH "Crossover Design")
S67(MH "Meta Analysis")
S66MH random assignment
S65(MH "Clinical Trials+")
S64S29 AND S63
S63S53 OR S54 OR S55 OR S56 OR S57 OR S58 OR S59 OR S60 OR S61 OR S62
S62Hanen*
S61(responsiv* N2 (education or teaching))
S60(milieu N2 teaching)
S59focus*ed stimulation
S58(naturalistic N2 (teach* or approach* or setting* or environment*))
S57TI((at home or (in N1 home) or home-based) N3 (coach* or educat* or intervention* or learn* or program* or teach* or train*))OR
AB((at home or (in N1 home) or home-based) N3 (coach* or educat* or intervention* or learn* or program* or teach* or train*))
S56TI(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or famil* or adult* or teacher*
or therapist*) N3 (speech* or languag* or communicat* or sign* or nonverbal* or non-verbal* or nonvocal* or non-vocal* or cue* or
symbol* or picture* or pictorial* or icon* or augment*) OR AB(parent* or maternal* or mother* or father* or paternal* or carer* or
caregiver* or care-giver* or famil* or adult* or teacher* or therapist*) N3 (speech* or languag* or communicat* or sign* or nonverbal*
or non-verbal* or nonvocal* or non-vocal* or cue* or symbol* or picture* or pictorial* or icon* or augment*)
S55TI(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*)
N3 (interact* or inter-act* or involv* or mediat* or respon*)OR AB(parent* or maternal* or mother* or father* or paternal* or carer*
or caregiver* or care-giver* or adult* or teacher* or therapist*) N3 (interact* or inter-act* or involv* or mediat* or respon*)
S54TI(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*)
N3 (coach* or educat* or intervention* or learn* or program* or teach* or train*) OR AB(parent* or maternal* or mother* or father*
or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*) N3 (coach* or educat* or intervention* or learn*
or program* or teach* or train*) or TI(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or
adult* or teacher* or therapist*) N3 (coach* or educat* or intervention* or learn* or program* or teach* or train*)
S53S35 AND S52
S52S36 OR S37 OR S38 OR S39 OR S40 OR S41 OR S42 OR S43 OR S44 OR S45 OR S46 OR S47 OR S48 OR S49 OR S50
OR S51
S51(MH "Language Disorders")

S50(MH "Speech Disorders")
 S49(MH "Alternative and Augmentative Communication")
 S48(MH "Communication Aids for Disabled")
 S47(MH "Communication Skills Training")
 S46(MH "Communication Skills")
 S45(MH "Sign Language")
 S44(MH "Nonverbal Communication+")
 S43(MH "Communication")
 S42(MH "Speech-Language Pathology")
 S41(MH "Language Therapy")
 S40(MH "Speech Therapy")
 S39(MH "Education, Special")
 S38(MH "Early Childhood Intervention")
 S37(MH "Teaching")
 S36(MH "Education")
 S35S30 OR S31 OR S32 OR S33 OR S34
 S34(MH "Caregivers")
 S33(MH "Family")
 S32(MH "Parenting")
 S31(MH "Parent-Child Relations+")
 S30(MH "Parents+")
 S29S25 AND S28
 S28S26 OR S27
 S27TI(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten* or kindergarten*) OR AB(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten* or kindergarten*)
 S26(MH "Infant+") OR (MH "Child+")
 S25S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13 OR S14 OR S15 OR S16 OR S17 OR S18 OR S19 OR S20 OR S21 OR S22 OR S23 OR S24
 S24TI(PRRT2 or PRRT-2) OR AB(PRRT2 or PRRT-2)
 S23TI(GLUT1 or GLUT-1) OR AB(GLUT1 or GLUT-1)
 S22TI((Angelman* OR HAPPY PUPPET) N1 syndrome*) OR AB((Angelman* OR HAPPY PUPPET) N1 syndrome*)
 S21(MH "Angelman Syndrome")
 S20TI(dyskinesia* or chorea*) OR AB(dyskinesia* or chorea*)
 S19(MH "Chorea")
 S18(MH "Dyskinesias")
 S17TI(down* syndrome*) OR AB(down* syndrome*)
 S16MH "Down Syndrome"
 S15TI (developmental* N3 (delay* or disab*)) OR AB (developmental* N3 (delay* or disab*))
 S14(MH "Developmental Disabilities")
 S13TI (acquired brain injur* or ABI) OR AB (acquired brain injur* or ABI)
 S12(MH "Brain Damage, Chronic") OR (MH "Brain Injuries")
 S11TI (motor N3 (disfunction* or dysfunction* or disorder* or disabilit* or impair*)) OR AB (motor N3 (disfunction* or dysfunction* or disorder* or disabilit* or impair*))
 S10TI (movement N3 (disorder* or disabilit* or impair*)) OR AB (movement N3 (disorder* or disabilit* or impair*))
 S9(MH "Motor Skills")
 S8(MH "Motor Skills Disorders")
 S7(MH "Movement Disorders")
 S6(non-progressive N3 (disabilit* or disorder* or impair*))
 S5TI(dyskinetic*) OR AB(dyskinetic*)
 S4TI(spastic*) or AB(spastic*)
 S3TI (ataxic or ataxia*) or AB(ataxic or ataxia*)
 S2TI(cerebral pals*) or AB(cerebral pals*)

S1(MH “Cerebral Palsy”)

PsycINFO Ovid

- 1 cerebral palsy/
- 2 cerebral pals\$.tw,id.
- 3 (ataxic or ataxia\$).tw,id.
- 4 spastic\$.tw,id.
- 5 dyskinetic\$.tw,id.
- 6 (non-progressive adj3 (disabilit\$ or disorder\$ or impair\$)).tw,id.
- 7 movement disorders/
- 8 nervous system disorders/
- 9 (movement adj3 (disorder\$ or disabilit\$ or impair\$)).tw,id.
- 10 Motor skills/
- 11 (motor adj3 (disfunction\$ or dysfunction\$ or disorder\$ or disabilit\$ or impair\$)).tw,id.
- 12 Brain Damage/
- 13 Traumatic Brain Injury/
- 14 (acquired brain injur\$ or ABI).tw,id.
- 15 developmental disabilities/
- 16 (developmental\$ adj3 (delay\$ or disab\$)).tw,id.
- 17 Down's syndrome/
- 18 down\$ syndrome\$.tw,id.
- 19 dyskinesia/
- 20 chorea/
- 21 (dyskinesia\$ or chorea\$).tw,id.
- 22 ((Angelman\$ or Happy Puppet) adj1 syndrome\$).tw,id.
- 23 (GLUT1 or GLUT-1).tw,id.
- 24 (PRRT2 or PRRT-2).tw,id.
- 25 or/1-24
- 26 (“100” or “120” or “140” or “160”).ag.
- 27 (child\$ or infant\$ or babies or baby or toddler\$ or girl\$ or boy\$ or pre-school\$ or preschool\$ or nurser\$ or kindergarten\$ or kindergarten\$).tw,id.
- 28 26 or 27
- 29 25 and 28
- 30 exp parents/
- 31 family members/
- 32 PARENTING/
- 33 parenting skills/
- 34 parent child relations/
- 35 parent training/
- 36 caregivers/
- 37 or/30-36
- 38 Education/
- 39 special education/
- 40 early intervention/
- 41 teaching/
- 42 early intervent\$.tw,id.
- 43 Speech Therapy/
- 44 Language Therapy/
- 45 speech-language pathology
- 46 sign language/
- 47 nonverbal communication/
- 48 exp augmentative communication/

49 communication/
 50 communication disorders/
 51 communication skills/
 52 communication skills training/
 53 language disorders/
 54 speech disorders/
 55 or/38-54
 56 37 and 55
 57 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$) adj3 (coach\$ or educat\$ or intervention\$ or learn\$ or program\$ or teach\$ or train\$)).tw,id.
 58 ((at home or (in adj3 home) or home-based) adj3 (coach\$ or educat\$ or intervention\$ or learn\$ or program\$ or teach\$ or train\$)).tw,id.
 59 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$) adj3 (interact\$ or inter-act\$ or involv\$ or mediat\$ or respon\$)).tw,id.
 60 ((parent\$ or maternal\$ or mother\$ or father\$ or paternal\$ or carer\$ or caregiver\$ or care-giver\$ or adult\$ or teacher\$ or therapist\$) adj3 (augment\$ or communicat\$ or cue\$ or icon\$ or languag\$ or nonverbal\$ or non-verbal\$ or nonvocal\$ or non-vocal\$ or picture\$ or pictorial\$ or sign\$1 or signing or speech\$)).tw,id.
 61 (naturalistic adj2 (teach\$ or approach\$ or setting\$ or environment\$)).tw,id.
 62 focus?ed stimulation.tw,id.
 63 (milieu adj2 teaching).tw,id.
 64 (responsiv\$ adj2 education).tw,id.
 65 (responsiv\$ adj2 teaching).tw,id.
 66 Hanen\$.tw,id.
 67 or/57-66
 68 clinical trials/
 69 longitudinal studies/
 70 treatment effectiveness evaluation/
 71 exp program evaluation/
 72 (crossover or cross-over).tw.
 73 ((singl\$ or doubl\$ or tripl\$ or trebl\$) adj1 (blind\$ or mask\$)).tw.
 74 random\$.tw.
 75 assign\$.ab.
 76 allocat\$.ab.
 77 group\$.ab.
 78 or/68-77
 79 56 or 67
 80 29 and 79
 81 78 and 80

Science Citation Index Web of Science

#18#17 AND #16
 #17TS=(random* or control* or group* or placebo* or trial* or assign* or prospectiv* or meta-analys* or “systematic review” or longitudinal*)
 #16#15 OR #13
 #15#14 and #8
 #14TS= (“focus*ed stimulation” or “milieu teaching” or “responsive teaching” or “natural* teaching” or Hanen*)
 #13#12 AND #11
 #12TS=(coach* or educat* or intervention* or learn* or program* or teach* or train*)
 #11#10 AND #9 AND #8
 #10TS=(speech* or languag* or communicat* or sign or signing or nonverbal* or non-verbal* or nonvocal* or non-vocal* or cue* or symbol* or picture* or pictorial* or icon* or augment*)
 #9TS=(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*)

#8#7 AND #6

#7TS=(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten* or kinder-garten*)

#6#5 OR #4 OR #3 OR #2 OR #1

#5TS= ("acquired brain injur*" or ABI)

#4TS=(motor near/3 (disfunction* or dysfunction* or disorder* or disabilit* or impair*))

#3TS=(movement near/3 (disorder* or disabilit* or impair*))

#2ts=("non-progressive" Near/3 (disabilit* or disorder* or impair*))

#1TS= ("cerebral pals*" OR ataxic or ataxia or spastic* or dyskinetic or dyskinesia* or chorea* OR "down* syndrome" or "Angelman syndrome" or "happy puppet")

Conference Proceedings Citation Index - Science Web of Science (CPCI-S)

#18#17 AND #16

#17TS=(random* or control* or group* or placebo* or trial* or assign* or prospectiv* or meta-analys* or "systematic review" or longitudinal*)

#16#15 OR #13

#15#14 and #8

#14TS= ("focus*ed stimulation" or "milieu teaching" or "responsive teaching" or "natural* teaching" or Hanen*)

#13#12 AND #11

#12TS=(coach* or educat* or intervention* or learn* or program* or teach* or train*)

#11#10 AND #9 AND #8

#10TS=(speech* or languag* or communicat* or sign or signing or nonverbal* or non-verbal* or nonvocal* or non-vocal* or cue* or symbol* or picture* or pictorial* or icon* or augment*)

#9TS=(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*)

#8#7 AND #6

#7TS=(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten* or kinder-garten*)

#6#5 OR #4 OR #3 OR #2 OR #1

#5TS= ("acquired brain injur*" or ABI)

#4TS=(motor near/3 (disfunction* or dysfunction* or disorder* or disabilit* or impair*))

#3TS=(movement near/3 (disorder* or disabilit* or impair*))

#2ts=("non-progressive" Near/3 (disabilit* or disorder* or impair*))

#1TS= ("cerebral pals*" OR ataxic or ataxia or spastic* or dyskinetic or dyskinesia* or chorea* OR "down* syndrome" or "Angelman syndrome" or "happy puppet")

Language and Linguistic Behaviour Abstracts ProQuest (LLBA)

SU("Cerebral Palsy") OR SU("Developmental Disabilities") OR SU("Nervous System Disorders") OR TI,AB((motor OR movement OR non-progressive) N3 (disfunction* OR dysfunction* OR disorder* OR disabilit* OR impair*)) OR TI,AB("cerebral pals*" OR ataxic OR ataxia OR spastic* OR dyskinetic OR dyskenesia* OR chorea* OR "down* syndrome" OR "Angelman syndrome" OR "happy puppet") AND SU("Children") OR SU("Infants") OR TI,AB(child* OR infant* OR babies OR baby OR toddler* OR girl* OR boy* OR pre-school* OR preschool* OR nurser* OR kindergarten* OR kinder-garten*) AND SU("Parents") OR SU("Parent child interaction") OR TI,AB(parent* OR maternal* OR mother* OR father* OR paternal* OR carer* OR caregiver* OR care-giver* OR adult* OR teacher* OR therapist*) AND SU("Therapy") OR SU("Communicative competence") OR TI,AB(coach* OR educat* OR intervention* OR learn* OR program* OR teach* OR train* OR "focus*ed stimulation" OR "milieu teaching" OR "responsive teaching" OR "natural* teaching" OR "Hanen")

British Education Index EBSCOhost (BEI)

S37S33 AND S36

S36S34 OR S35

S35(random* or trial* or group* or assign* or allocat* or experiment* or control or PROSPECTIVE* OR longitudinal or BLIND* or CONTROL*)
 S34DE "RESEARCH" OR DE "COMMUNICATION -- Research" OR DE "COMPARATIVE studies" OR DE "EXPERIMENTS" OR DE "LANGUAGE & languages -- Research"
 S33S23 AND S32
 S32S24 OR S25 OR S26 OR S27 OR S28 OR S29 OR S30 OR S31
 S31(speech* or languag* or communicat* or sign or signing or nonverbal* or non-verbal* or nonvocal* or non-vocal* or cue* or symbol* or picture* or pictorial* or icon* or augment*)
 S30DE "VISUAL communication"
 S29DE "COMMUNICATION"
 S28DE "COMMUNICATION devices for people with disabilities"
 S27(DE "LANGUAGE disorders")
 S26DE "SPEECH disorders"
 S25(DE "SPEECH therapy")
 S24(DE "EDUCATION") OR (DE "TEACHING")
 S23S18 AND S22
 S22S19 OR S20 OR S21
 S21TI(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*) or AB(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*)
 S20DE "CAREGIVERS"
 S19DE "PARENTS" OR DE "FATHERS" OR DE "FOSTER parents" OR DE "GRANDPARENTS" OR DE "GRANDPARENTS as parents" OR DE "MOTHERS" " OR DE "SINGLE parents" OR DE "TEENAGE parents" OR DE "WORKING parents" OR DE "FAMILIES"
 S18S14 AND S17
 S17S15 OR S16
 S16TI(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten* or kindergarten*) OR AB(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten* or kinder-garten*)
 S15DE "CHILDREN" OR DE "PRESCHOOL children"
 S14S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13
 S13TI(PRRT2 or PRRT-2) OR AB(PRRT2 or PRRT-2)
 S12TI(GLUT1 or GLUT-1) OR AB(GLUT1 or GLUT-1)
 S11TI((Angelman* OR HAPPY PUPPET) n1 syndrome*) OR AB((Angelman* OR HAPPY PUPPET) n1 syndrome*)
 S10DE "DOWN syndrome"
 S9DE "DEVELOPMENTAL disabilities"
 S8TI (developmental* N3 (delay* or disab*)) OR AB (developmental* N3 (delay* or disab*))
 S7TI(acquired brain injur* or ABI) OR AB (acquired brain injur* or ABI)
 S6TI (motor N3 (disfunction* or dysfunction* or disorder* or disabilit* or impair*)) OR AB (motor N3 (disfunction* or dysfunction* or disorder* or disabilit* or impair*))
 S5TI (movement N3 (disorder* or disabilit* or impair*)) OR AB (movement N3 (disorder* or disabilit* or impair*))
 S4DE "MOTOR ability" OR DE "FINE motor ability" OR DE "GROSS motor ability" OR DE "MOTOR ability in children"
 S3(non-progressive N3 (disabilit* or disorder* or impair*))
 S2(ataxic or ataxia* or dyskinesia* or chorea* or spastic*)
 S1DE "MOVEMENT disorders in children" OR DE "CEREBRAL palsied children" or (cerebral pals*)

ERIC EBSCOhost (Education Resources Information Center)

S53S49 AND S52
 S52S50 OR S51
 S51TI (random* or trial* or experiment* or PROSPECTIVE* OR longitudinal or BLIND* or CONTROL*) OR AB (random* or trial* or experiment* or PROSPECTIVE* OR longitudinal or BLIND* or CONTROL*)
 S50DE "Meta Analysis" OR DE "Evaluation Research" OR DE "Control Groups" OR DE "Experimental Groups" OR DE "Longitudinal Studies" OR DE "Followup Studies" OR DE "Program Effectiveness" OR DE "Program Evaluation"

S49S23 AND S48
 S48S38 OR S39 OR S40 OR S41 OR S42 OR S43 OR S44 OR S45 OR S46 OR S47
 S47Hanen*
 S46(responsiv* N2 (education or teaching))
 S45(milieu N2 teaching)
 S44"focus*ed stimulation"
 S43(naturalistic N2 (teach* or approach* or setting* or environment*))
 S42TI((at home or (in N1 home) or home-based) N3 (coach* or educat* or intervention* or learn* or program* or teach* or train*))OR
 AB((at home or (in N1 home) or home-based) N3 (coach* or educat* or intervention* or learn* or program* or teach* or train*))
 S41TI(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or famil* or adult* or teacher* or
 therapist*) N3 (speech* or languag* or communicat* or sign or signing or nonverbal* or non-verbal* or nonvocal* or non-vocal* or
 cue* or symbol* or picture* or pictorial* or icon* or augment*) OR AB(parent* or maternal* or mother* or father* or paternal* or
 carer* or caregiver* or care-giver* or famil* or adult* or teacher* or therapist*) N3 (speech* or languag* or communicat* or sign or
 signing or nonverbal* or non-verbal* or nonvocal* or non-vocal* or cue* or symbol* or picture* or pictorial* or icon* or augment*)
 S40TI(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*)
 N3 (interact* or inter-act* or involv* or mediat* or respon*)OR AB(parent* or maternal* or mother* or father* or paternal* or carer*
 or caregiver* or care-giver* or adult* or teacher* or therapist*) N3 (interact* or inter- act* or involv* or mediat* or respon*)
 S39TI(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*)
 N3 (coach* or educat* or intervention* or learn* or program* or teach* or train*) OR AB(parent* or maternal* or mother* or father*
 or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*) N3 (coach* or educat* or intervention* or learn*
 or program* or teach* or train*)or TI(parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or
 adult* or teacher* or therapist*) N3 (coach* or educat* or intervention* or learn* or program* or teach* or train*)
 S38S26 AND S37
 S37S27 OR S28 OR S29 OR S30 OR S31 OR S32 OR S33 OR S34 OR S35 OR S36
 S36DE "Augmentative and Alternative Communication" OR DE "Nonverbal Communication" OR DE "Communication Skills" OR
 DE "Manual Communication" OR DE "Communication Strategies" OR DE "Communication Disorders"
 S35DE "Speech Impairments"
 S34DE "Language Impairments"
 S33DE "Speech Language Pathology"
 S32DE "Speech Therapy"
 S31DE "Special Education"
 S30DE "Early Intervention"
 S29DE "Parenting Skills"
 S28DE "Teaching (1966 1980)"
 S27DE "Education"
 S26S24 OR S25
 S25DE "Caregivers"
 S24DE "Parents" OR DE "Parents as Teachers"
 S23S19 AND S22
 S22S20 OR S21
 S21(DE "Young Children" OR DE "Infants" OR DE "Preschool Children" OR DE "Toddlers")
 S20TI(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten* or kinder-
 garten*) OR AB(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten*
 or kinder-garten*)
 S19S1 OR S2 OR S3 OR S4 OR S5 OR S6 OR S7 OR S8 OR S9 OR S10 OR S11 OR S12 OR S13 OR S14 OR S15 OR S16 OR
 S17 OR S18
 S18TI(PRRT2 or PRRT-2) OR AB(PRRT2 or PRRT-2)
 S17TI(GLUT1 or GLUT-1) OR AB(GLUT1 or GLUT-1)
 S16TI(Angelman* OR HAPPY PUPPET) n1 syndrome*) OR AB((Angelman* OR HAPPY PUPPET) n1 syndrome*)
 S15TI(dyskinesia* or chorea*) OR AB(dyskinesia* or chorea*)
 S14TI(down* syndrome*) OR AB(down* syndrome*)
 S13DE "Down Syndrome"
 S12DE "Developmental Disabilities"

S11TI (developmental* N3 (delay* or disab*)) OR AB (developmental* N3 (delay* or disab*))
 S10TI (acquired brain injur* or ABI) OR AB (acquired brain injur* or ABI)
 S9TI (motor N3 (disfunction* or dysfunction* or disorder* or disabilit* or impair*)) OR AB (motor N3 (disfunction* or dysfunction* or disorder* or disabilit* or impair*))
 S8TI (movement N3 (disorder* or disabilit* or impair*)) OR AB (movement N3 (disorder* or disabilit* or impair*))
 S7DE "Psychomotor Skills" OR DE "Motor Development"
 S6(non-progressive N3 (disabilit* or disorder* or impair*))
 S5TI(dyskinetic*) OR AB(dyskinetic*)
 S4TI(spastic*) or AB(spastic*)
 S3TI (ataxic or ataxia*) or AB(ataxic or ataxia*)
 S2TI(cerebral pals*) or AB(cerebral pals*)
 S1DE "Cerebral Palsy"

Cochrane Database of Systematic Reviews (CDSR) and Database of Abstracts of Reviews of Effects (DARE), part of the Cochrane Library

#1[mh ^"Cerebral Palsy"]
 #2cerebral pals*:ti,ab
 #3(non-progressive near/3 (disabilit* or disorder* or impair*)):ti,ab
 #4[mh ^"Movement Disorders"]
 #5[mh "motor disorders"]
 #6[mh ^"motor skills disorder"]
 #7[mh "Motor Skills"]
 #8(movement near/1 (disorder* or disabilit* or impair*)):ti,ab
 #9(motor near/1 (disfunction* or dysfunction* or disorder* or disabilit* or impair*)):ti,ab
 #10[mh ^"Brain Injuries"]
 #11[mh "Brain Damage, Chronic"]
 #12[mh "Brain injury, chronic"]
 #13("acquired brain injur*" or ABI):ti,ab
 #14[mh ^"Developmental disabilities"]
 #15(developmental* near/3 (delay* or disab*)):ti,ab
 #16[mh dyskinesias]
 #17[mh Chorea]
 #18(dyskinesia* or chorea*):ti,ab
 #19[mh "Angelman Syndrome"]
 #20((Angelman* or "Happy Puppet") near/1 syndrome*):ti,ab
 #21(GLUT1 or GLUT-1):ti,ab
 #22(PRRT2 or PRRT-2):ti,ab
 #23{or #1-#22}
 #24[mh Infant]
 #25[mh child]
 #26(child* or infant* or babies or baby or toddler* or girl* or boy* or pre-school* or preschool* or nurser* or kindergarten* or kindergarten*):ti
 #27{or #24-#26}
 #28#23 and #27
 #29[mh "Parent-Child Relations"]
 #30[mh Parents]
 #31[mh Parenting]
 #32[mh Family]
 #33[mh Caregivers]
 #34{or #29-#33}
 #35[mh education]
 #36[mh teaching]

#37[mh "Early intervention (Education)"]
 #38[mh "Education of Intellectually Disabled"]
 #39[mh "education, special"]
 #40[mh "speech therapy"]
 #41[mh "language therapy"]
 #42[mh "speech-language pathology"]
 #43[mh "speech disorders"]
 #44[mh "language disorders"]
 #45[mh "Sign language"]
 #46[mh "Manual Communication"]
 #47[mh "Nonverbal communication"]
 #48[mh "Communication Aids for Disabled"]
 #49[mh "Self-Help Devices"]
 #50[mh "communication"]
 #51[mh "communication disorders"]
 #52{or #35-#51}
 #53#34 and #52
 #54((parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*)
 near/3 (coach* or educat* or intervention* or learn* or program* or teach* or train*)):ti,ab
 #55((parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or adult* or teacher* or therapist*)
 near/3 (interact* or inter-act* or involv* or mediat* or respon*)):ti,ab
 #56((parent* or maternal* or mother* or father* or paternal* or carer* or caregiver* or care-giver* or famil* or adult* or teacher* or
 therapist*) near/3 (speech* or languag* or communicat* or sign or signing or nonverbal* or non-verbal* or nonvocal* or non-vocal* or
 cue* or symbol* or picture* or pictorial* or icon* or augment*)):ti,ab
 #57(naturalistic near/2 (teach* or approach* or setting* or environment*)):ti,ab
 #58"focus*ed stimulation":ti,ab
 #59(milieu near/2 teaching):ti,ab
 #60(responsiv* near/2 education):ti,ab
 #61(responsiv* near/2 teaching):ti,ab
 #62Hanan*:ti,ab
 #63{or #53-#62}
 #64#28 and #63 in Cochrane Reviews (Reviews and Protocols) and Other Reviews

LILACS (Latin American and Caribbean Health Science Information database; lilacs.bvsalud.org/en)

(tw:(((motor OR movement))) AND (tw:(disorder* OR dysfunction* OR disfunction* OR impair* OR disabil*))) OR (tw:("cerebral
 pals*" OR ataxic OR ataxia* OR dyskinetic* OR dyskinesia* OR chorea* OR spastic*)) AND (tw:((speech* OR languag* OR com-
 municat* OR sign OR signing OR nonverbal* OR non-verbal* OR nonvocal* OR non-vocal* OR cue* OR symbol* OR picture* OR
 pictorial* OR icon* OR augment*))) AND (instance:"regional") AND (db:("LILACS") AND type`of`study:("clinical`trials"))

Rehabdata National Rehabilitation Information Center (www.naric.com/?q=REHABDATA)

3 separate search strings
 WITH ALL OF THESE WORDS PARENTS
 WITH ANY OF THESE WORDS Movement OR motor OR cerebral
 WITH THE EXACT PHRASE RANDOM*

SpeechBITE (speechbite.com)

Keyword(s): "cerebral palsy"
 Age group: Children
 Research Design: Randomised Controlled Trial
 Keyword(s): "movement disorders"

Age group: Children
Research Design: Randomised Controlled Trial
Keyword(s): "motor disorders"
Age group: Children
Research Design: Randomised Controlled Trial

ClinicalTrials.gov (clinicaltrials.gov)

Interventional Studies | cerebral palsy OR movement OR motor OR spastic OR Angelman OR Down Syndrome OR ataxic OR ataxia OR chorea OR dyskinesia | communication OR speech OR language OR Hanen OR naturalistic OR responsive OR milieu OR augmentative | Child

UK Clinical Trials Gateway (www.ukctg.nihr.ac.uk/home)

Search 1 (motor disorders OR movement disorders OR cerebral palsy) AND (speech OR language OR communication)
Search 2 (motor disorders OR movement disorders OR cerebral palsy) AND (parents OR caregivers OR carers)

World Health Organisation International Clinical Trials Registry Platform (WHO ICTRP; www.who.int/ictRP/en)

CONDITION: cerebral palsy OR movement OR motor OR spastic OR Angelman OR Down Syndrome OR ataxic OR ataxia OR chorea OR dyskinesia
INTERVENTION: communication OR speech OR language OR Hanen OR naturalistic OR responsive OR milieu
Limited to Clinical Trials in Children

Appendix 2. Criteria for judging risk of bias

Random sequence generation (method of assigning participants to groups)

1. Low risk of bias: well-described, randomised process (e.g. coin toss; table of random numbers; computerised, random number generator).
2. High risk of bias: non-random method (e.g. days of the week, alternate).
3. Unclear risk of bias: allocation was not described or description led to uncertainty in quality of allocation, and possibility of bias.

Allocation concealment

1. Low risk of bias: we rated concealment of allocation as adequate if random allocation schedules were developed by an independent researcher, and if allocation was recorded within opaque envelopes created by the independent researcher or the computerised allocation system was controlled by the independent researcher.
2. High risk of bias: providers of intervention undertook allocation or research team allocated participants and had access to participant characteristics.
3. Unclear risk of bias: methods of concealment were not described or description did not allow bias to be ruled out.

Blinding of participants and personnel

In the case of parent-mediated training interventions, neither the parent nor the trainer can be blinded to the type of treatment given. Blinding in studies in this review, therefore, referred to components of the intervention being tested (e.g. video coaching; focus on an individual communication strategy). We assessed studies on an individual basis; however, most studies were at high risk of bias on this criterion due to the nature of the intervention.

1. Low risk of bias: parents and trainers were blinded to which of the two comparison treatments they were allocated or were blinded to the exact feature of the intervention that was being tested.

2. High risk of bias: parents or trainers, or both, were not blinded to the exact focus of intervention being tested, or it was likely that blinding could have been broken during the study.
3. Unclear risk of bias: insufficient information provided to judge the knowledge of parents and trainers on the feature of the intervention being tested or the difference between individual treatments.

Blinding of outcome assessors

1. Low risk of bias: reports stated that outcome assessors were blinded to group allocation.
2. High risk of bias: reports suggested that assessors were likely to have known the group to which the participant was allocated (e.g. provided treatment, worked with person delivering treatment).
3. Unclear risk of bias: no information on blinding of assessors or role of assessors in allocation or treatment provision.

Completeness of data collection

We reported the numbers in each intervention group at each end point compared with total number of participants randomised and the reason(s) for attrition/exclusion where provided by study authors. If missing data were imputed, we consulted a statistician about the appropriateness of the method used. Had we retrieved and entered data into the review, we would have reported such reinclusions in analyses.

1. Low risk of bias: no missing outcome data; or loss to follow-up was unlikely to be related to the true outcome; or attrition was similar in both conditions and proportion of missing data (dichotomous outcomes) or effect size (continuous outcomes) were unlikely to have a clinically-relevant effect; imputation of missing data judged to be appropriate.
2. High risk of bias: loss of participants to follow-up was likely to be related to the true outcome, or was distributed unevenly across groups; or we considered imputation of missing data to be inappropriate. Had we encountered studies showing uneven loss to follow-up, we would have considered these separately in sensitivity analyses.
3. Unclear risk of bias: loss of participants to follow-up was not reported or insufficient information was provided to judge the reason for loss and a judgement of low or high risk of bias.

Selective reporting

We checked published protocols to assess if all planned comparisons were reported. If protocols were not published, we contacted study authors to ask if additional comparisons were planned.

1. Low risk of bias: all prespecified outcomes were reported.
2. High risk of bias: selective reporting of outcomes was evident.
3. Unclear risk of bias: not possible to judge if all planned comparisons were reported.

Other sources of bias

We described any additional problems that might have put a study at risk of bias.

1. Low risk of bias: study appeared free from other sources of bias.
2. High risk of bias: at least one important risk of bias (e.g. groups clearly different at baseline, children receiving direct therapy during the study period).
3. Unclear risk of bias: insufficient information provided on which to judge other sources of bias.

We judged studies that scored a low risk of bias for all criteria at low risk of bias overall. Where studies were deemed at low or unclear risk of bias for all criteria, we judged them as being at unclear risk of bias overall. Where studies were deemed at high risk of bias for one or more criteria, we judged them at high risk of bias overall.

CONTRIBUTIONS OF AUTHORS

LP and JG conceived and designed the review.

LP and WA screened titles and abstracts. JG was involved in disagreement resolution.

All authors considered full-text reports for inclusion in the review, extracted data from the included studies, and assessed the risk of bias and quality of the evidence. In the event of disagreement regarding inclusion, a third review author (WA, JG, KL or LP) reviewed the report independently, and we reached consensus through discussion and by reassessing the study against the inclusion criteria together.

LP and WA wrote the review with drafts reviewed by KL and JG.

LP has overall responsibility for the review.

DECLARATIONS OF INTEREST

LP is a Speech and Language Therapist and Senior Lecturer at the Institute of Health and Society, Newcastle University, UK. LP has led an early phase trial of parent-mediated intervention for children with cerebral palsy. LP's institution received funding from the National Institute for Health Research (NIHR) Cochrane Incentive Awards to complete this review and from Sparks, The Children's Medical Research Charity, to develop a smart phone app for use in parent-mediated therapy (14NCL01). The latter study does not fit the criteria for this review, as assessed by KL and JG, and is not included. LP received payment of accommodation fees by Cerebral Palsy Alliance to present at a symposium on Early Intervention at the International Cerebral Palsy Conference, 31 May to 1 June 2016, Stockholm, Sweden.

WA is currently a PhD Student at Newcastle University.

KL is an Advanced Speech and Language Therapist, at the Northumberland Tyne and Wear NHS Foundation Trust, UK. KL's previous employer (City Hospitals Sunderland) received a set fee to release KL for 30 days (January to June 2016), as an internship with Dr Pennington (Health Education North and NIHR Clinical Academic Training Programme).

JG is Faculty Head of Research and Knowledge Exchange (Health, Psychology and Social Care Faculty) at Manchester Metropolitan University, UK. JG currently receives funding (14/70/153) from the NIHR Health Services and Delivery Research (HS&DR) Programme, to conduct the following research: "Identifying appropriate symbol communication aids for children who are non-speaking: enhancing clinical decision making." The project, which commenced in February 2016 and will run for three years, addresses the tangentially-related research area of clinical decision-making in augmentative and alternative communication. JG's role is primarily in systematic reviews to inform the project and, as such, JG does not perceive any conflict of interest. JG also declares that she received funding for external evaluation of the 'ENABLIN+' project, aimed at staff training for working with children with complex and intense support needs, Contract 541981-LLP-1-2013-1-BE-LEONARDO-LMP, and that this does not bring any conflict of interest. JG's institution was paid by the International Society for Augmentative and Alternative Communication, Norway, for two lectures on early communication and evidence-based practice in the spring of 2016. JG receives royalties for *Coupe-O'Kane and Goldbart (1997) "Communication Before Speech."* JG declares the book focuses on children with profound disabilities rather than motor disorders and addresses assessment and generic strategies rather than specific interventions.

SOURCES OF SUPPORT

Internal sources

- Newcastle University, UK.
Salary for Lindsay Pennington
- Manchester Metropolitan University, UK.
Salary for Juliet Goldbart

External sources

- National Institute for Health Research (NIHR), UK.
- This review presents independent research funded by Health Education England (HEE) and the National Institute for Health Research (NIHR) under the Clinical Academic Training Programme; Internship Awards 2015/16. The completion of the review was supported also by a £5000 grant from the NIHR Cochrane Incentive Scheme 2017. The views expressed herein are those of the authors and not necessarily those of the National Health Service (NHS), HEE, the NIHR or the Department of Health.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

1. Types of participants

i) We included only studies that had a least one child with motor disorders in each group whose motor disorders may have influenced their communication, to increase similarity between groups in individual studies.

2. Electronic searches

i) We did not search EU Clinical Trials Register (www.clinicaltrialsregister.eu/ctr-search/search) because it does not register psychotherapeutic trials. We were not able to search NIH Clinical Research Center (www.cc.nih.gov/home/clinicalstudies.html) because the website returned the message "Site unavailable."

3. Summary of findings for the main comparison

i) We produced one 'Summary of findings' table for our main comparison, rather than one table for outcome, as stated in our protocol ([Pennington 2017](#))