IMMUNOMETABOLISM

Metabolic requirements to expand and arm a clone army

Andrew L. Mellor and Lei Huang (Institute of Cellular Medicine, Newcastle University)

A new target to control T cell responses involves de novo biosynthesis of tetrahydrobiopterin (BH4), adding to the list of processes dependent on BH4 synthesis, which includes neurotransmitter and nitric oxide production and pain control.

T cells are the storm troopers of the immune system since they can kill other cells. Consequently, T cells must be licensed to expand and bear arms through a tightly regulated process ¹. A recent study in *Nature* by Cronin et al. reveal a critical role for the enzyme cofactor tetrahydrobiopterin (BH4) in licensing T cell clones to proliferate and acquire weapons of destruction ². These findings add to the extensive list of biological functions mediated by BH4, including production of neurotransmitters, nitric oxide and pain control.

Circulating T cells are metabolically inert. On activation, T cells proliferate rapidly and acquire effector functions, including, in the case of cytotoxic T cells, the ability to destroy target cells (Fig. 1). This licensing process has strict requirements for coordinated delivery of activation signals from antigens, co-stimulatory ligands and pro-inflammatory cytokines ¹. These constraints ensure that potentially dangerous effector T cells are generated only as needed. Cronin et al. report that de novo, cell-intrinsic, BH4 synthesis is necessary for T cells to undergo clonal expansion. The initial prompt was the observation that activated T cells expressed GTP cyclohydrolase I (GCH1), the first step in BH4 synthesis ². *Gch1*-deficient mice were generated and activated T cells from these mice exhibited profound proliferative defects in vitro and in vivo,

confirming that T cell-intrinsic BH4 synthesis was required for efficient T cell licensing. This proliferative defect was highly selective for T cells, as *Gch1* ablation did not impact development of T and B lymphocytes or the ability of B cells and regulatory T cells (T_{reg} cells) to acquire effector functions on activation.

The potential clinical significance of BH4 emerged from mouse models of inflammatory colitis, allergy and autoimmune syndromes. In each model, inflammatory infiltrates and disease severity were attenuated significantly due to defective clonal expansion of T cells lacking *Gch1*. However, selective *Gch1* ablation did not impair the ability of T cells to make IL-17 and IFN- γ , two pro-inflammatory cytokines that mediate T cell effector functions and drive auto-inflammatory diseases. It was not reported if *Gch1* ablation impacted expression of perforin and granzyme, which mediate target cell killing. Intriguingly, experimental autoimmune encephalitis (EAE), a popular mouse model of multiple sclerosis (MS), followed normal kinetics for two weeks after disease induction, irrespective of the ability of T cells to make BH4. Thus, the ability to make BH4 was not essential to drive EAE onset or to maintain disease for 6 weeks, implying that effector T cells do not induce or maintain EAE. However, intrinsic BH4 synthesis by T cells was required for EAE severity to increase progressively after disease onset. It is unclear from the data presented if a delayed role for BH4 synthesis in T cells also manifested in the colitis and allergy models. Collectively, these findings reveal that BH4 synthesis is necessary for activated T cells to expand but BH4 is not required for T cells to acquire weapons of destruction.

The finding that T cells must make BH4 to undergo clonal expansion identifies a potential 'Achilles' heel' that might be exploited for clinical benefit. Accordingly, the study authors show that inhibiting BH4 synthesis attenuated T cell responses, while boosting BH4 synthesis potentiated T cell responses. A new drug (QM385) was developed to inhibit sepiapterin reductase (SPR), the terminal enzyme in the BH4 synthesis pathway. As expected, QM385 reduced clonal expansion of activated mouse and human T cells and attenuated inflammatory

responses and disease severity in the allergy model. On the other hand, enhancing *Gch1* expression increased T cell clonal expansion and increased T cell-mediated inflammation and disease severity in the allergy and colitis models, consistent with the notion that BH4 synthesis boosts destructive T cell immunity in these models. Enhanced *Gch1* expression in mice also conferred greater resistance to tumor growth and administering BH4 to mice bearing established (albeit small) tumors attenuated subsequent tumor growth. These studies identify the BH4 synthesis pathway as a promising target to manipulate T cell immunity for clinical benefit in a range of diseases.

Further insights into the molecular mechanism of BH4-mediated licensing of T cells emerged from gene expression profiling of activated *Gch1*-deficient T cells. Enzymes involved in iron metabolism were affected by *Gch1* ablation. Mitoferrin, a mitochondrial iron transporter, was up-regulated prominently in T cells lacking *Gch1*, whereas iron levels were reduced in activated T cells lacking intact *Gch1* genes. These findings implied that *Gch1* mobilizes iron required for normal mitochondrial respiration, a notion borne out by studies showing that ATP synthesis was attenuated and glycolysis was augmented in T cells from *Gch1*-deficient mice. Moreover, mitochondrial respiration and oxygen consumption were lower in T cells unable to synthesize BH4 due to defective mitochondrial electron transport linked to imbalance in cytochrome c redox states and consequent elevated production of reactive oxygen species (ROS). These defects were corrected by adding BH4 or reduced cytochrome c to *Gch1*-deficient T cells. Taken together, these studies suggest that BH4 maintains iron homeostasis needed for mitochondrial respiration in activated T cells.

In an intriguing aside Cronin et al. mention a link between BH4 synthesis and tryptophan (Trp) metabolism, which impacts immunologic and neurologic processes. Trp catabolism suppresses T cell immunity via the kynurenine (Kyn) pathway, which is initiated by indoleamine 2,3 dioxygenase (IDO) and tryptophan 2,3 dioxygenase (TDO). Elevated IDO and TDO activity

suppresses T cell immunity in a range of chronic inflammatory diseases, including cancers, allergic and autoimmune syndromes ^{3,4}. Cells expressing IDO or TDO produce bioactive Trp catabolites that suppress immunity and cause neurologic dysfunction. For example, Kyn and other downstream Trp catabolites suppress T cell immunity, though molecular mechanisms of T cell suppression remain controversial ³. Cronin et al. report that Kyn treatment inhibited SPR activity in T cells and that adding BH4 overcame the T cell suppressive effects of adding Kyn to restore T cell proliferation and reduce ROS production. Xanthurenic acid (XA) is a Trp catabolite made downstream of Kyn and XA is a potent SPR inhibitor ⁵, prompting the suggestion that XA might mediate the T cell suppressive effects of Kyn (Fig. 1). If verified, the Kyn pathway might be a natural regulator of BH4 synthesis in T cells, offering a potential new mechanism to explain the T cell suppressive effects of IDO and TDO activity. These findings also imply that the T cell stimulatory effects of IDO inhibitors, currently under intense scrutiny as potential drugs to boost anti-tumor immunity in cancer patients ⁶, may in part arise due to blockade of XA synthesis, thereby releasing the brakes on BH4-mediated clonal expansion of effector T cells (Fig. 1).

Though potential links between the BH4 and Kyn pathways cited above are plausible they raise new questions. For example, access to amino acids is essential for T cell licensing, as amino acid transporters are induced in activated T cell and loss of amino acid transporter functions blocks T cell clonal expansion and impairs development of malignant T cell leukemias ^{7,8}. In particular, T cells are sensitive to Trp depletion since the ribosomal amino acid sensors GCN2 and mTOR block T cell clonal expansion if IDO and TDO are active ³. Tryptophan hydroxylase-1 (TPH1) also consumes Trp as the obligate substrate to make the monoamine neurotransmitter serotonin (5-hydroxytryptamine, 5HT). Intriguingly, BH4 is an essential TPH1 co-factor and TPH1 activity was necessary for mast cells to suppress T cell immunity in mouse models of transplant tolerance, tumor growth and EAE via a mechanism independent of 5HT synthesis ⁹. These observations beg the question of whether de novo BH4 biosynthesis in mast

cells might be necessary to suppress T cell immunity that protects tumors, or transplanted and healthy tissues from destruction in these models. Other questions relate to links between elevated Kyn pathway activity during chronic inflammation and neurologic dysfunction leading to increased pain sensitivity, chronic fatigue and neurologic depression ^{10,11}. Clearly, the BH4, 5HT and Kyn pathways are a confluence of metabolic processes that regulate immunologic and neurologic processes in many chronic inflammatory conditions, which may be exploited for therapeutic benefit. As evidenced by ongoing efforts to develop IDO and TDO inhibitors as treatments to block Trp catabolism in cancer patients⁶, metabolic pathways underpinning immune processes provide a growing number of new targets for immunotherapy. It is also likely that common neurologic comorbidities arising during disease progression or in response to therapy may be rooted in metabolic changes in BH4, 5HT and Kyn pathway activity.

In summary, Cronin et al. identify BH4 synthesis as a metabolic ‘Achilles’ heel’ that might be exploited to prevent T cells from acquiring storm trooper status as they transit through the immunologic licensing process. Their study also raises intriguing links between BH4 and 5HT synthesis and the Kyn pathway that might have wide ranging implications for understanding how immune balance and neurologic comorbidities manifest in many chronic inflammatory diseases and how immunotherapies modify these linked metabolic pathways. In light of these reminders of the critical importance of metabolic controls it may be worth remembering Napoleon’s alleged quote that, ‘an army marches on its stomach’.

Competing interests

ALM is a scientific consultant for NewLink Genetics and receives income and share options for this service. LH declares no competing interests.

Figure 1. BH4 plays a key role in T cell licensing. Cronin et al., report that de novo BH4 synthesis is essential for activated T cells to expand and acquire effector functions. BH4 regulates iron metabolism needed for mitochondrial (Mito) functions that support T cell licensing. Indoleamine 2,3 dioxygenase (IDO) activity inhibits BH4 synthesis by generating the tryptophan (Trp) catabolite xanthurenic acid (XA), which inhibits sepiapterin reductase (SPR) activity to block T cell licensing. BH4 is also a cofactor for TPH, the first step in serotonin (5HT) synthesis.

References

- 1 Malissen, B. & Bongrand, P. Early T cell activation: integrating biochemical, structural, and biophysical cues. *Annual review of immunology* **33**, 539-561, doi:10.1146/annurev-immunol-032414-112158 (2015).
- 2 Cronin, S. J. F. *et al.* The metabolite BH4 controls T cell proliferation in autoimmunity and cancer. *Nature*, doi:10.1038/s41586-018-0701-2 (2018).
- 3 Munn, D. H. & Mellor, A. L. Indoleamine 2,3 dioxygenase and metabolic control of immune responses. *Trends in immunology* **34**, 137-143, doi:10.1016/j.it.2012.10.001 (2012).
- 4 Platten, M., von Knebel Doeberitz, N., Oezen, I., Wick, W. & Ochs, K. Cancer Immunotherapy by Targeting IDO1/TDO and Their Downstream Effectors. *Frontiers in immunology* **5**, 673, doi:10.3389/fimmu.2014.00673 (2014).
- 5 Haruki, H., Hovius, R., Pedersen, M. G. & Johnsson, K. Tetrahydrobiopterin Biosynthesis as a Potential Target of the Kynurenine Pathway Metabolite Xanthurenic Acid. *J Biol Chem* **291**, 652-657, doi:10.1074/jbc.C115.680488 (2016).

- 6 Muller, A. J., Manfredi, M. G., Zakharia, Y. & Prendergast, G. C. Inhibiting IDO pathways to treat cancer: lessons from the ECHO-301 trial and beyond. *Semin Immunopathol*, doi:10.1007/s00281-018-0702-0 (2018).
- 7 Sinclair, L. V. *et al.* Control of amino-acid transport by antigen receptors coordinates the metabolic reprogramming essential for T cell differentiation. *Nat Immunol* **14**, 500-508, doi:10.1038/ni.2556 (2013).
- 8 Grzes, K. M. *et al.* Control of amino acid transport coordinates metabolic reprogramming in T-cell malignancy. *Leukemia*, doi:10.1038/leu.2017.160 (2017).
- 9 Nowak, E. C. *et al.* Tryptophan hydroxylase-1 regulates immune tolerance and inflammation. *J Exp Med* **209**, 2127-2135, doi:10.1084/jem.20120408 (2012).
- 10 Kim, H. *et al.* Brain indoleamine 2,3-dioxygenase contributes to the comorbidity of pain and depression. *J Clin Invest* **122**, 2940-2954, doi:10.1172/JCI61884 (2012).
- 11 Huang, L. *et al.* Virus Infections Incite Pain Hypersensitivity by Inducing Indoleamine 2,3 Dioxygenase. *PLoS Pathog* **12**, e1005615, doi:10.1371/journal.ppat.1005615 (2016).